
NPTE MUSCULOSKELETAL (MSK) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://nptemsk.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **Heberden's nodes are commonly associated with which type of arthritis?**
   - A. Rheumatoid arthritis
   - B. Osteoarthritis
   - C. Gout
   - D. Lupus arthritis

2. **What distinguishes the Biceps Load Test I from Biceps Load Test II?**
   - A. Angle of shoulder abduction
   - B. Position of the forearm
   - C. Degree of arm rotation
   - D. Type of resistance applied

3. **Which hip position is associated with the transition from heel strike to foot flat?**
   - A. Flexion and lateral rotation
   - B. Extension and medial rotation
   - C. Flexion and medial rotation
   - D. Extension and neutral

4. **What does flat foot in old gait terminology occur during?**
   - A. Midstance
   - B. Initial contact
   - C. Loading response
   - D. Heel off

5. **What condition might involve fibrous changes in the internal organs?**
   - A. Systemic lupus erythematosus
   - B. Progressive systemic sclerosis
   - C. Osteoarthritis
   - D. Fibromyalgia

SAMPLE

6. What is a common exercise recommendation for patients with spondylolisthesis?

A. High-impact aerobics

B. Core stabilization exercises

C. Heavy weightlifting

D. Extension exercises

7. What is a characteristic of the S1 nerve root?

A. Hip flexion

B. Foot eversion

C. Knee extension

D. Hip adduction

8. During the standing flexion test for the lumbar spine, what indicates a positive test?

A. No movement of both PSIS

B. The PSIS that moves inferiorly first

C. The PSIS that moves cranially first

D. Symmetry in PSIS movement

9. Finkelstein's test primarily evaluates which tendons?

A. Flexor pollicis longus and flexor carpi ulnaris

B. Abductor pollicis longus and extensor pollicis brevis

C. Extensor pollicis longus and extensor carpi radialis

D. Extensor digitorum and abductor digiti minimi

10. What are the primary symptoms of juvenile rheumatoid arthritis?

A. Joint stiffness and swelling

B. Fever and rash

C. Severe pain and immobility

D. Muscle weakness and discomfort

SAMPLE

Answers

SAMPLE

1. B
2. A
3. D
4. C
5. B
6. B
7. B
8. C
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. Heberden's nodes are commonly associated with which type of arthritis?

A. Rheumatoid arthritis

B. Osteoarthritis

C. Gout

D. Lupus arthritis

Heberden's nodes are bony swellings that occur at the distal interphalangeal (DIP) joints of the fingers and are a classic sign associated with osteoarthritis. This condition is characterized by the degeneration of cartilage and the formation of bone spurs, leading to joint pain and stiffness. The presence of Heberden's nodes indicates that there has been wear and tear on the joints, which is typical in osteoarthritis. While rheumatoid arthritis and other types of arthritis can cause joint swelling and changes, Heberden's nodes specifically are not found in these conditions. In rheumatoid arthritis, for example, the joint involvement tends to affect the proximal interphalangeal (PIP) joints more prominently and is characterized by systemic symptoms and symmetrical joint involvement. Similarly, in gout and lupus arthritis, the patterns of joint involvement differ significantly from that seen in osteoarthritis. Thus, the association of Heberden's nodes with osteoarthritis highlights the unique pathophysiology of this degenerative joint disease.

2. What distinguishes the Biceps Load Test I from Biceps Load Test II?

A. Angle of shoulder abduction

B. Position of the forearm

C. Degree of arm rotation

D. Type of resistance applied

The distinguishing feature between Biceps Load Test I and Biceps Load Test II is indeed the angle of shoulder abduction. In Biceps Load Test I, the shoulder is positioned at 90 degrees of abduction, while in Biceps Load Test II, the shoulder is positioned at 120 degrees of abduction. This variation in degrees affects the tension in the biceps tendon and the structures surrounding the shoulder, specifically the labrum. The purpose of these tests is to assess the integrity of the biceps tendon and its association with the labrum, particularly in cases of SLAP (Superior Labrum Anterior to Posterior) lesions. By changing the abduction angle, practitioners can create different biomechanical loads on the biceps tendon and surrounding soft tissues, helping to differentiate between potential pathologies. This procedural variation is essential for accurate assessment and diagnosis in patients with shoulder complaints.

SAMPLE

3. Which hip position is associated with the transition from heel strike to foot flat?

A. Flexion and lateral rotation

B. Extension and medial rotation

C. Flexion and medial rotation

D. Extension and neutral

The transition from heel strike to foot flat during walking involves specific movements of the hip that facilitate smooth weight transfer and the deceleration of the leg. When considering this phase of gait, the hip is primarily in an extended position, which allows for optimal stability and alignment as the body weight shifts onto the forward leg after initial contact with the ground. At the moment of heel strike, the knee is typically flexed to absorb the shock, while the hip is in a neutral or slightly extended position. This alignment helps maintain a stable center of mass and prepares the lower limb for the subsequent loading response. Extension of the hip at this stage is crucial as it counters the forward momentum and stabilizes the pelvis, ensuring a controlled transition as the weight shifts toward the foot flat position. In contrast, the other options imply varying combinations of flexion and rotation that do not align with the mechanics of hip function during this specific phase of walking. These positions would compromise stability and the effective loading of the limb, making them less suited for the heel strike to foot flat transition.

4. What does flat foot in old gait terminology occur during?

A. Midstance

B. Initial contact

C. Loading response

D. Heel off

Flat foot, in the context of gait terminology, refers to a period in the gait cycle where the foot comes into full contact with the ground. This typically occurs during the loading response phase of the gait cycle. During the loading response, the foot adopts a flat position to accommodate weight transfer from the heel to the forefoot. This phase starts immediately after the foot makes initial contact with the ground. As the body weight shifts, the entire plantar surface of the foot engages with the ground, facilitating balance and shock absorption. It's important to understand this concept within the context of the gait cycle. While midstance refers to the phase when the body is directly over the weight-bearing foot, and initial contact occurs as the heel strikes the ground, flat foot specifically relates to the loading response where the foot has settled and is effectively flat against the ground. This position is critical for providing support and stability while distributing the forces exerted during walking.

SAMPLE

5. What condition might involve fibrous changes in the internal organs?

A. Systemic lupus erythematosus

B. Progressive systemic sclerosis

C. Osteoarthritis

D. Fibromyalgia

Progressive systemic sclerosis, also known as scleroderma, is characterized by the abnormal accumulation of collagen leading to fibrous changes in the skin and internal organs. This condition involves a cascade of pathological processes where fibrosis can develop within various organ systems, including the heart, lungs, kidneys, and gastrointestinal tract. As a result, individuals may experience a range of symptoms such as skin tightening, pulmonary hypertension, renal impairment, and difficulty with digestion. The fibrotic changes are a direct consequence of the autoimmune nature of progressive systemic sclerosis, where the immune system erroneously attacks the body's own tissues, resulting in excessive collagen deposition and organ dysfunction. This distinguishes it from conditions like systemic lupus erythematosus, which primarily affects the skin, joints, and multiple organs but does not typically cause the level of fibrosis seen in scleroderma. In other conditions mentioned, osteoarthritis primarily affects joints without causing systemic fibrotic changes, while fibromyalgia is characterized by widespread musculoskeletal pain without the fibrotic response or involvement of internal organs. Hence, progressive systemic sclerosis is the condition that directly correlates with fibrous changes in internal organs.

6. What is a common exercise recommendation for patients with spondylolisthesis?

A. High-impact aerobics

B. Core stabilization exercises

C. Heavy weightlifting

D. Extension exercises

Core stabilization exercises are commonly recommended for patients with spondylolisthesis because they help enhance the stability of the lumbar spine and support the affected vertebrae. This condition involves the forward slippage of one vertebra over another, which can lead to instability and discomfort. By strengthening the core muscles, including the abdominal, back, and pelvic floor muscles, these exercises provide crucial support to the spine, helping to reduce excessive motion and alleviate pressure on the affected area. In addition, core stabilization exercises can improve overall posture and biomechanics, which can be beneficial in managing symptoms and preventing further progression of the condition. These exercises typically focus on promoting controlled movements and endurance rather than high-impact activities or heavy lifting, which could exacerbate the instability and pain associated with spondylolisthesis. The other options, such as high-impact aerobics, heavy weightlifting, or extension exercises, may not be suited for patients with this condition. High-impact activities can place undue stress on the spine, while heavy lifting could lead to further injury. Extension exercises might also worsen symptoms for some patients, particularly if they lead to increased extension of the lumbar spine, which can exacerbate pain in cases of spondylolisthesis, especially if there is associated

SAMPLE

7. What is a characteristic of the S1 nerve root?

A. Hip flexion

B. Foot eversion

C. Knee extension

D. Hip adduction

The S1 nerve root is primarily associated with the common peroneal nerve, which has a significant role in foot eversion and plantarflexion. This nerve innervates the muscles responsible for lifting the outer edge of the foot, allowing for movements such as eversion. The action of foot eversion involves the peroneus longus and peroneus brevis muscles, both of which are innervated by the S1 nerve root. In contrast, hip flexion is predominantly controlled by the lumbar plexus and involves primarily the L2 and L3 nerve roots, which innervate the iliopsoas muscle. Knee extension is mainly governed by the L3 and L4 nerve roots, which control the quadriceps muscle. Hip adduction is associated with the obturator nerve, primarily originating from the L2-L4 nerve roots. This highlights the distinct functions of each nerve root and their associated muscle actions, clearly establishing foot eversion as a characteristic movement of the S1 nerve root.

8. During the standing flexion test for the lumbar spine, what indicates a positive test?

A. No movement of both PSIS

B. The PSIS that moves inferiorly first

C. The PSIS that moves cranially first

D. Symmetry in PSIS movement

A positive standing flexion test for the lumbar spine is indicated when one of the posterior superior iliac spines (PSIS) moves cranially (upwards) first as the patient bends forward. This suggests asymmetrical motion, which can be indicative of a restriction or dysfunction in the lumbar spine or sacroiliac joints. In this context, the cranial movement of one PSIS may suggest that there is an issue such as sacroiliac joint dysfunction or an abnormality in the pelvic mechanics. When one side moves before the other, it reflects a lack of coordinated motion that is typically expected when the pelvis and lumbar spine accommodate flexion together. Other responses may reflect various forms of movement or symmetry, which do not signify the same degree of dysfunction as the presence of one PSIS moving cranially before the other. For instance, if both PSIS do not move or move symmetrically, it may indicate a different type of mechanical issue, but not necessarily the same level of dysfunction that the cranial movement first would imply. Thus, the movement of the PSIS cranially first is a key indicator of dysfunction recognized during this test.

9. Finkelstein's test primarily evaluates which tendons?

A. Flexor pollicis longus and flexor carpi ulnaris

B. Abductor pollicis longus and extensor pollicis brevis

C. Extensor pollicis longus and extensor carpi radialis

D. Extensor digitorum and abductor digiti minimi

Finkelstein's test is specifically designed to evaluate the presence of de Quervain's tenosynovitis, which is an inflammatory condition affecting the tendons of the wrist. The primary tendons involved in this condition are the abductor pollicis longus and the extensor pollicis brevis. During Finkelstein's test, the patient is asked to make a fist with the thumb inside the fingers, and then the wrist is ulnarly deviated. This maneuver places stress on the abductor pollicis longus and extensor pollicis brevis tendons as they pass through the first dorsal compartment of the wrist. If this movement results in pain along the radial side of the wrist, it is indicative of de Quervain's tenosynovitis, confirming that the tendons in question are indeed the abductor pollicis longus and extensor pollicis brevis. Therefore, this specific association substantiates the correctness of the answer provided.

SAMPLE

10. What are the primary symptoms of juvenile rheumatoid arthritis?

A. Joint stiffness and swelling

B. Fever and rash

C. Severe pain and immobility

D. Muscle weakness and discomfort

Juvenile rheumatoid arthritis (JRA), also known as juvenile idiopathic arthritis, is characterized by inflammation in the joints and can have systemic effects. The primary symptoms include joint stiffness and swelling, which often occur in the morning or after periods of rest, and may be accompanied by systemic symptoms such as fever and rash. Fever and rash are particularly notable as they indicate the systemic involvement of the disease. JRA can manifest in various subtypes, some of which are associated with fever and characteristic rash, especially in the systemic form of the disease. This systemic aspect distinguishes JRA from other forms of arthritis that may primarily focus on joint symptoms without systemic signs. Though joint stiffness and swelling are classic and significant symptoms of JRA, the presence of fever and rash highlights the inflammatory nature of the condition and the potential for wider systemic involvement, which is critical in diagnosing and managing the disease effectively.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nptemsk.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE