NUCLEAR MEDICINE PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://nuclearmed.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the required time for written reporting of a medical event to the NRC?

A. 24 hours

B. 30 calendar days

C. 72 hours

D. 5 regular work days

2. Pulse rates may be determined by all of the following methods except:

A. Listening to the heart with a stethoscope

B. Using a pulse oximeter

C. Multiplying the respiration rate by 4

D. Analyzing the electrocardiogram

3. What region of interest is used for detecting a left to right cardiac shunt?

A. Superior vena cava

B. Left ventricle

C. Right ventricle and great vessels

D. One or both lungs

4. What could happen if a 99mTc sulfur colloid preparation is boiled for too long?

A. It will result in chemically unstable particles

B. It will result in colloidal particles that are clumpy

C. It will result in colloidal particles that are too large

D. It will result in optimal size particles

5. How often should wipe tests of areas where radiopharmaceuticals are prepared be performed?

A. Weekly

B. Only if contamination occurs

C. Every day on which radiopharmaceuticals are used

D. On a reasonable schedule

SAMPLE

6. Which type of imaging is typically used to assess liver function in a hepatobiliary study?

A. Liver biopsy

B. Ultrasound

C. CT scan

D. Radionuclide imaging

7. If visualized, the gallbladder fossa may be seen on what view(s) of a liver spleen image?

A. Anterior and posterior

B. Right lateral only

C. Anterior only

D. Anterior and right lateral

8. What is the significance of the half-life of a radionuclide in clinical practice?

A. It affects patient scheduling for imaging

B. It determines the radiation dose to the staff

C. It influences the decay rate of the radionuclide

D. It impacts the type of shielding needed

9. In evaluating anaphylactic reactions, which symptom is more characteristic of a mild allergic reaction rather than anaphylaxis?

A. Angioedema

B. Wheezing

C. Skin rash

D. Hypotension

10. What sign is required to be posted in imaging rooms according to the NRC?

A. No posting is required

B. Caution: Radiation area

C. Caution: Radioactive materials

D. Caution: High Radiation Area

SAMPLE

Answers

SAMPLE

1. B
2. C
3. D
4. C
5. D
6. D
7. D
8. C
9. C
10. C

SAMPLE

Explanations

SAMPLE

1. What is the required time for written reporting of a medical event to the NRC?

A. 24 hours

B. 30 calendar days

C. 72 hours

D. 5 regular work days

The correct answer is that the required time for written reporting of a medical event to the Nuclear Regulatory Commission (NRC) is 30 calendar days. This time frame is established to ensure that incidents involving patient safety and quality of care are documented adequately and addressed in a timely manner. In nuclear medicine, any significant medical event—especially those that could lead to an adverse effect on a patient's health—must be reported to the NRC as part of their oversight responsibilities. The 30-day window allows facility personnel to gather all necessary information related to the event, conduct a thorough investigation if needed, and prepare a comprehensive report that includes details about the incident, the patients involved, and any corrective actions taken to mitigate future occurrences. This protocol is vital for enhancing safety practices and maintaining patient trust in the healthcare system.

2. Pulse rates may be determined by all of the following methods except:

A. Listening to the heart with a stethoscope

B. Using a pulse oximeter

C. Multiplying the respiration rate by 4

D. Analyzing the electrocardiogram

Each option describes a different method of determining pulse rates, but one stands out as inappropriate. Listening to the heart with a stethoscope is a common practice in clinical settings, allowing for the direct observation of heart sounds and rhythm, which can aid in determining pulse rates. Similarly, utilizing a pulse oximeter provides a non-invasive way to measure the oxygen saturation levels in the blood and can also give readings of heart rate. Analyzing an electrocardiogram (ECG) is another reliable method for assessing the heart's electrical activity, which directly reflects heart rate. However, multiplying the respiration rate by 4 does not provide a clinical or physiologically accurate measure of pulse rate. While respiration and heart rates are often related, the two do not have a fixed ratio that can be applied across different individuals or situations. Each person's respiratory and heart rates can vary independently, making this method unreliable for determining pulse rates. Therefore, this option does not accurately describe a valid technique for measuring heart rate.

SAMPLE

3. What region of interest is used for detecting a left to right cardiac shunt?

A. Superior vena cava

B. Left ventricle

C. Right ventricle and great vessels

D. One or both lungs

Detecting a left to right cardiac shunt primarily involves evaluating the pulmonary circulation, which is directly impacted by such shunts. In the case of a left to right shunt, blood flows from the left heart (typically from the left atrium or left ventricle) into the right heart (right atrium or right ventricle) and subsequently into the lungs. When assessing for a left to right shunt, the region of interest is often one or both lungs, as a significant amount of blood from the left side of the heart is diverted to the right, leading to increased blood volume in the pulmonary circulation. This can result in various detectable changes in lung perfusion and can be visualized using imaging techniques. This increased blood flow to the lungs can cause signs of pulmonary overcirculation or congestion on imaging studies. The other regions mentioned, such as the superior vena cava, the left ventricle, or the right ventricle and great vessels, do not provide the requisite information to effectively demonstrate the impact of a left to right shunt. These areas are more related to the anatomic structure and initial flow of blood rather than the functional consequences on pulmonary vasculature.

4. What could happen if a 99mTc sulfur colloid preparation is boiled for too long?

A. It will result in chemically unstable particles

B. It will result in colloidal particles that are clumpy

C. It will result in colloidal particles that are too large

D. It will result in optimal size particles

When a 99mTc sulfur colloid preparation is boiled for an extended period, one of the significant effects is that it can lead to the formation of particles that are too large. This occurs because prolonged boiling can cause excessive aggregation of the colloidal particles. The heat and the duration can promote the interaction among the colloidal sulfur particles, leading to larger agglomerates forming rather than maintaining a stable, uniform size distribution. In nuclear medicine, the size of colloidal particles is crucial, particularly for applications such as liver and spleen imaging or lymphoscintigraphy. Ideally, the particles should be small enough to be taken up by reticuloendothelial cells without clogging the vascular spaces. Larger particles can lead to complications, such as reduced imaging quality or incorrect localization, and may hinder proper distribution within the organs of interest. Maintaining the optimal size of these colloidal particles is vital for achieving the desired imaging results and ensuring that adequate biological processes are visualized accurately.

SAMPLE

5. How often should wipe tests of areas where radiopharmaceuticals are prepared be performed?

A. Weekly

B. Only if contamination occurs

C. Every day on which radiopharmaceuticals are used

D. On a reasonable schedule

Wipe tests in areas where radiopharmaceuticals are prepared are essential for ensuring safety and contamination control. The principle of performing these tests on a reasonable schedule underscores the importance of both regular monitoring and flexibility based on operational needs and contamination risks. A reasonable schedule means that wipe tests should be systematic to maintain compliance with safety standards and regulations, but the frequency can vary based on factors such as the volume of radiopharmaceuticals being handled, the type of procedures performed, and the specific protocols of the facility. This approach allows for both diligence in monitoring and the efficient use of resources, ensuring that potential contamination is detected and addressed without being overly burdensome to operations. By opting for a reasonable schedule, facilities can tailor their practices to match their unique circumstances, thereby enhancing safety while also being practical. Using this strategy is a well-recognized best practice in nuclear medicine, ensuring ongoing vigilance against contamination without necessitating excessive or unnecessary testing.

6. Which type of imaging is typically used to assess liver function in a hepatobiliary study?

A. Liver biopsy

B. Ultrasound

C. CT scan

D. Radionuclide imaging

Radionuclide imaging is a specialized technique used in nuclear medicine that allows for the assessment of liver function through the evaluation of hepatobiliary function. This method typically involves the use of a radiopharmaceutical that is taken up by the liver and subsequently excreted into the bile. By tracking the distribution and activity of the radiopharmaceutical, practitioners can gain valuable insights into the liver's ability to perform its vital functions, such as bile production and excretion. In a hepatobiliary study, radionuclide imaging can provide specific information regarding conditions like acute or chronic liver disease, biliary obstructions, and other hepatic disorders. The imaging technique showcases the dynamic process of liver function in real time, allowing for a functional assessment, which is crucial for diagnosing liver-related conditions. Other imaging modalities, while useful in visualizing the structure of the liver and surrounding organs, do not primarily assess liver function. For instance, a liver biopsy involves taking a tissue sample to evaluate liver pathology directly but does not provide functional insights. Ultrasound is beneficial for examining liver size, structure, and blood flow but lacks the functional assessment capabilities of radionuclide imaging. Similarly, a CT scan can visualize liver anatomy and identify lesions but does not assess liver function in the

SAMPLE

7. If visualized, the gallbladder fossa may be seen on what view(s) of a liver spleen image?

A. Anterior and posterior

B. Right lateral only

C. Anterior only

D. Anterior and right lateral

The gallbladder fossa is a region where the gallbladder is located on the liver, and it can be visualized through imaging techniques commonly used in nuclear medicine, particularly during liver spleen scans. When examining the views for imaging these organs, both anterior and right lateral views are the most effective for visualizing anatomical structures in the upper abdominal area, including the gallbladder fossa. The anterior view allows for a direct look at the liver and its landmarks, making it possible to identify the gallbladder fossa easily. The right lateral view complements this by providing an alternative perspective that can enhance visualization of the gallbladder's position relative to the liver and surrounding organs. These two views together maximize the likelihood of clearly seeing the gallbladder fossa. In contrast, the other options either limit the views to just one perspective or exclude one of the two key views necessary for optimal visualization. Therefore, the combination of anterior and right lateral views is indeed the best approach for assessing the gallbladder fossa on a liver spleen image.

8. What is the significance of the half-life of a radionuclide in clinical practice?

A. It affects patient scheduling for imaging

B. It determines the radiation dose to the staff

C. It influences the decay rate of the radionuclide

D. It impacts the type of shielding needed

The half-life of a radionuclide is a critical factor in clinical practice as it directly influences the decay rate of the radionuclide. The half-life is defined as the time required for half of the radioactive atoms in a sample to decay. This property is fundamental in nuclear medicine because it determines how quickly the radionuclide will lose its radioactivity and how long it will remain effective for diagnostic or therapeutic purposes. For example, radionuclides with short half-lives decay quickly, making them useful for imaging purposes where rapid accumulation and clearance are needed, such as in PET scans. Conversely, radionuclides with longer half-lives are utilized in therapies where sustained radiation exposure is necessary to achieve therapeutic effects, such as in the treatment of certain cancers. The influence of half-life on the decay rate also has implications for dosing and safety protocols, as practitioners must account for how long a radionuclide will be active in the body or in the environment when planning procedures and ensuring patient and staff safety. Thus, understanding the half-life allows healthcare professionals to adequately plan the timing of imaging or treatment sessions, optimize the use of radionuclides in patient care, and manage waste effectively.

SAMPLE

9. In evaluating anaphylactic reactions, which symptom is more characteristic of a mild allergic reaction rather than anaphylaxis?

A. Angioedema

B. Wheezing

C. Skin rash

D. Hypotension

In the context of allergic reactions, a skin rash is more characteristic of a mild allergic reaction compared to anaphylaxis. Allergic reactions can typically manifest in various ways, including skin symptoms like rashes or hives, which often accompany milder reactions. These skin-related symptoms, though they can occur in anaphylaxis, are not exclusive to it and are commonly seen in less severe allergic responses. On the other hand, symptoms such as angioedema (which involves swelling deeper in the skin), wheezing (indicating airway constriction), and hypotension (a significant drop in blood pressure) are associated with more severe reactions and are hallmark signs of anaphylaxis. The presence of these symptoms suggests a systemic response that could be life-threatening, distinguishing them clearly from the milder manifestations that predominantly include localized rashes. Thus, a skin rash serves as an indicator of a less severe allergic reaction, making it the correct identification in this scenario.

10. What sign is required to be posted in imaging rooms according to the NRC?

A. No posting is required

B. Caution: Radiation area

C. Caution: Radioactive materials

D. Caution: High Radiation Area

In nuclear medicine imaging rooms, it is critical to ensure that proper signage is displayed as a safety measure for both patients and staff. The Nuclear Regulatory Commission (NRC) requires specific warnings to be posted in areas where radioactive materials are used. The correct sign, "Caution: Radioactive materials," is essential because it informs individuals entering the area that they may be in proximity to materials that emit radiation. This helps to prevent unnecessary exposure and ensures that everyone is aware of the potential hazards associated with radioactive substances. These signs serve as a protective measure, emphasizing the importance of following safety protocols in imaging rooms where radioactive materials are present during diagnostic or therapeutic procedures. Other options like "Caution: Radiation area" or "Caution: High Radiation Area" are similar but pertain to different levels of radiation exposure potential. "Caution: Radiation area" is used for areas where an individual could receive a dose greater than a certain threshold in an hour, while "Caution: High Radiation Area" indicates zones where the radiation doses are significantly higher. However, in the context of imaging rooms specifically dealing with radioactive materials, the indication of "Caution: Radioactive materials" is the most relevant and legally mandated sign that must be present.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nuclearmed.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE