OBJECTIVE STRUCTURED CLINICAL ASSESSMENT (OSCA) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://osca.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the normal diastolic blood pressure range?

A. 50-70 mmHg (diastolic)

B. 60-90 mmHg (diastolic)

C. 70-95 mmHg (diastolic)

D. 40-80 mmHg (diastolic)

2. Bradycardia is defined as:

A. Above 100 BPM

B. 60-100 BPM

C. Below 60 BPM

D. 100-120 BPM

3. What is the first-line treatment for suspected anaphylaxis?

A. Intramuscular epinephrine

B. Oral antihistamines

C. Inhaled bronchodilators

D. IV antibiotics

4. Which statement best reflects the essential content when documenting informed consent?

A. Consent is valid with verbal agreement only.

B. A signed form alone suffices; capacity not required.

C. Key elements include explanation, alternatives, risks, benefits, and documentation of capacity.

D. Consent is not required for emergency procedures.

5. How should you introduce yourself and confirm the patient's identity at the start of a consultation?

A. Rely on receptionist to identify the patient.

B. Confirm the patient's identity and preferred name, and obtain consent to proceed.

C. State your name and role, confirm the patient's identity and preferred name, and obtain consent to proceed.

D. Skip introduction if the patient seems unwell.

SAMPLE

6. Which ISBAR component includes vital signs and current observations?

A. Assessment

B. Background

C. Recommendations

D. Situation

7. In a pain assessment using the PQRST mnemonic, what does R stand for and what question would you ask?

A. Radiating - Does the pain move or radiate? On movement?

B. Reaction - How does the pain respond to touch?

C. Range - How wide is the area affected?

D. Radius - Does the pain extend to distant sites?

8. Which safety check is essential before prescribing a high-risk medication?

A. Confirm patient identity.

B. Review allergies.

C. Verify allergies, review drug interactions and organ function, assess dose.

D. Confirm patient identity, verify allergies, review drug interactions and organ function, dose appropriateness, consent, and documentation.

9. Which ISBAR component may include reason for admission, medical history, and medications?

A. Situation - presenting problem

B. Assessment - vitals

C. Background - may include reason for admission, medical history, medications

D. Recommendations - plan

10. What is the recommended sequence of an abdominal examination, and why?

A. Palpation, auscultation, percussion, inspection.

B. Inspect, auscultate, percussion, palpation; palpation deferred to avoid altering bowel sounds.

C. Auscultation first, then inspection.

D. Percussion last.

SAMPLE

Answers

SAMPLE

1. B
2. C
3. A
4. C
5. C
6. A
7. A
8. D
9. D
10. B

SAMPLE

Explanations

SAMPLE

1. What is the normal diastolic blood pressure range?

A. 50-70 mmHg (diastolic)

B. 60-90 mmHg (diastolic)

C. 70-95 mmHg (diastolic)

D. 40-80 mmHg (diastolic)

Diastolic pressure is the arterial pressure when the heart is resting between beats, and it reflects the baseline pressure the arteries maintain. In adults, a diastolic value around 60–90 mmHg is generally considered normal. Values lower than about 60 mmHg can indicate hypotension, while sustained values at or above about 90–100 mmHg suggest hypertension. The 60–90 mmHg range sits in the conventional normal-to-borderline zone, avoiding the clearly low end and not extending into higher values that would indicate hypertension. The other ranges include numbers that are clearly too low or push into ranges associated with elevated blood pressure, so this range best fits the typical normal diastolic expectation.

2. Bradycardia is defined as:

A. Above 100 BPM

B. 60-100 BPM

C. Below 60 BPM

D. 100-120 BPM

Bradycardia means a resting heart rate that's slower than normal. A typical resting range is about 60 to 100 beats per minute, so when the rate falls below 60, that's bradycardia. This threshold helps distinguish it from normal variation and from tachycardia, which is a rate above 100. In some people, such as well-trained athletes or during sleep, a rate under 60 can be normal due to higher vagal tone. However, if bradycardia causes symptoms (dizziness, fatigue, fainting) or reflects underlying issues (conduction system disease, hypothyroidism, or medication effects), it warrants evaluation.

3. What is the first-line treatment for suspected anaphylaxis?

A. Intramuscular epinephrine

B. Oral antihistamines

C. Inhaled bronchodilators

D. IV antibiotics

Prompt recognition and rapid reversal of the life-threatening collapse in anaphylaxis is the focus here. The best first step is giving epinephrine by intramuscular injection as soon as anaphylaxis is suspected. Epinephrine tackles the main urgent problems at once: it constricts swollen blood vessels to raise blood pressure, reduces airway edema, and relaxes bronchial smooth muscle to relieve wheeze and breathing difficulty. It also stimulates the heart to improve perfusion and dampens further mediator release from mast cells. The intramuscular route, typically into the mid-outer thigh, provides the quickest and most reliable absorption in an emergent setting. In adults a standard initial dose is about 0.3 to 0.5 mg, with repeat doses every 5 to 15 minutes as needed while symptoms persist or worsen; dosing for children is weight-based (about 0.01 mg/kg per dose, up to a practical maximum), with repeats as necessary. After administering epinephrine, call for emergency help, ensure the airway, give high-flow oxygen, and establish IV access for fluids and monitoring. While antihistamines or steroids and inhaled bronchodilators may be used as adjuncts, they do not rapidly reverse the life-threatening features of anaphylaxis and should not replace epinephrine as the initial treatment. IV antibiotics have no role in the acute management of anaphylaxis.

SAMPLE

4. Which statement best reflects the essential content when documenting informed consent?

A. Consent is valid with verbal agreement only.

B. A signed form alone suffices; capacity not required.

C. Key elements include explanation, alternatives, risks, benefits, and documentation of capacity.

D. Consent is not required for emergency procedures.

Informed consent rests on patient autonomy and understanding. The essential content ensures the patient receives a clear explanation of what will be done, the alternatives (including the option of not proceeding), and the risks and benefits involved, with an evaluation that the patient has the capacity to understand and make the decision, plus proper documentation of that capacity and the discussion. A signed form or a verbal assent alone does not guarantee understanding or voluntariness; capacity must be assessed and the decision recorded. This combination—information, alternatives, risks and benefits, capacity, and documentation—provides a valid, ethically sound basis for consent.

5. How should you introduce yourself and confirm the patient's identity at the start of a consultation?

A. Rely on receptionist to identify the patient.

B. Confirm the patient's identity and preferred name, and obtain consent to proceed.

C. State your name and role, confirm the patient's identity and preferred name, and obtain consent to proceed.

D. Skip introduction if the patient seems unwell.

Introducing yourself and confirming identity right at the start is about clear, respectful, patient-centered communication that promotes safety. You should say your name and role, check the patient's identity and preferred name, and obtain consent to proceed. This combination sets who is speaking, reduces the risk of misidentification, respects how the patient wants to be addressed, and ensures they agree to continue with the consultation. Relying on the receptionist to identify the patient bypasses direct confirmation and can lead to errors. Skipping an introduction, even if the patient is unwell, can undermine trust and fail to establish a collaborative tone. Confirming identity and consent without stating your name and role can leave the patient unsure about who is leading their care.

6. Which ISBAR component includes vital signs and current observations?

A. Assessment

B. Background

C. Recommendations

D. Situation

In ISBAR, the vital signs and current observations are interpreted and summarized in the Assessment. This part is about evaluating the patient's present condition based on the data you've collected—why the readings matter, whether they show stability or deterioration, and what they imply for the plan. For example, a growing heart rate, rising respiratory rate, and dropping oxygen saturation would be interpreted here as potential deterioration and a need for escalation or intervention, rather than just listing numbers. The other sections serve different roles: the Situation states what is happening now, the Background provides context from the patient's history, and the Recommendation outlines what actions are needed. So the component that includes vital signs and current observations is the Assessment.

SAMPLE

7. In a pain assessment using the PQRST mnemonic, what does R stand for and what question would you ask?

A. Radiating - Does the pain move or radiate? On movement?

B. Reaction - How does the pain respond to touch?

C. Range - How wide is the area affected?

D. Radius - Does the pain extend to distant sites?

R stands for Radiation (region and radiation). It's about where the pain is and whether it spread beyond the initial spot. Ask something like: "Does the pain radiate to other areas?" for example down the arm, into the jaw, or other regions. This helps distinguish localized pain from nerve-related or referred pain, which is important for guiding diagnosis and management. The other terms listed aren't used in this mnemonic.

8. Which safety check is essential before prescribing a high-risk medication?

A. Confirm patient identity.

B. Review allergies.

C. Verify allergies, review drug interactions and organ function, assess dose.

D. Confirm patient identity, verify allergies, review drug interactions and organ function, dose appropriateness, consent, and documentation.

Before prescribing a high-risk medication, a comprehensive safety check is essential to prevent harm. The best answer includes confirming patient identity, verifying allergies, reviewing potential drug interactions and organ function, ensuring the dose is appropriate, and obtaining consent with proper documentation. This full checklist helps prevent errors from misidentification, undisclosed allergies, harmful drug interactions, or inappropriate dosing, and it supports ethical practice and legal accountability. Shorter checks miss critical components: identity or allergy checks alone don't address interactions, dosing, consent, or documentation.

9. Which ISBAR component may include reason for admission, medical history, and medications?

A. Situation - presenting problem

B. Assessment - vitals

C. Background - may include reason for admission, medical history, medications

D. Recommendations - plan

In ISBAR, the Background section provides the patient's context and history to frame the current situation. It is where you include the reason for admission, past medical history, current medications, allergies, and prior relevant info. This context helps the team understand baseline risk, potential drug interactions, and why the patient is here, which informs the subsequent assessment and recommendations. The Situation describes the present issue or reason you are communicating now; the Assessment covers your current status, including data like vitals; and the Recommendations outline what you think should happen next. So the details about why the patient was admitted and their medications belong in Background, not in the other sections.

SAMPLE

10. What is the recommended sequence of an abdominal examination, and why?

A. Palpation, auscultation, percussion, inspection.

B. Inspect, auscultate, percussion, palpation; palpation deferred to avoid altering bowel sounds.

C. Auscultation first, then inspection.

D. Percussion last.

Doing the abdominal exam in the right order is essential to avoid distorting the findings. The best sequence is to inspect first, then auscultate, then percussion, and finally palpate. Inspecting lets you note the abdomen's shape, contour, scars, distension, and any visible masses or signs of distress. Auscultation should come next because listening for bowel sounds and vascular noises is best done on an untouched abdomen; touching or peeking can alter motility and sound transmission, leading to misinterpretation. Percussion follows to help identify tympany versus dullness and to assess for fluid, masses, or organ borders without having affected the bowel sounds. Palpation is performed last, starting with light palpation to detect surface tenderness and guarding, then deeper palpation to feel for organs and masses. Since palpation and percussion can change bowel activity and the perception of tenderness, delaying them until after auscultation preserves accuracy.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://osca.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE