OBSTETRICS CAPSTONE PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://obcapstone.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is chorioamnionitis?

A. An infection of the amniotic fluid and membranes surrounding the fetus

B. A type of fetal presentation during labor

C. A hormonal change that occurs during pregnancy

D. An abnormality detected through ultrasound

2. What information should be included when teaching a client experiencing stress incontinence at 32 weeks of gestation?

A. Decrease dietary fiber.

B. Practice Kegel exercises.

C. Restrict daily fluid intake.

D. Avoid daily exercise.

3. What are the risks associated with prolonged rupture of membranes?

A. Increased risk of maternal hemorrhage

B. Increased risk of infection and preterm labor

C. Increased risk of fetal brain injury

D. Increased risk of placenta previa

4. What is the typical duration of a full-term pregnancy?

A. 32 weeks

B. 36 weeks

C. 40 weeks

D. 44 weeks

5. Which anticoagulant is frequently used to prevent thrombosis in pregnancy?

A. Warfarin

B. Low molecular weight heparin (LMWH)

C. Aspirin

D. Clopidogrel

SAMPLE

6. What is a common symptom associated with gestational hypertension?

A. Increased energy levels

B. Severe headaches

C. Weight loss

D. Reduced thirst

7. Which vitamin is particularly important for the development of the fetal brain during pregnancy?

A. Vitamin C

B. Folic acid

C. Vitamin D

D. Vitamin K

8. What assessment finding in a postpartum patient warrants immediate intervention?

A. Temperature of 99.0°F

B. Fundus located at the umbilicus

C. Bright red vaginal bleeding

D. Increased appetite

9. Which condition is associated with an increased risk during labor in late-term pregnancies?

A. Gestational diabetes

B. Decreased amniotic fluid

C. Preterm labor

D. Placenta previa

10. What is the primary role of an obstetrician during delivery?

A. To administer anesthesia for the mother

B. To manage labor and delivery, ensuring the health of mother and fetus

C. To perform surgeries unrelated to labor

D. To monitor fetal heart rates exclusively

SAMPLE

Answers

SAMPLE

1. A
2. B
3. B
4. C
5. B
6. B
7. B
8. C
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is chorioamnionitis?

A. An infection of the amniotic fluid and membranes surrounding the fetus

B. A type of fetal presentation during labor

C. A hormonal change that occurs during pregnancy

D. An abnormality detected through ultrasound

Chorioamnionitis is defined as an infection of the amniotic fluid and the membranes surrounding the fetus. This condition typically arises when bacteria ascend from the vagina into the uterus, leading to inflammation and infection of the amniotic sac. It is a significant concern during labor because it can result in serious complications for both the mother and the fetus, including preterm labor, increased risk of cesarean section, and potential neonatal infections. Understanding chorioamnionitis is important for obstetric care, as it requires prompt diagnosis and treatment, often necessitating the administration of antibiotics and, in some cases, early delivery. Recognizing the signs and symptoms, such as maternal fever, fetal tachycardia, and foul-smelling amniotic fluid, is crucial for healthcare providers in managing this condition effectively.

2. What information should be included when teaching a client experiencing stress incontinence at 32 weeks of gestation?

A. Decrease dietary fiber.

B. Practice Kegel exercises.

C. Restrict daily fluid intake.

D. Avoid daily exercise.

Practicing Kegel exercises is highly beneficial for clients experiencing stress incontinence, particularly during pregnancy. These exercises strengthen the pelvic floor muscles, which support the bladder and may help alleviate incontinence symptoms. Strengthening these muscles can promote better control over urination, particularly during physical activities that may induce stress incontinence such as walking, coughing, or laughing. Kegel exercises are safe for most pregnant women and can be easily incorporated into their daily routine. By increasing muscle tone and improving the pelvic floor strength, Kegel exercises can enhance bladder control and potentially reduce episodes of incontinence. In contrast, decreasing dietary fiber might lead to constipation, which can exacerbate urinary incontinence. Restricting fluid intake is generally discouraged as it can lead to dehydration and may not effectively resolve incontinence issues. Avoiding daily exercise is not advisable either, as physical activity is important for overall health during pregnancy, and it can help improve muscle tone, including the pelvic floor muscles, when performed safely.

SAMPLE

3. What are the risks associated with prolonged rupture of membranes?

A. Increased risk of maternal hemorrhage

B. Increased risk of infection and preterm labor

C. Increased risk of fetal brain injury

D. Increased risk of placenta previa

Prolonged rupture of membranes, defined as the rupture of the amniotic sac more than 18 hours before delivery, poses significant risks primarily related to infection and the likelihood of preterm labor. When the protective amniotic sac is compromised for an extended period, the risk of ascending infections increases, leading to conditions such as chorioamnionitis, which can affect both the mother and fetus. Additionally, prolonged rupture can initiate uterine contractions, resulting in preterm labor, particularly if the pregnancy is not at term. The other options, while they relate to various obstetric complications, are not directly associated with prolonged rupture of membranes in the same way. Maternal hemorrhage and fetal brain injury are more related to other risk factors and complications that may arise during labor and delivery, rather than specifically a consequence of prolonged rupture. Similarly, placenta previa is a condition defined by the position of the placenta and is not influenced by the status of the membranes. Thus, the primary concern with prolonged rupture of membranes centers on the heightened risk of infection and preterm labor, affirming the choice made.

4. What is the typical duration of a full-term pregnancy?

A. 32 weeks

B. 36 weeks

C. 40 weeks

D. 44 weeks

A full-term pregnancy typically lasts about 40 weeks, which is counted from the first day of the last menstrual period (LMP) to the birth of the baby. This timeframe includes three trimesters, with the first trimester encompassing weeks 1 to 12, the second trimester from weeks 13 to 26, and the third trimester from weeks 27 to 40. Deliveries are generally considered full-term if they occur between 39 and 40 weeks; however, up to 42 weeks is still within the span of full-term classification. This duration allows for optimal fetal development, including necessary growth, organ maturation, and overall readiness for life outside the womb. Accurate understanding of gestational age is critical for monitoring both maternal and fetal health throughout pregnancy and helps in planning for labor and delivery.

SAMPLE

5. Which anticoagulant is frequently used to prevent thrombosis in pregnancy?

A. Warfarin

B. Low molecular weight heparin (LMWH)

C. Aspirin

D. Clopidogrel

Low molecular weight heparin (LMWH) is frequently used to prevent thrombosis in pregnancy due to its safety profile and effectiveness. During pregnancy, there is an increased risk of venous thromboembolism due to physiological changes such as enhanced coagulation and venous stasis. LMWH is preferred because it does not cross the placenta and is less likely to cause harm to the developing fetus compared to other anticoagulants. Additionally, LMWH has predictable pharmacokinetics, allowing for easier management with less frequent dosing compared to unfractionated heparin. While it is important to monitor certain parameters, the risk of complications is significantly reduced, making LMWH a suitable choice for pregnant patients, especially those with a history of thrombosis or other risk factors. Other anticoagulants listed are not the first-line options for pregnancy. For instance, warfarin is contraindicated because it can cross the placenta and is associated with teratogenic effects. Aspirin is generally used for different indications in pregnancy, such as preventing preeclampsia rather than as a primary anticoagulant. Clopidogrel's use in pregnancy is also limited and generally not recommended for routine prophylaxis against thrombosis.

6. What is a common symptom associated with gestational hypertension?

A. Increased energy levels

B. Severe headaches

C. Weight loss

D. Reduced thirst

Gestational hypertension, also known as pregnancy-induced hypertension, is characterized by high blood pressure that develops after the 20th week of pregnancy. One common symptom associated with this condition is the occurrence of severe headaches. These headaches may be caused by elevated blood pressure, and they can also signal potential complications, such as progression to preeclampsia. Women experiencing severe headaches during pregnancy should be evaluated by a healthcare provider, as this symptom can indicate the need for immediate attention. In contrast, increased energy levels typically do not correlate with gestational hypertension; in fact, women may often feel more fatigued due to the stress on their bodies. Weight loss is not a symptom associated with gestational hypertension as, generally, women experience weight gain during pregnancy. Reduced thirst is also unlikely; women usually maintain their hydration, and a decrease in thirst could signal other health issues unrelated to gestational hypertension. Thus, severe headaches stand out as a significant and concerning symptom of this condition.

SAMPLE

7. Which vitamin is particularly important for the development of the fetal brain during pregnancy?

A. Vitamin C

B. Folic acid

C. Vitamin D

D. Vitamin K

Folic acid plays a crucial role in the development of the fetal brain during pregnancy. It is a B vitamin that is essential for DNA synthesis and repair, as well as for cell division and growth. Adequate folic acid intake during the early stages of pregnancy is particularly vital because it helps prevent neural tube defects, which can affect the brain and spinal cord. Folic acid supports the formation of neural pathways and is involved in the production of neurotransmitters, which are crucial for brain development and function. This nutrient's significance in the early formation of the fetal neural structure underlines why it is strongly recommended for all women of childbearing age to ensure they receive sufficient folic acid before conception and throughout pregnancy. In comparison, the other vitamins listed, though important for various aspects of health and development, do not have the same direct relation to fetal brain development as folic acid does. Vitamin C is important for the immune system and tissue formation, Vitamin D supports bone health and immune function, and Vitamin K is essential for blood clotting and bone metabolism. None of these vitamins have the same strong evidence linking them directly to preventing neural tube defects and directly impacting brain development as folic acid.

8. What assessment finding in a postpartum patient warrants immediate intervention?

A. Temperature of 99.0°F

B. Fundus located at the umbilicus

C. Bright red vaginal bleeding

D. Increased appetite

Bright red vaginal bleeding in a postpartum patient is a significant assessment finding that requires immediate intervention. This type of bleeding, known as lochia rubra, typically occurs during the first few days following delivery, but it should gradually decrease in volume and transition in color from red to pink and then to yellowish-white as the body heals. When bright red bleeding is excessive or sudden, it can indicate complications such as retained placental fragments, uterine atony, or lacerations, all of which can lead to postpartum hemorrhage. Such conditions pose serious risks to the mother's health, potentially resulting in hypovolemic shock if not addressed promptly. Therefore, the presence of bright red vaginal bleeding prompts urgent evaluation and management to ensure the safety and wellbeing of the postpartum patient. By comparison, a mild fever of 99.0°F is generally not concerning as it may be a normal response to physiological changes after labor. A fundus located at the umbilicus is typical around days one to two postpartum, indicating appropriate uterine involution. An increased appetite can also be a normal finding as a woman begins to recover and resume her routine following childbirth. It is the bright red vaginal bleeding that stands out as a critical indicator requiring immediate attention to

SAMPLE

9. Which condition is associated with an increased risk during labor in late-term pregnancies?

A. Gestational diabetes

B. Decreased amniotic fluid

C. Preterm labor

D. Placenta previa

The condition associated with an increased risk during labor in late-term pregnancies is decreased amniotic fluid, also known as oligohydramnios. Amniotic fluid plays a crucial role in cushioning the fetus, facilitating movements, and promoting lung development. In late-term pregnancies, insufficient fluid can lead to complications such as umbilical cord compression, which can affect fetal heart rate and oxygenation during labor. It may also increase the risk of labor augmentation and the need for interventions, such as cesarean delivery, to ensure the safety of both mother and baby. While conditions like gestational diabetes and placenta previa can present risks, they are more relevant to specific management strategies and outcomes rather than being directly associated with the physiological changes during labor itself. Preterm labor is characterized by contractions and cervical changes before 37 weeks of gestation, and it does not pertain to late-term pregnancies, where gestation has progressed beyond 40 weeks. Therefore, the critical element linking decreased amniotic fluid to increased labor risk in late-term pregnancies underscores the importance of monitoring and management in these circumstances.

10. What is the primary role of an obstetrician during delivery?

A. To administer anesthesia for the mother

B. To manage labor and delivery, ensuring the health of mother and fetus

C. To perform surgeries unrelated to labor

D. To monitor fetal heart rates exclusively

The primary role of an obstetrician during delivery is to manage labor and delivery, ensuring the health of both the mother and the fetus. This encompasses a wide range of responsibilities, including monitoring the progress of labor, assessing the wellbeing of the fetus, and addressing any complications that may arise during the delivery process. An obstetrician is trained to provide comprehensive care throughout labor and delivery. They are responsible for making critical decisions about interventions that may be necessary, such as performing a cesarean section if complications occur. Their expertise allows them to respond effectively to changes in the mother's or baby's condition, aiming to achieve the best possible outcomes for both. While administering anesthesia may be a part of care during delivery, it is typically performed by an anesthesiologist or nurse anesthetist, not the obstetrician themselves. The obstetrician's focus is on obstetric care rather than surgical procedures outside the realm of childbirth. Monitoring fetal heart rates is indeed a vital part of care, but it is just one aspect of the overall management of labor and does not encapsulate the full role of the obstetrician in this context.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://obcapstone.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE