

OCCLUSION & TMD EXAM 1 PRACTICE (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE

<https://occlusionandtmd1.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. What is the rationale for treating disc displacement without reduction?**
 - A. Educate the patient about the disorder and retrodiscal tissue adaptation.
 - B. Reduce joint loading with soft foods and small bites.
 - C. Stabilization splint at night with follow-up in 3 weeks; adaptation 8-10 weeks.
 - D. NSAIDs if inflammation suspected.
- 2. For referral pain identification, how many seconds should constant pressure be applied?**
 - A. 10 seconds
 - B. 3 seconds
 - C. 15 seconds
 - D. 5 seconds
- 3. Which statement best explains why occlusion alone does not predict TMD?**
 - A. Occlusion alone predicts TMD
 - B. Occlusion is not involved in TMD
 - C. Many factors contribute to TMD
 - D. TMD is solely a joint problem
- 4. Loading forces on the retrodiscal tissue are perceived as what?**
 - A. Pain in the innervated area.
 - B. Numbness unrelated to pain.
 - C. Tinnitus.
 - D. Altered taste.
- 5. What does incorrect facebow mounting misrepresent and what can it lead to?**
 - A. Occlusal plane and condylar relationship are unaffected.
 - B. It records only vertical dimension.
 - C. Incorrect facebow mounting misrepresents the occlusal plane/condylar relationship and can lead to occlusal errors.
 - D. There is no impact on restorative outcomes.

- 6. After the lower incisors pass the upper incisors, the mandible moves upward until posterior teeth contact. In which direction does the movement occur?**
- A. Downward
 - B. Forward
 - C. Upward
 - D. Rotational movement
- 7. When is manual manipulation most successful for disc displacement without reduction?**
- A. In an acute case, manual manipulation is not recommended.
 - B. In an acute case, manual manipulation to recapture the disc is most successful within about 1 week or less.
 - C. Chronic locking has a higher chance of successful recapture.
 - D. Manual manipulation is never successful.
- 8. Maxillary impression steps: Step 2?**
- A. Seat the patient upright and position yourself to the side/back.
 - B. Retract the patient's right cheek and rotate the filled tray into the mouth.
 - C. If material flows toward the oropharynx, gently move the patient's head forward to reduce gagging.
 - D. Hold until alginate sets.
- 9. A restricted mouth opening less than 40 mm is most consistent with which problem area?**
- A. Gingival inflammation.
 - B. TMJ and/or muscle problem.
 - C. Tooth caries.
 - D. Maxillary sinusitis.
- 10. Muscle spindles are located in which muscles and what do they sense?**
- A. Located in elevator muscles; sense changes in muscle length.
 - B. Located in periodontal ligament; sense pressure.
 - C. Located in jaw joint capsule; sense joint movement.
 - D. Located in tongue muscles; sense taste.

Answers

SAMPLE

1. C
2. D
3. C
4. A
5. C
6. C
7. B
8. B
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. What is the rationale for treating disc displacement without reduction?

- A. Educate the patient about the disorder and retrodiscal tissue adaptation.
- B. Reduce joint loading with soft foods and small bites.
- C. Stabilization splint at night with follow-up in 3 weeks; adaptation 8-10 weeks.**
- D. NSAIDs if inflammation suspected.

Disc displacement without reduction is best addressed by giving the jaw a stable, repeatable position that reduces abnormal loading and allows the joint's soft tissues to adapt. Using a stabilization splint worn at night creates an orthopedically favorable bite, lowers muscle hyperactivity, and distributes forces away from the retrodiscal tissues and the anteriorly displaced disc. This creates a protected environment where the disc-condyle relationship can remodel toward a more stable arrangement, while symptoms such as limited opening and pain improve. The plan also includes a short follow-up to assess response and compliance, and an extended adaptation period (about eight to ten weeks) because tissue remodeling and reorganization of the disc-condyle complex take time. This approach is preferred because it directly targets the mechanical problem with a reversible, conservative method that facilitates true functional adaptation, rather than merely addressing symptoms or temporarily reducing loading without promoting long-term structural change.

2. For referral pain identification, how many seconds should constant pressure be applied?

- A. 10 seconds
- B. 3 seconds
- C. 15 seconds
- D. 5 seconds**

During referral pain identification, you want to apply constant pressure for a short, fixed period on suspected trigger points to elicit any referred pain. Five seconds is the optimal duration because it gives enough time for a referral pattern to appear if the trigger point is involved, without causing excessive discomfort or guarding that could blur the findings. Shorter pressures may miss referral pain, while longer pressures can fatigue the patient, increase local pain, or produce non-specific responses. Holding for five seconds and then checking where the patient feels pain helps distinguish myofascial referral from a purely dental source in TMD assessment.

3. Which statement best explains why occlusion alone does not predict TMD?

- A. Occlusion alone predicts TMD
- B. Occlusion is not involved in TMD
- C. Many factors contribute to TMD**
- D. TMD is solely a joint problem

TMD is a multifactorial condition. Pain and dysfunction arise from a mix of muscle function, joint health, neurology of pain processing, sleep-related behaviors, stress, parafunctional habits, and even trauma or systemic factors. Because of this complexity, occlusion alone cannot reliably predict who will develop TMD or how it will present. You can see significant occlusal differences in people with no TMD symptoms and, conversely, people with TMD who have only minor or no notable occlusal issues. That variability shows that bite alignment is only one piece of the puzzle, not the sole driver. In practice, this means evaluating and addressing muscular function, joint integrity, parafunctions, sleep quality, and psychosocial factors in addition to any occlusal considerations. Choices that imply occlusion alone predicts TMD, or that occlusion is not involved, or that it's exclusively a joint problem, don't fit the broader, multifactorial nature of the disorder.

4. Loading forces on the retrodiscal tissue are perceived as what?

- A. Pain in the innervated area.
- B. Numbness unrelated to pain.
- C. Tinnitus.
- D. Altered taste.

Loading forces on the retrodiscal tissue are perceived as pain in the innervated area. The retrodiscal pad behind the disc is rich in nerve endings, so when it's compressed or stressed by bite forces or joint movement, nociceptors send pain signals that the brain interprets as pain in the region supplied by those nerves (around the preauricular/temporal area). This pain response is not about numbness (loss of sensation), tinnitus (ringing in the ears), or altered taste, which involve different pathways and structures. So the brain's perception follows the sensory nerves that innervate that tissue, making pain in the innervated area the correct description.

5. What does incorrect facebow mounting misrepresent and what can it lead to?

- A. Occlusal plane and condylar relationship are unaffected.
- B. It records only vertical dimension.
- C. Incorrect facebow mounting misrepresents the occlusal plane/condylar relationship and can lead to occlusal errors.
- D. There is no impact on restorative outcomes.

Facebow transfer ties the maxillary occlusal plane to the articulator and reproduces the patient's condylar guidance on that instrument. When mounting is off, that spatial relationship is misrepresented on the articulator, so the restorations you fabricate are aligned to a jaw path that doesn't match the patient's real movements. That misalignment can shift the occlusal plane relative to the condylar guidance and distort the condylar relationship on the model, leading to occlusal errors such as improper contacts, premature posterior contacts, and inadequate anterior guidance or posterior disclusion. It's not about vertical dimension—the facebow's purpose isn't to record height—but about correctly reproducing how the jaw moves in space. This misrepresentation directly impacts restorative outcomes, causing interferences and instability that may require adjustments or remakes.

6. After the lower incisors pass the upper incisors, the mandible moves upward until posterior teeth contact. In which direction does the movement occur?

- A. Downward
- B. Forward
- C. Upward
- D. Rotational movement

In protrusive jaw movement, the mandible translates forward under incisal guidance, and then needs to rise to bring the back teeth into contact. Once the lower incisors pass the uppers, the vertical component becomes essential: the posterior teeth occlude only after the mandible elevates toward the maxilla. So the movement is upward as the jaw closes onto the posterior teeth. Downward or forward alone wouldn't achieve posterior contact, and rotational motion alone doesn't capture the necessary vertical lift to reach the occlusion.

7. When is manual manipulation most successful for disc displacement without reduction?

- A. In an acute case, manual manipulation is not recommended.
- B. In an acute case, manual manipulation to recapture the disc is most successful within about 1 week or less.**
- C. Chronic locking has a higher chance of successful recapture.
- D. Manual manipulation is never successful.

The key idea is timing. Manual manipulation to recapture a disc displaced without reduction works best when the problem is in its acute phase, typically within about a week of onset. In this early window, the surrounding tissues are more compliant and inflammation hasn't yet caused dense adhesions or fibrosis. That makes it possible to gently guide the condyle and reposition the disc back onto the fossa, resolving the locked bite and restoring functional movement. As time passes and the condition becomes chronic, the retrodiscal tissues and surrounding joint structures tend to stiffen with fibrosis and adhesions. Those changes make recapturing the disc much harder with non-surgical manipulation, reducing success rates. So the statement that acute onset within about a week offers the best chance aligns with the biology of tissue plasticity and remodeling in TMD. The other options don't fit because they contradict the reality that early intervention improves success, while chronic cases are less amenable to recapture and improvable with manipulation.

8. Maxillary impression steps: Step 2?

- A. Seat the patient upright and position yourself to the side/back.
- B. Retract the patient's right cheek and rotate the filled tray into the mouth.**
- C. If material flows toward the oropharynx, gently move the patient's head forward to reduce gagging.
- D. Hold until alginate sets.

The action being tested is the correct technique for inserting a loaded maxillary impression tray. Retracting the cheek on the chosen side creates a clear corridor in the vestibule, preventing soft tissue from catching the tray as you bring it into the mouth. Rotating the filled tray into the mouth from the side helps you seat the anterior edge first and then bring the tray onto the palate and posterior teeth, promoting even contact and full coverage while minimizing distortion or air pockets. This controlled, side-entry maneuver sets up a successful impression by ensuring the tray seats smoothly and securely. Other steps described—adjusting general chair position, managing gagging by moving the head forward, or waiting for the material to set—pertain to different aspects of the procedure and do not describe the insertion technique focused on proper cheek retraction and orientation of the tray.

9. A restricted mouth opening less than 40 mm is most consistent with which problem area?

- A. Gingival inflammation.
- B. TMJ and/or muscle problem.**
- C. Tooth caries.
- D. Maxillary sinusitis.

Restricted mouth opening points to problems in the jaw joints or the muscles that move the jaw. A measured opening under 40 mm is a classic sign of a TMJ (temporomandibular joint) issue or a related masticatory muscle disorder, such as muscle spasm or inflammation limiting movement. Normal active mouth opening in adults is typically about 40–50 mm, so falling below 40 mm indicates a mechanical or muscular restriction rather than a dental or sinus issue. Gingival inflammation mainly causes gums problems and pain with brushing or chewing, not a restricted jaw opening. Tooth caries causes localized toothache or sensitivity but usually does not limit how wide you can open your mouth. Maxillary sinusitis can cause facial aching and pressure but does not typically produce a measurable restriction of jaw opening like this.

10. Muscle spindles are located in which muscles and what do they sense?

- A. Located in elevator muscles; sense changes in muscle length.**
- B. Located in periodontal ligament; sense pressure.
- C. Located in jaw joint capsule; sense joint movement.
- D. Located in tongue muscles; sense taste.

Muscle spindles are stretch receptors embedded in skeletal muscles, including the elevator muscles of the jaw, and they sense changes in muscle length (and how quickly that length is changing). When a muscle is stretched, the spindles send signals to the spinal cord to trigger a reflex that resists the stretch, helping regulate muscle tone and coordinate jaw movements. This proprioceptive feedback is crucial for controlled bite and jaw position. The other options mix in receptors that aren't muscle spindles: the periodontal ligament has mechanoreceptors that detect pressure from tooth contact; the jaw joint capsule has receptors that monitor joint movement and position; and taste is detected by taste buds in the tongue, not by muscle spindles.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://occlusionandtmd1.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE