ONTARIO PARAMEDIC PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://ontarioparamedic.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is a key step in performing a rapid trauma assessment?

A. Consulting with the physician beforehand

B. Immediately begin administering medication

C. Quickly evaluate the patient for life-threatening injuries

D. Focus only on the patient's complaints

2. Why is it important to obtain a patient's consent before treatment?

A. To ensure patient satisfaction

B. To respect the patient's autonomy and rights

C. To comply with legal regulations

D. To avoid liability

3. In pediatric patients, what is the recommended compression-to-ventilation ratio for CPR?

A. 30:2

B. 15:2

C. 10:1

D. 20:2

4. What are the indications to a traumatic cardiac arrest?

A. Cardiac arrest due to drowning

B. Cardiac arrest secondary to drug overdose

C. Cardiac arrest secondary to severe blunt or penetrating trauma

D. Cardiac arrest due to hypothermia

5. Why is it important to determine the mechanism of injury in trauma patients?

A. To categorize patients by age

B. To assess potential internal injuries

C. To calculate transportation times

D. To review the patient's insurance details

SAMPLE

6. What is the correct dose of epinephrine for a severe allergic reaction in adults?

A. 0.1 to 0.2 mg

B. 0.3 to 0.5 mg

C. 1.0 to 1.5 mg

D. 0.5 to 0.7 mg

7. Which of the following conditions is considered active bleeding?

A. Internal bleeding from a controlled peptic ulcer

B. External trauma that has been dressed and controlled

C. Intermittent bleeding from renal colic

D. Bleeding gums after dental surgery

8. When assessing a patient with chest pain, what vital sign is critical to monitor?

A. Temperature

B. Blood pressure

C. Pulse oximetry

D. Respiratory rate

9. What is the dosage of Acetaminophen for a person aged 12 years to less than 18?

A. PO 960-1000 one time

B. PO 500-650 one time

C. PO 400mg once only

D. PO 800mg once only

10. What is the max number of doses for a traumatic defib?

A. 2

B. 3

C. 1

D. 4

SAMPLE

Answers

SAMPLE

1. C
2. B
3. B
4. C
5. B
6. B
7. D
8. B
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. What is a key step in performing a rapid trauma assessment?

A. Consulting with the physician beforehand

B. Immediately begin administering medication

C. Quickly evaluate the patient for life-threatening injuries

D. Focus only on the patient's complaints

A key step in performing a rapid trauma assessment is to quickly evaluate the patient for life-threatening injuries. This initial assessment is crucial in emergency situations where time is of the essence, as it allows the paramedic to identify and address any conditions that could lead to severe complications or death. The primary goal is to ensure patient safety by focusing on critical areas such as airway patency, breathing effectiveness, circulation status, and any immediate threats to life. By prioritizing life-threatening injuries in a systematic and swift manner, paramedics can implement necessary interventions, such as securing the airway, controlling bleeding, or initiating cardiopulmonary resuscitation if needed. This assessment lays the groundwork for further care and stabilization of the patient. Other potential actions, such as consulting with a physician beforehand or immediately administering medication, are not appropriate during the rapid assessment phase. This phase requires swift evaluation and action rather than waiting for additional input or commencing treatment that may not yet be necessary. Focusing solely on the patient's complaints would also be inadequate because it could overlook critical injuries not directly expressed by the patient, hence compromising their safety and care.

2. Why is it important to obtain a patient's consent before treatment?

A. To ensure patient satisfaction

B. To respect the patient's autonomy and rights

C. To comply with legal regulations

D. To avoid liability

Obtaining a patient's consent before treatment is crucial because it respects the patient's autonomy and rights. Autonomy is a fundamental principle in healthcare that acknowledges an individual's right to make informed decisions regarding their own body and treatment options. By securing consent, healthcare providers empower patients to participate actively in their care, ensuring that they understand the proposed interventions, benefits, risks, and alternatives. This process goes beyond simply getting permission; it affirms the respect for the individual's right to choose or refuse treatment, which is essential in establishing a trusting and collaborative relationship between the patient and the healthcare provider. Informed consent also reflects ethical standards in medicine, promoting the idea that patients should have control over their own health decisions—aligning with the values of dignity and respect that underpin patient-centered care. While other options touch on important aspects of patient care, such as compliance with legal statutes and the protection from liability, the primary focus of consent is the respect for patient autonomy, making it a cornerstone of ethical healthcare practice.

SAMPLE

3. In pediatric patients, what is the recommended compression-to-ventilation ratio for CPR?

A. 30:2

B. 15:2

C. 10:1

D. 20:2

In pediatric patients, the recommended compression-to-ventilation ratio for CPR is 15:2. This ratio applies specifically when two rescuers are present, ensuring that adequate chest compressions are delivered alongside effective ventilations. The rationale behind this specific ratio is rooted in the developmental physiology of children. Children have a higher respiratory rate and a higher demand for oxygen than adults, which makes effective ventilation critical during resuscitation efforts. Using a 15:2 ratio allows for more frequent ventilations compared to adults, where a ratio of 30:2 is standard. This emphasis on providing more breaths is particularly important for pediatric patients, who are more prone to respiratory issues that can lead to cardiac arrest. The focus on a higher number of compressions in this configuration also helps maintain adequate blood circulation while oxygenation is prioritized. In a practical scenario, having two rescuers allows for one to focus on delivering efficient chest compressions, while the second can ensure that ventilations are provided without prolonged interruptions. This structured approach increases the chances of successful resuscitation in pediatric emergencies.

4. What are the indications to a traumatic cardiac arrest?

A. Cardiac arrest due to drowning

B. Cardiac arrest secondary to drug overdose

C. Cardiac arrest secondary to severe blunt or penetrating trauma

D. Cardiac arrest due to hypothermia

The correct indication for a traumatic cardiac arrest is related to cardiac arrest secondary to severe blunt or penetrating trauma. This is because traumatic cardiac arrest is specifically defined as a loss of heart function due to significant injuries from trauma, such as those sustained in accidents, falls, or assaults. The physiological effects of severe trauma can lead to loss of critical hemodynamic stability, resulting in cardiac arrest. Traumatic cardiac arrest often necessitates rapid assessment and intervention, as these patients may still have a chance of survival if the underlying cause, such as bleeding, can be addressed quickly. A thorough understanding of this condition is crucial for emergency responders to initiate appropriate therapies promptly, including advanced resuscitative techniques and potential surgical interventions. In contrast, other conditions listed, such as cardiac arrest due to drowning, drug overdose, or hypothermia, are not classified as traumatic cardiac arrests but rather as non-traumatic causes of cardiac arrest. These situations involve different underlying pathophysiological mechanisms and management approaches.

SAMPLE

5. Why is it important to determine the mechanism of injury in trauma patients?

A. To categorize patients by age

B. To assess potential internal injuries

C. To calculate transportation times

D. To review the patient's insurance details

Determining the mechanism of injury in trauma patients is crucial primarily because it aids in assessing potential internal injuries. Understanding how an injury occurred provides valuable insight into the forces involved, which can help predict the types of injuries that may not be immediately apparent. For example, a high-speed motor vehicle collision or a fall from a significant height often indicates a higher likelihood of internal organ damage or fractures that may not be visible externally. Recognizing the mechanism of injury allows paramedics and healthcare providers to prioritize assessments, guide imaging and interventions, and develop an appropriate treatment plan. This understanding also supports the development of a comprehensive history to communicate effectively with other medical personnel during transport and in the emergency department, ensuring that appropriate care is initiated promptly. In contrast, categorizing patients by age, calculating transportation times, and reviewing insurance details may be relevant to patient management but do not have the same immediate and critical relevance to identifying and addressing potential life-threatening injuries as understanding the mechanism of injury does.

6. What is the correct dose of epinephrine for a severe allergic reaction in adults?

A. 0.1 to 0.2 mg

B. 0.3 to 0.5 mg

C. 1.0 to 1.5 mg

D. 0.5 to 0.7 mg

For a severe allergic reaction, commonly known as anaphylaxis, the correct dose of epinephrine for adults is typically between 0.3 to 0.5 mg. This dosage is effective in counteracting the severe symptoms associated with anaphylaxis, such as airway constriction and cardiovascular instability. Administering epinephrine helps to rapidly reverse the effects of the allergic reaction by stimulating alpha and beta-adrenergic receptors, leading to vasoconstriction, bronchodilation, and increased heart rate, among other physiological responses. In the event of anaphylaxis, it is important to administer the epinephrine intramuscularly, usually in the mid-anterolateral thigh, for optimal absorption and efficacy. The 0.3 to 0.5 mg range is the most widely accepted and recommended dosage in clinical practice and guidelines for adults experiencing a severe allergic reaction. This dose may be repeated every 5 to 15 minutes if symptoms persist, depending on the severity of the reaction and the clinical response of the patient. Dosing outside of the recommended range, such as doses lower than 0.3 mg or higher than 0.5 mg, may not provide the necessary impact or could lead to adverse effects

7. Which of the following conditions is considered active bleeding?

A. Internal bleeding from a controlled peptic ulcer

B. External trauma that has been dressed and controlled

C. Intermittent bleeding from renal colic

D. Bleeding gums after dental surgery

Active bleeding refers to ongoing or continuous bleeding that has not been stopped or controlled. In this case, bleeding gums after dental surgery are considered active bleeding because the gums are still bleeding and have not been stopped from bleeding by a dressing or other means. Options A, B, and C all indicate bleeding that has been controlled or stopped, thus they are not considered active bleeding.

SAMPLE

8. When assessing a patient with chest pain, what vital sign is critical to monitor?

A. Temperature

B. Blood pressure

C. Pulse oximetry

D. Respiratory rate

Monitoring blood pressure is crucial when assessing a patient with chest pain because it provides essential information about the patient's cardiovascular status. Changes in blood pressure can indicate the severity of the underlying condition, such as a myocardial infarction or other cardiac events. A significant drop in blood pressure may suggest hypoperfusion or shock, which requires immediate intervention. Conversely, elevated blood pressure can indicate increased stress on the heart or the presence of conditions such as hypertension. While monitoring temperature, pulse oximetry, and respiratory rate can offer valuable insight into the patient's overall health and other potential complications, blood pressure directly reflects the status of the heart and circulatory system. In the context of chest pain, this vital sign is particularly telling and can guide critical treatment decisions, making it the most important parameter to watch closely during the assessment.

9. What is the dosage of Acetaminophen for a person aged 12 years to less than 18?

A. PO 960-1000 one time

B. PO 500-650 one time

C. PO 400mg once only

D. PO 800mg once only

The appropriate dosage of Acetaminophen for a person aged 12 years to less than 18 is generally in the range of 500-650 mg taken by mouth (PO) once. This dosage is consistent with standard guidelines for pediatric patients, ensuring effective relief from pain or fever while minimizing the risk of toxicity. The rationale for this dosage range takes into account both safety and efficacy, as Acetaminophen is widely used for its analgesic and antipyretic properties. While there are other dosages listed, options that significantly exceed this range could lead to an increased risk of adverse effects. It is also important to ensure that the dosing aligns with current recommendations on maximum daily intake and avoids exceeding recommended limits for this age group. Thus, the choice of 500-650 mg once is suitable for addressing the needs of young adolescents safely and effectively.

10. What is the max number of doses for a traumatic defib?

A. 2

B. 3

C. 1

D. 4

In the context of traumatic defibrillation, the maximum number of doses typically administered during a resuscitation attempt is one. This is because traumatic cardiac arrest often presents with significant underlying injuries, and the effectiveness of defibrillation is greatly influenced by the nature of those injuries. If the initial defibrillation attempt is unsuccessful, paramedics will frequently prioritize other life-saving measures such as airway management, control of hemorrhage, and volume resuscitation rather than continuing with additional shocks. This approach aligns with established protocols that guide advanced cardiac life support in traumatic scenarios, emphasizing the need for rapid assessment and intervention to address the causes of arrest rather than multiple defibrillation attempts. Thus, recognizing that the initial shock is crucial, the maximum number of doses for defibrillation in this context is indeed one.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://ontarioparamedic.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE