
ORAL AND MAXILLOFACIAL SURGERY (OMFS) BOARD PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://omfsboard.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Hamman's sign is associated with which condition?

A. Spontaneous pneumothorax

B. Spontaneous mediastinal emphysema

C. Pneumonia

D. Chronic lung disease

2. What level of the skin does a moderate chemical peel treat?

A. Superficial epidermis

B. Upper reticular dermis

C. Mid-reticular dermis

D. Lower dermis

3. Why is it not advisable to use a trapezius flap after a radical neck dissection?

A. Damage to spinal accessory nerve

B. Transverse cervical artery may have been sacrificed

C. Salvage procedure complications

D. Limited vascular supply

4. What is the onset time for the action of Pradaxa?

A. Immediate

B. Within hours

C. 24 hours

D. Several days

5. What is the preferred resuscitation fluid for patients with acute brain injury?

A. Ringer's lactate

B. 0.9% saline

C. Hypertonic saline

D. Crystalloids

SAMPLE

6. How is pain typically perceived when there is a disc perforation in the TMJ?

A. Peripheral nociceptors

B. Subchondral nociceptors

C. Central nervous system receptors

D. Visceral nociceptors

7. What Glasgow Coma Scale score range reflects a moderate injury?

A. 3-8

B. 9-12

C. 13-15

D. 16-20

8. What is the maximum amount of bone that can be harvested from the posterior iliac crest?

A. 75 cc

B. 100 cc

C. 125 cc

D. 150 cc

9. Which type of skin structure survives when harvesting a dermis graft?

A. Hair follicles

B. Sweat glands

C. Sebaceous glands

D. Blood vessels

10. Which cranial nerve provides taste sensation to the anterior two-thirds of the tongue?

A. Glossopharyngeal nerve (CN IX)

B. Facial nerve (CN VII)

C. Vagus nerve (CN X)

D. Trigeminal nerve (CN V)

SAMPLE

Answers

SAMPLE

1. B
2. B
3. B
4. B
5. B
6. B
7. B
8. B
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. Hamman's sign is associated with which condition?

A. Spontaneous pneumothorax

B. Spontaneous mediastinal emphysema

C. Pneumonia

D. Chronic lung disease

Hamman's sign is a clinical finding characterized by a crackling sound that can be heard over the precordium or chest during auscultation when the patient is in a supine position. This sign is specifically associated with spontaneous mediastinal emphysema, a condition that occurs when air leaks into the mediastinum, often due to rupture of air-containing structures such as the alveoli or bronchi. The presence of Hamman's sign suggests that there is air in the mediastinum, which can create an audible sound with each heartbeat, similar to a crunching or popping noise. This finding is associated with conditions like those that cause sudden increases in intrathoracic pressure or trauma, and thus, it is a key clinical indicator for mediastinal emphysema. In contrast, spontaneous pneumothorax, pneumonia, or chronic lung disease may involve respiratory symptoms and findings but do not typically produce Hamman's sign. In spontaneous pneumothorax, you might hear decreased breath sounds or hyper-resonance on percussion, while pneumonia typically presents with dullness, crackles, or decreased breath sounds related to lung consolidation. Chronic lung disease often manifests in chronic cough or wheezing but does not relate directly to Hamman's sign. Thus

2. What level of the skin does a moderate chemical peel treat?

A. Superficial epidermis

B. Upper reticular dermis

C. Mid-reticular dermis

D. Lower dermis

A moderate chemical peel primarily targets the upper reticular dermis, along with the deeper layers of the epidermis. This level of peel is typically achieved using ingredients such as glycolic acid, trichloroacetic acid (TCA), or other agents that penetrate deeper than superficial peels, which only affect the outermost layers of the skin. By reaching the upper reticular dermis, moderate chemical peels help in improving skin texture, addressing pigmentation irregularities, and promoting collagen remodeling. The depth of penetration allows for more significant changes in the skin compared to superficial peels, making moderate peels effective for treating issues like fine lines, moderate photoaging, and superficial scars. Understanding the depth of the peel is important for determining the expected outcomes, recovery times, and potential side effects associated with the procedure. The other layers mentioned do not accurately represent the level of treatment provided by a moderate chemical peel, which is an essential aspect of its efficacy and safety profile.

SAMPLE

3. Why is it not advisable to use a trapezius flap after a radical neck dissection?

A. Damage to spinal accessory nerve

B. Transverse cervical artery may have been sacrificed

C. Salvage procedure complications

D. Limited vascular supply

The trapezius flap is a surgical technique often used for reconstructive purposes in the head and neck region. However, its use post-radical neck dissection is limited due to potential complications associated with vascular supply, particularly the preservation of the transverse cervical artery. In a radical neck dissection, the transverse cervical artery may be sacrificed along with other structures such as lymph nodes and surrounding tissue. This artery is crucial for supplying blood to the trapezius muscle. Without adequate vascular supply, the viability of the trapezius flap can be compromised, potentially leading to flap necrosis and failure. Moreover, a sacrificed transverse cervical artery means that even if the trapezius muscle is viable at the time of transfer, it may not receive enough blood flow post-operatively due to its reliance on this artery for nutrition. Therefore, the flaps may not heal properly, and the overall success of the reconstructive procedure can be adversely affected. Understanding the role of blood supply in surgical planning highlights why the transverse cervical artery's status is paramount when considering the use of a trapezius flap following a radical neck dissection.

4. What is the onset time for the action of Pradaxa?

A. Immediate

B. Within hours

C. 24 hours

D. Several days

The onset time for the action of Pradaxa, which is an oral anticoagulant containing dabigatran etexilate, is indeed within hours. Pradaxa works as a direct thrombin inhibitor, meaning it directly affects the blood clotting process. After ingestion, the pro-drug dabigatran etexilate is rapidly absorbed and converted into its active form, dabigatran, which has an onset of action typically between one to four hours after administration. This rapid onset is particularly beneficial in clinical settings where quick anticoagulation is necessary, such as in the prevention of thromboembolic events in patients with atrial fibrillation. Understanding the pharmacokinetics of Pradaxa is crucial, as it allows healthcare providers to make informed decisions about patient management, especially in terms of timing for procedures and potential bleeding risks. Choices suggesting a longer onset timeframe, such as 24 hours or several days, do not align with the pharmacological profile of Pradaxa. An immediate effect is also not the case, as there is a slight delay in the time required for the drug to reach therapeutic levels and exert its effect.

SAMPLE

5. What is the preferred resuscitation fluid for patients with acute brain injury?

A. Ringer's lactate

B. 0.9% saline

C. Hypertonic saline

D. Crystalloids

In the context of managing patients with acute brain injury, the preferred resuscitation fluid is 0.9% saline. This choice is based on the need to maintain adequate cerebral perfusion pressure and to avoid potential complications that can arise from other fluid types. 0.9% saline, also known as normal saline, is isotonic and maintains a normal serum sodium level, which is crucial in patients with brain injuries to prevent the risk of cerebral edema. It aids in providing volume resuscitation effectively and is typically well-tolerated in clinical settings. Moreover, its pH is neutral, minimizing the risk of aggravating any acidosis that might accompany severe head injuries. In comparison, although Ringer's lactate and crystalloids are useful in resuscitation for various medical conditions, they may influence acid-base balance and electrolyte levels in a manner that could complicate the management of head injuries. Hypertonic saline, while beneficial in specific scenarios like managing severe intracranial hypertension, is not routinely used as the first-line fluid for general resuscitation due to its somewhat complex effects on fluid dynamics and cerebral edema. Thus, the use of 0.9% saline offers a straightforward and effective approach to fluid resuscitation in

6. How is pain typically perceived when there is a disc perforation in the TMJ?

A. Peripheral nociceptors

B. Subchondral nociceptors

C. Central nervous system receptors

D. Visceral nociceptors

In the context of disc perforation in the temporomandibular joint (TMJ), pain is primarily perceived through subchondral nociceptors. These nociceptors are located within the subchondral bone beneath the cartilage and are activated in response to inflammatory changes or mechanical stress associated with joint pathology. When a disc perforation occurs, it often leads to changes in the biomechanics of the TMJ, resulting in increased stress on the joint surfaces and the underlying bone. This can stimulate the subchondral nociceptors, leading to the sensation of pain. Additionally, inflammatory mediators released in response to joint damage can further sensitize these nociceptors, enhancing the perception of pain in the affected area. While peripheral nociceptors do contribute to pain perception in some contexts, in TMJ disorders specifically, the subchondral nociceptors are more directly involved when there is a structural issue like disc perforation. Central nervous system receptors play a role in processing pain signals but are not the primary source for the initial pain perception at the site of the injury. Visceral nociceptors, which respond to pain from internal organs, are not relevant in the context of TMJ function

SAMPLE

7. What Glasgow Coma Scale score range reflects a moderate injury?

A. 3-8

B. 9-12

C. 13-15

D. 16-20

The Glasgow Coma Scale (GCS) is a neurological scale used to assess the level of consciousness in a person following a traumatic brain injury. The scale consists of three components: eye opening, verbal response, and motor response, each of which is scored individually. The total score can range from 3 (indicating deep unconsciousness) to 15 (indicating full consciousness). A score of 9 to 12 on the GCS indicates a moderate head injury. This range suggests that the patient is not fully alert and exhibits some degree of impaired consciousness, which may include confusion and a decreased ability to respond to external stimuli. Such patients typically require closer monitoring and may benefit from medical intervention to address the underlying causes of their impaired consciousness. In contrast, lower scores (3-8) indicate severe brain injury with a significantly reduced level of responsiveness, while higher scores (13-15) reflect mild brain injury or a fully alert state. A score of 16-20 is not applicable, as the maximum score on the Glasgow Coma Scale is 15. Thus, the score range of 9-12 is clearly representative of moderate injury, necessitating appropriate medical assessment and management.

8. What is the maximum amount of bone that can be harvested from the posterior iliac crest?

A. 75 cc

B. 100 cc

C. 125 cc

D. 150 cc

Harvesting bone from the posterior iliac crest is a common procedure in oral and maxillofacial surgery, particularly for grafting purposes. The amount of bone that can be harvested depends on several factors, including the patient's anatomy and the surgical technique employed. The maximum volume that is typically considered safe and feasible to harvest from the posterior iliac crest is around 100 cc. This amount is aligned with reported data in the literature, making it a practical standard for both surgeons and patients. Exceeding this volume can increase the risk of complications, such as hematoma formation, infection, and damage to surrounding structures. Options that suggest greater amounts, such as 125 cc or 150 cc, exceed the generally accepted limits and pose a higher risk of complications, while those suggesting lower amounts, such as 75 cc, may underestimate the potential for harvesting bone effectively in most patients. Hence, the choice of 100 cc reflects a balance between sufficient grafting material and minimizing patient risk.

SAMPLE

9. Which type of skin structure survives when harvesting a dermis graft?

A. Hair follicles

B. Sweat glands

C. Sebaceous glands

D. Blood vessels

When harvesting a dermis graft, sebaceous glands are among the skin structures that are likely to survive. Dermis grafts are primarily composed of the dermal layer of the skin, which contains these glands. Sebaceous glands are responsible for producing oil that helps to keep the skin moisturized and are situated within the dermis. On the other hand, hair follicles typically do not survive in dermal grafts because they reside in the deeper layers that may not be preserved when only the dermis is harvested. Sweat glands, while located in the dermis, are more sensitive to the harvesting process, and therefore, their survival is not guaranteed. Blood vessels, although crucial for graft viability, do not remain intact as independent structures in the graft; rather, for a graft to survive, revascularization from the recipient site must occur post-operation. Understanding the nature of each structure and its relationship to the dermal layer is essential for comprehending which components can endure the harvesting process. The presence of sebaceous glands within the harvested dermal graft contributes to its potential for integration and functionality when transplanted.

10. Which cranial nerve provides taste sensation to the anterior two-thirds of the tongue?

A. Glossopharyngeal nerve (CN IX)

B. Facial nerve (CN VII)

C. Vagus nerve (CN X)

D. Trigeminal nerve (CN V)

The facial nerve, also known as cranial nerve VII, is responsible for providing taste sensation to the anterior two-thirds of the tongue. The taste buds in this region send their signals through the chorda tympani branch of the facial nerve, which then relays the taste information to the brain. This function is distinct from other cranial nerves listed. The glossopharyngeal nerve (CN IX) primarily provides taste sensation to the posterior one-third of the tongue. The vagus nerve (CN X) is involved in taste sensation from the epiglottis and the lower pharynx, while the trigeminal nerve (CN V) primarily handles general sensory information such as touch, pain, and temperature from the tongue rather than taste. Understanding these specific functions of each cranial nerve helps clarify why the facial nerve is correctly associated with taste in the anterior two-thirds of the tongue.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://omfsboard.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE