
ORANGE COUNTY FLORIDA EMERGENCY MEDICAL SERVICES (EMS) PROTOCOL PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://orangecountyflemsprotocol.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. When should you reassess a patient's ABCs?

A. Only when the patient arrives at the hospital

B. After each vital sign check

C. Anytime there is a change in the patient's condition or after any intervention

D. Once every hour

2. During CPR, where should the hands be positioned for effective chest compressions?

A. On the stomach

B. On the center of the chest, interlocked

C. On the lower ribs

D. On the shoulders

3. What should always accompany the chief complaint in a handoff report?

A. The patient's symptoms

B. The treatment given

C. The patient's age

D. The time of incident

4. Which age group requires the consideration of child-specific protocols in EMS?

A. Adolescents (13-19 years)

B. Adults (20-64 years)

C. Pediatric patients (infants and children)

D. Geriatric patients (65 years and older)

5. What is required before administering epinephrine to a patient in respiratory distress over 55 years old?

A. Doctor's consent

B. Medical director's permission

C. Patient's written consent

D. Confirmation of symptoms

SAMPLE

6. What condition might lead to the administration of good chest compressions without interruptions?

A. Asthma attack

B. Pulseless electrical activity (PEA)

C. Anaphylactic shock

D. Severe pneumonia

7. How often should CPR be reassessed in a patient?

A. Every 10 minutes

B. Every two minutes or after every 5 cycles of 30 compressions and 2 breaths

C. Every 5 minutes regardless of condition

D. Only after the ambulance arrives

8. Which condition is a contraindication for administering nitroglycerin?

A. Hypertension (high blood pressure)

B. Hypotension (low blood pressure)

C. Chest pain

D. Diabetes

9. What vital signs are critical for assessing a patient's hemodynamic status?

A. Temperature, heart rate, oxygen saturation, and blood pressure

B. Heart rate, blood pressure, respiratory rate, and oxygen saturation

C. Blood glucose, heart rate, respiratory rate, and blood pressure

D. Heart rate, temperature, blood pressure, and level of consciousness

10. What should be done if a patient has an obstructed airway?

A. Administer medication immediately

B. Perform abdominal thrusts for adults or back blows for infants

C. Call for advanced medical support only

D. Encourage the patient to cough forcefully

SAMPLE

Answers

SAMPLE

1. C
2. B
3. B
4. C
5. B
6. B
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. When should you reassess a patient's ABCs?

A. Only when the patient arrives at the hospital

B. After each vital sign check

C. Anytime there is a change in the patient's condition or after any intervention

D. Once every hour

Reassessing a patient's ABCs (Airway, Breathing, Circulation) is a critical component of ongoing patient evaluation in emergency medical situations. The correct approach is to conduct this reassessment anytime there is a change in the patient's condition or after any intervention. This ensures that changes in the patient's status are promptly recognized and managed. For instance, if a patient's breathing becomes labored, or if an intervention, such as administering oxygen, is initiated, reassessing the ABCs would allow for immediate evaluation of the effectiveness of the intervention. This practice aligns with the dynamic nature of patient care, where conditions can change rapidly, especially in emergency scenarios. Regularly reassessing ABCs can help identify potentially life-threatening issues early, thereby facilitating timely and necessary interventions. In contrast, waiting until the patient arrives at the hospital or reassessing only after specific intervals (like once every hour) could lead to delays in recognizing deterioration in the patient's condition. Similarly, while checking vital signs is essential, it should not be the only trigger for assessing ABCs, as significant changes can occur at any moment and warrant immediate action.

2. During CPR, where should the hands be positioned for effective chest compressions?

A. On the stomach

B. On the center of the chest, interlocked

C. On the lower ribs

D. On the shoulders

For effective chest compressions during CPR, the hands should be positioned on the center of the chest, interlocked. This position allows for the greatest force to be applied directly to the sternum, facilitating the compression of the heart and improving blood circulation to vital organs. The technique also ensures that compressions are delivered at the correct depth and rate, which is crucial for increasing the chances of survival during cardiac arrest. Interlocking the fingers provides stability and helps prevent fatigue in the rescuer's hands, allowing for sustained compressions over time. Additionally, positioning the hands in the center of the chest avoids potential injury to the ribs or internal organs that could occur if compressions were applied elsewhere, such as on the stomach or lower ribs. Proper technique not only maximizes the effectiveness of CPR but also adheres to established guidelines and protocols.

SAMPLE

3. What should always accompany the chief complaint in a handoff report?

A. The patient's symptoms

B. The treatment given

C. The patient's age

D. The time of incident

The chief complaint is a critical component of a handoff report, as it directly informs the receiving medical personnel about the primary concern that necessitated emergency medical services. However, it is essential to accompany the chief complaint with the treatment given. This information provides context about the interventions that have already been performed, which can significantly influence ongoing care decisions. Understanding what treatment was administered allows the receiving team to gauge the effectiveness of the initial interventions, recognize any possible complications or side effects, and determine the next appropriate steps in the patient's care. It also helps ensure continuity of care, as the new medical team must understand the patient's prior management and response to treatment. Other details, such as the patient's symptoms, age, and time of the incident, while important, do not carry the same immediate implications for treatment continuity and decision making as the treatment previously given.

4. Which age group requires the consideration of child-specific protocols in EMS?

A. Adolescents (13-19 years)

B. Adults (20-64 years)

C. Pediatric patients (infants and children)

D. Geriatric patients (65 years and older)

The consideration of child-specific protocols in EMS is crucial for pediatric patients, which include infants and children. This age group exhibits distinct physiological and developmental characteristics that differ significantly from those of adults. These differences can greatly impact emergency medical care, including drug dosages, assessment techniques, and treatment modalities. In pediatrics, for example, vital signs, normal ranges, and responses to illnesses or trauma can vary widely compared to older populations. Infants and children may also present symptoms that differ from those seen in adults, necessitating a tailored approach to treatment. Child-specific protocols ensure EMS personnel provide appropriate care that recognizes these variations and adapts interventions accordingly. By following these protocols, EMS providers can better ensure positive outcomes for younger patients in emergency situations.

SAMPLE

5. What is required before administering epinephrine to a patient in respiratory distress over 55 years old?

A. Doctor's consent

B. Medical director's permission

C. Patient's written consent

D. Confirmation of symptoms

In cases involving the administration of epinephrine for respiratory distress in patients over 55 years old, obtaining the medical director's permission is crucial due to the potential for underlying health complications that could arise in older patients. This age group often presents with various comorbidities, which necessitate a more cautious approach to medication administration. The involvement of the medical director ensures that the decision to administer epinephrine is not only based on immediate symptoms but also takes into account the patient's overall health status and any protocols specific to older adults. Medical directors typically provide guidelines that reflect evidence-based practice and patient safety considerations, which is particularly important when administering potent medications like epinephrine that can have significant cardiovascular effects. In contrast, options such as doctor's consent or patient's written consent may not necessarily align with established protocols, as the medical director is the designated authority overseeing EMS protocols and medication administration. Confirmation of symptoms, while important, is typically a prerequisite for treatment but does not address the additional layer of oversight required for an older population. Therefore, obtaining the medical director's permission is the appropriate step to ensure compliance with EMS guidelines and to prioritize patient safety.

6. What condition might lead to the administration of good chest compressions without interruptions?

A. Asthma attack

B. Pulseless electrical activity (PEA)

C. Anaphylactic shock

D. Severe pneumonia

The correct answer is based on the understanding of pulseless electrical activity (PEA), which is a specific condition that occurs during cardiac arrest. PEA is characterized by the presence of electrical activity on an electrocardiogram (ECG), but the heart fails to effectively pump blood, resulting in a lack of pulse. In such cases, immediate and effective chest compressions are crucial for maintaining circulation and perfusing vital organs until more definitive intervention can occur, such as the administration of medications or advanced cardiac life support procedures. Continuous chest compressions without interruptions are emphasized in the treatment of PEA because they are vital for enhancing survival outcomes. Interruptions in compressions can lead to decreased blood flow and adversely affect the chances of restoring a perfusing rhythm. In contrast, the other conditions may not necessarily require a constant focus on uninterrupted chest compressions. For instance, during an asthma attack, the primary concern is airway management and bronchodilator therapy, while anaphylactic shock requires immediate administration of epinephrine and may involve airway interventions alongside potential compressions. Severe pneumonia typically calls for supportive care and treatment of infection rather than chest compressions. Therefore, the management of PEA uniquely highlights the importance of delivering effective chest compressions continuously.

SAMPLE

7. How often should CPR be reassessed in a patient?

A. Every 10 minutes

B. Every two minutes or after every 5 cycles of 30 compressions and 2 breaths

C. Every 5 minutes regardless of condition

D. Only after the ambulance arrives

The correct interval for reassessing CPR in a patient is every two minutes or after every five cycles of 30 compressions and 2 breaths. This practice is based on established CPR protocols that emphasize the importance of regular reassessment to ensure that the quality of compressions is maintained and that the patient's condition is being monitored effectively. Reassessing CPR every two minutes allows for the opportunity to check for any changes in the patient's responsiveness and pulse, and it aligns with the rhythm of performing compressions and rescue breaths. Following this rhythm ensures that the provider can adjust their technique based on the patient's response, modifying interventions as necessary based on the evolving situation. Regular reassessment also minimizes the risk of fatigue and helps maintain effective compression rates and depth, which are critical for improving the chances of successful resuscitation. Thus, sticking to the structured intervals ensures adherence to life-saving guidelines and maximizes patient outcomes in cardiac arrest situations.

8. Which condition is a contraindication for administering nitroglycerin?

A. Hypertension (high blood pressure)

B. Hypotension (low blood pressure)

C. Chest pain

D. Diabetes

Hypotension, or low blood pressure, is indeed a contraindication for administering nitroglycerin. Nitroglycerin works by dilating blood vessels, which decreases systemic vascular resistance and lowers blood pressure. If a patient is already experiencing low blood pressure, administering nitroglycerin could exacerbate this condition, potentially leading to severe hypotension and resulting in inadequate perfusion to vital organs. Therefore, it is critical to assess blood pressure before administering nitroglycerin to ensure the safety of the patient. Conditions such as hypertension, chest pain, and diabetes do not inherently prohibit the use of nitroglycerin. In fact, nitroglycerin is often used to relieve chest pain associated with angina or myocardial infarction, provided the patient's blood pressure is within an acceptable range. Furthermore, having diabetes does not directly affect the appropriateness of nitroglycerin administration; however, it is essential to always consider the overall clinical picture, including any complications associated with diabetes, such as potential cardiovascular issues.

SAMPLE

9. What vital signs are critical for assessing a patient's hemodynamic status?

A. Temperature, heart rate, oxygen saturation, and blood pressure

B. Heart rate, blood pressure, respiratory rate, and oxygen saturation

C. Blood glucose, heart rate, respiratory rate, and blood pressure

D. Heart rate, temperature, blood pressure, and level of consciousness

The assessment of a patient's hemodynamic status involves evaluating their cardiovascular and respiratory function, which is essential for determining how well the body is perfusing tissues and delivering oxygen. The correct choice includes heart rate, blood pressure, respiratory rate, and oxygen saturation, as these parameters directly reflect cardiac output, vascular resistance, and pulmonary function. Heart rate indicates the heart's effectiveness in pumping blood, while blood pressure illustrates the force exerted by circulating blood on the walls of blood vessels, providing insight into overall blood volume and vascular health. Respiratory rate offers information about a patient's respiratory effort and oxygen exchange, which is critical in assessing both oxygenation and potential respiratory distress. Oxygen saturation gives a direct measurement of the hemoglobin's oxygen-carrying capacity, essential for determining the adequacy of tissue oxygenation. Together, these vital signs create a comprehensive picture of a patient's hemodynamic status, helping identify conditions such as shock, respiratory failure, or cardiac dysfunction that require immediate intervention. Other choices may include parameters that are important but do not provide as direct an assessment of hemodynamic status specifically.

10. What should be done if a patient has an obstructed airway?

A. Administer medication immediately

B. Perform abdominal thrusts for adults or back blows for infants

C. Call for advanced medical support only

D. Encourage the patient to cough forcefully

In the situation where a patient has an obstructed airway, performing abdominal thrusts for adults or back blows for infants is the appropriate and immediate action. This procedure is specifically designed to dislodge the object causing the obstruction. Abdominal thrusts, also known as the Heimlich maneuver, create an upward force on the diaphragm, leading to increased pressure in the lungs and providing a strong expulsion of air. This can help to expel the obstruction from the airway of an adult or child. For infants, back blows are more suitable, as their anatomy and size require different techniques that are both safer and more effective in dislodging an obstruction. Other options do not address the immediate threat of airway blockage adequately. While medications may be helpful in various emergency scenarios, they do not resolve an obstructed airway. Simply calling for advanced medical support would delay critical intervention when immediate action is necessary to ensure the patient's safety. Encouraging a patient to cough forcefully might be effective in milder cases of obstruction, but it is not a reliable or safe approach for all patients, especially if they are unable to cough effectively. Thus, performing abdominal thrusts or back blows is the established best practice in this situation.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://orangecountyflemsprotocol.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE