PAIN SCIENCE RA 3 PRACTICE (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://painsciencera3.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1
Table of Contents 2
Introduction 3
How to Use This Guide 4
Questions 5
Answers 8
Explanations 10
Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which statement best reflects the clinical implication of the Fear-Avoidance Model in chronic pain?

A. It emphasizes that pain is purely physical.

B. It suggests avoidance is beneficial.

C. It highlights that pain is modulated by social factors.

D. Catastrophizing leads to fear and avoidance, causing disuse and disability; targets include graded exposure, CBT, and activity re-engagement.

2. What is a recommended approach to progressing an exercise program for chronic pain?

A. Start with high intensity and long duration

B. Start with low intensity, short durations; gradually increase weekly; include aerobic and strengthening; monitor response and adapt; set SMART goals

C. Avoid monitoring responses

D. Only perform isometric exercises

3. Which example best represents nociplastic pain?

A. Osteoarthritis

B. Diabetic neuropathy

C. Fibromyalgia

D. Acute muscle strain

4. Which sex is more commonly affected by juvenile idiopathic arthritis (JIA)?

A. Boys

B. Unknown

C. Equal

D. Girls

5. What is the correct order of pain types from most to least common?

A. Back ache > Headache > Stomach ache

B. Headache > Back ache > Stomach ache

C. Stomach ache > Headache > Back

D. Headache > Stomach ache > Back

SAMPLE

6. What are essential components of a pain neuroscience education session?

A. Explain pain biology, emphasize brain as processor, reduce threat, relate to movement and function, address myths, provide actionable strategies

B. Focus only on pharmacology

C. Avoid discussing myths and movement

D. Emphasize tissue damage as sole cause

7. In CRPS, which difference between children and adults is reported?

A. Psychological influence is greater in children

B. Injury history is more common in children

C. Skin is warmer in children

D. Neurological symptoms are more pronounced in children

8. Which statement best describes the frequency of serious medical conditions in older adults presenting with back pain?

A. Common

B. Rare

C. Frequent but overlooked

D. Obvious

9. About what percent of older adults report pain at more than two sites?

A. 60%

B. 50%

C. 65%

D. 70%

10. Which of the following is a treatment component for Osgood-Schlatters?

A. Antibiotics

B. Chemotherapy

C. Exercise

D. Radiation therapy

SAMPLE

Answers

SAMPLE

1. D
2. B
3. C
4. D
5. B
6. A
7. A
8. B
9. D
10. C

SAMPLE

Explanations

SAMPLE

1. Which statement best reflects the clinical implication of the Fear-Avoidance Model in chronic pain?

A. It emphasizes that pain is purely physical.

B. It suggests avoidance is beneficial.

C. It highlights that pain is modulated by social factors.

D. Catastrophizing leads to fear and avoidance, causing disuse and disability; targets include graded exposure, CBT, and activity re-engagement.

Beliefs and emotions about pain can shape how people respond, and those responses can sustain pain and disability. In the Fear-Avoidance Model, thinking catastrophically about pain produces fear that movement will cause harm, which leads to avoiding activity. This avoidance causes disuse and deconditioning, social withdrawal, and greater disability, which can feed back into more pain and fear—a self-perpetuating cycle. Clinically, the implication is to interrupt this cycle by addressing the cognitive and emotional drivers. Targeted approaches aim to reduce catastrophizing and fear, and to get people back into meaningful activity through strategies like graded exposure to feared movements, cognitive-behavioral therapy, and activity re-engagement. These interventions help patients recalibrate beliefs about pain, increase safe activity, and improve function. The other statements miss the main idea: pain is not treated as purely physical, avoidance is not advantageous, and while social factors can influence pain, the most actionable clinical takeaway from the model is the link between fear, avoidance, disuse, and disability and the corresponding treatment targets.

2. What is a recommended approach to progressing an exercise program for chronic pain?

A. Start with high intensity and long duration

B. Start with low intensity, short durations; gradually increase weekly; include aerobic and strengthening; monitor response and adapt; set SMART goals

C. Avoid monitoring responses

D. Only perform isometric exercises

Progression of an exercise program for chronic pain works best when loading is gradual and tailored to the person's responses. Start with low intensity and short sessions, making the activities doable but challenging enough to stimulate adaptation. From there, increase the dose each week in small steps—whether by a bit more time, a little more resistance, or a combination of both—while watching how the body responds. The goal is to build capacity over time without triggering painful flare-ups, so the pace should be guided by how pain, fatigue, and function change after each session. Including both aerobic activities and strengthening exercises is important. Aerobic work improves overall fitness, endurance, and pain modulation, while strengthening supports movement and can reduce vulnerability to future pain problems. Monitoring responses is essential: track pain levels, day-to-day function, mood, and any new or worsening symptoms, and then adapt the plan if responses suggest the current load is too much or not enough. SMART goals—specific, measurable, achievable, relevant, and time-bound—help structure progression, keeping goals clear and motivating and making it easy to decide when to advance. Starting with high intensity or long sessions tends to provoke pain and reduce adherence, and ignoring responses can hide signs that the plan needs adjustment. Limiting to only isometric exercises misses the broader benefits of dynamic loading and cardio conditioning that support real-world function.

SAMPLE

3. Which example best represents nociplastic pain?

A. Osteoarthritis

B. Diabetic neuropathy

C. Fibromyalgia

D. Acute muscle strain

Nociplastic pain comes from altered processing of pain signals in the central nervous system, without clear tissue damage or nerve injury driving the pain. Fibromyalgia fits this pattern because people often have widespread pain and heightened sensitivity to pressure and other stimuli, along with fatigue and sleep issues, but there isn't a single identifiable tissue injury or nerve lesion. In contrast, osteoarthritis pain mainly stems from actual joint tissue wear and inflammation (nociceptive pain from damaged tissue), diabetic neuropathy involves nerve injury causing neuropathic pain, and an acute muscle strain results from recent tissue damage triggering nociceptive pain. So the example that best represents nociplastic pain is fibromyalgia.

4. Which sex is more commonly affected by juvenile idiopathic arthritis (JIA)?

A. Boys

B. Unknown

C. Equal

D. Girls

In juvenile idiopathic arthritis, the pattern across many populations is that girls are more commonly affected than boys. This female predominance is seen across several common JIA subtypes, especially oligoarticular and polyarticular forms, though the degree of difference can vary by subtype and population. The reasons aren't fully understood, but likely involve hormonal influences, genetic factors on the X chromosome, and general immune regulation differences between sexes. Because of this, saying that girls are more commonly affected best fits the observed epidemiology, while saying it's boys, unknown, or equal doesn't align with the data.

5. What is the correct order of pain types from most to least common?

A. Back ache > Headache > Stomach ache

B. Headache > Back ache > Stomach ache

C. Stomach ache > Headache > Back

D. Headache > Stomach ache > Back

Headache is one of the most frequent pain experiences people report, occurring across ages and settings. Back pain is also extremely common and a leading cause of disability, so it ranks high as well but is typically reported after headaches in terms of overall frequency. Abdominal (stomach) pain happens too, but as a standalone chronic pain type it is less frequently reported than headaches or back pain. So the order from most to least common is headache, then back ache, then stomach ache.

SAMPLE

6. What are essential components of a pain neuroscience education session?

A. Explain pain biology, emphasize brain as processor, reduce threat, relate to movement and function, address myths, provide actionable strategies

B. Focus only on pharmacology

C. Avoid discussing myths and movement

D. Emphasize tissue damage as sole cause

Pain neuroscience education centers on teaching that pain is produced by the brain as a response to perceived threat, shaped by beliefs, emotions, and context, not simply the size of tissue damage. The best answer is effective because it covers all essential components: explain how pain biology works; emphasize that the brain acts as the primary processor interpreting threats; work to reduce perceived threat so pain becomes more manageable; connect how pain and movement relate to function in real life, helping people stay active rather than fear movement; address myths that can keep people stuck in fear or misunderstanding; and provide actionable strategies people can use to influence their pain and function day to day. If a session leans only on pharmacology, or avoids discussing myths and movement, or treats tissue damage as the sole cause, it misses the mechanism by which pain is generated and modulated and fails to empower self-management.

7. In CRPS, which difference between children and adults is reported?

A. Psychological influence is greater in children

B. Injury history is more common in children

C. Skin is warmer in children

D. Neurological symptoms are more pronounced in children

In CRPS, how a child experiences and manages the pain is often shaped more by psychological and social context than in adults. Developmentally, children depend on families, schools, and caregivers to interpret symptoms, cope with distress, and decide how much they engage in activity. These factors can strongly influence pain reporting, avoidance, and disability, sometimes more than the underlying physical signs. That's why psychological influences tend to be more evident in pediatric cases: mood, anxiety, stress, family dynamics, and coping strategies play a larger role in the level of disability and recovery. The other possibilities don't show a consistent, recognizable difference between children and adults. Injury history is common in CRPS across ages; skin temperature changes and other neurological signs vary widely and aren't reliably greater in one group.

8. Which statement best describes the frequency of serious medical conditions in older adults presenting with back pain?

A. Common

B. Rare

C. Frequent but overlooked

D. Obvious

The main idea is that serious medical conditions causing back pain in older adults are not common. Most back pain in this group is mechanical or degenerative and benign in nature, so the overall frequency of something serious lurking behind the pain is relatively low. That's why "rare" best fits: while you must remain vigilant for red flags and investigate promptly if red flags are present, the typical presentation does not equate to a common finding of a serious condition. The other descriptions overstate the frequency or imply the seriousness is always obvious, which isn't the case in real clinical practice.

SAMPLE

9. About what percent of older adults report pain at more than two sites?

A. 60%

B. 50%

C. 65%

D. 70%

Pain across multiple sites is a common pattern among older adults, reflecting widespread pain rather than a single localized issue. Population data frequently show that around seven in ten older adults report pain in three or more sites. This high prevalence highlights the importance of a thorough, body-wide pain assessment and integrated management that considers multiple painful areas, functional impact, and related health conditions. Other commonly reported percentages, such as sixty, fifty, or sixty-five percent, underestimate how common mult-site pain is in this group, making seventy percent the best estimate for this question.

10. Which of the following is a treatment component for Osgood-Schlatters?

A. Antibiotics

B. Chemotherapy

C. Exercise

D. Radiation therapy

Managing Osgood-Schlatter disease centers on controlled activity and physical therapy to reduce traction at the tibial tubercle and restore knee function. A key component is a progressive exercise program that includes stretching the quadriceps and hamstrings to improve flexibility, plus strengthening of the surrounding muscles (thigh and hip) and a gradual return to activity as symptoms allow. This approach helps lessen the stress on the tibial tubercle during growth and supports healing. Antibiotics, chemotherapy, and radiation therapy do not address this condition and are not used here; they're for infections or cancer, not apophysitis from overuse.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://painsciencera3.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE