PEDIATRIC ADVANCED LIFE SUPPORT (PALS) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://pediatricadvancedlifesupport.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What tool is used to measure ST segment elevation or depression on an ECG?**

   A. P wave analysis

   B. QRS duration

   C. ST segment measurement

   D. J wave assessment

2. **In which patients can an oropharyngeal airway be used?**

   A. With a strong gag reflex

   B. No gag reflex

   C. Patients under anesthesia

   D. Patients with mild respiratory distress

3. **Which phenomenon is seen during cardiac tamponade, resulting in a larger than normal drop in systolic blood pressure during inspiration?**

   A. Pulsus alternans

   B. Pulsus paradoxus

   C. Pulsus bisferiens

   D. Orthostatic hypotension

4. **What is the primary goal of using chest compressions during CPR?**

   A. To deliver oxygen to the lungs

   B. To maintain blood circulation

   C. To stabilize cardiac rhythm

   D. To clear the airway

5. **When assessing a child's mental status, what does the letter 'P' in AVPU stand for?**

   A. Pain

   B. Pleasant

   C. Persistent

   D. Positive

SAMPLE

6. In PALS, which of the following should be assessed immediately if a child demonstrates decreased responsiveness?

A. Oxygenation

B. Nutrition

C. Hydration status

D. Emotional state

7. What heart rate in children over 2 years old may indicate a serious condition?

A. HR > 140/min

B. HR > 160/min

C. HR > 170/min

D. HR > 180/min

8. What is the typical size of an endotracheal tube for a child compared to their anatomy?

A. Size of the child's little toe

B. Size of the child's 5th finger

C. Size of the child's ear

D. Size of the child's foot

9. What age is classified as infants in pediatric assessments?

A. Under 6 months old

B. Under 1 year old

C. From 1 to 2 years old

D. From newborn to 1 year old

10. Which pharmacologic property of aspirin is important in the treatment of myocardial infarction?

A. Antithrombotic

B. Antipyretic

C. Analgesic

D. Anti-inflammatory

SAMPLE

Answers

SAMPLE

1. D
2. B
3. B
4. B
5. A
6. A
7. B
8. B
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. What tool is used to measure ST segment elevation or depression on an ECG?

A. P wave analysis

B. QRS duration

C. ST segment measurement

D. J wave assessment

The correct answer focuses on the ST segment measurement tool used specifically to assess ST segment elevation or depression on an ECG. This measurement is critical in diagnosing various cardiac conditions, especially in the context of ischemia or myocardial infarction. The ST segment represents the period between the end of ventricular depolarization and the beginning of ventricular repolarization, and changes in its elevation or depression can indicate underlying pathophysiological processes. Utilizing a specific tool for measuring the ST segment allows healthcare professionals to quantify these changes accurately, which is essential for making informed clinical decisions regarding treatment and intervention strategies. Other options, while related to various components of the ECG, do not serve the specific purpose of measuring ST segment changes. P wave analysis looks at atrial depolarization, QRS duration pertains to ventricular depolarization, and J wave assessment refers to a different aspect of the ECG that may indicate particular conditions but does not focus on ST segment evaluation directly. Understanding the role of these various components is crucial in accurately interpreting ECG results and managing pediatric cardiac emergencies effectively.

2. In which patients can an oropharyngeal airway be used?

A. With a strong gag reflex

B. No gag reflex

C. Patients under anesthesia

D. Patients with mild respiratory distress

An oropharyngeal airway is specifically indicated for patients who do not have a gag reflex. This is because the device is designed to keep the airway open by preventing the tongue from blocking the throat and allowing for unobstructed breathing. In patients who have a strong gag reflex, the insertion of an oropharyngeal airway could trigger vomiting or laryngospasm, which can lead to further airway obstruction or complications. In clinical practice, understanding the patient's level of consciousness and airway reflexes is essential before determining the appropriate airway management strategy. Patients who are under anesthesia often have their gag reflex diminished or absent, making an oropharyngeal airway suitable in the controlled setting of anesthesia; however, the primary focus for securing the airway is typically through endotracheal intubation in such situations. Mild respiratory distress may not warrant the use of an oropharyngeal airway unless accompanied by altered consciousness, as maintaining the patient's natural airway efforts is usually preferable when they still have an intact gag reflex. Therefore, the safest group for whom an oropharyngeal airway is intended is those without a gag reflex.

SAMPLE

3. Which phenomenon is seen during cardiac tamponade, resulting in a larger than normal drop in systolic blood pressure during inspiration?

A. Pulsus alternans

B. Pulsus paradoxus

C. Pulsus bisferiens

D. Orthostatic hypotension

The phenomenon that occurs during cardiac tamponade, characterized by a larger than normal drop in systolic blood pressure during inspiration, is known as pulsus paradoxus. In this condition, the drop in blood pressure exceeds the usual variation that occurs with respiration and is indicative of impaired filling of the heart chambers due to external pressure. During inspiration, negative intrathoracic pressure facilitates increased venous return to the heart. However, in cardiac tamponade, the fluid accumulation in the pericardial space restricts the heart's ability to expand, particularly affecting the right ventricle during diastole. This results in decreased stroke volume and consequentially a marked decrease in systolic blood pressure, which can be more than 10 mmHg. As for the other options, pulsus alternans refers to alternating strong and weak heartbeats, pulsus bisferiens is characterized by a double pulse wave, and orthostatic hypotension is the drop in blood pressure upon standing. None of these specifically relate to the pronounced drop in systolic blood pressure during inspiration linked to cardiac tamponade.

4. What is the primary goal of using chest compressions during CPR?

A. To deliver oxygen to the lungs

B. To maintain blood circulation

C. To stabilize cardiac rhythm

D. To clear the airway

The primary goal of using chest compressions during CPR is to maintain blood circulation. When a person goes into cardiac arrest, their heart stops pumping blood effectively, which deprives vital organs, including the brain, of oxygen. Chest compressions generate artificial circulation by compressing the chest and forcing blood to flow through the heart and into the rest of the body. This is crucial in preserving the function of vital organs until advanced medical help arrives or until a normal heart rhythm is restored. While other aspects of CPR, such as delivering oxygen to the lungs and clearing the airway, are important components, they serve to complement the primary objective of ensuring that blood continues to circulate effectively. Stabilizing cardiac rhythm is also a goal but is not within the immediate scope of what chest compressions directly achieve; rather, it is hoped that compressions will support the heart's natural rhythm until defibrillation or advanced interventions can be applied.

SAMPLE

5. When assessing a child's mental status, what does the letter 'P' in AVPU stand for?

A. Pain

B. Pleasant

C. Persistent

D. Positive

In the AVPU scale used to assess a child's mental status, the letter 'P' stands for 'Pain.' This particular assessment tool is utilized to evaluate a patient's responsiveness and level of consciousness in emergency situations. The AVPU acronym stands for Alert, Voice, Pain, and Unresponsive. When a child is assessed as responding to 'Pain,' it indicates that the child does not respond to verbal stimuli but does react to physical stimuli, such as a gentle shake or painful stimulus. This response is a critical indicator of the child's neurological function and helps in determining the severity of their condition. Recognizing the levels of responsiveness, from being fully alert to unresponsive, allows healthcare providers to make informed decisions regarding the necessary interventions and urgency of care needed.

6. In PALS, which of the following should be assessed immediately if a child demonstrates decreased responsiveness?

A. Oxygenation

B. Nutrition

C. Hydration status

D. Emotional state

In the context of Pediatric Advanced Life Support (PALS), immediate assessment of oxygenation is crucial when a child displays decreased responsiveness. Oxygenation refers to the adequacy of oxygen supply to the tissues, which is vital for cellular function and overall organ performance. In children, decreased responsiveness can often be an indicator of inadequate oxygenation due to various factors such as respiratory distress, airway obstruction, or hypoventilation. Promptly assessing oxygen levels and the child's respiratory status can help identify whether supplemental oxygen or advanced airway management is required. Rapid intervention to restore adequate oxygenation can prevent further deterioration of the child's condition and support vital organ systems. While nutrition, hydration status, and emotional state are important considerations for a child's overall health, they do not require immediate assessment in an emergency situation where decreased responsiveness is evident. Addressing oxygenation as a priority maximizes the chances of stabilizing the child and potentially reversing the underlying causes of their decreased responsiveness.

SAMPLE

7. What heart rate in children over 2 years old may indicate a serious condition?

A. HR > 140/min

B. HR > 160/min

C. HR > 170/min

D. HR > 180/min

In children over 2 years old, a heart rate greater than 160 beats per minute can indicate a serious medical condition. This threshold is significant because it is higher than the upper limit of normal resting heart rates for this age group, which typically ranges from 70 to 130 beats per minute. When a child's heart rate exceeds this range, it can suggest potential underlying issues such as fever, dehydration, infection, or even cardiac arrhythmias. Monitoring heart rate is essential in assessing a child's overall health status, especially in emergency settings. A sustained heart rate above 160 indicates that the child might be experiencing distress, whether due to physical exertion or illness, and warrants further evaluation and possibly immediate medical intervention.

8. What is the typical size of an endotracheal tube for a child compared to their anatomy?

A. Size of the child's little toe

B. Size of the child's 5th finger

C. Size of the child's ear

D. Size of the child's foot

The typical size of an endotracheal tube for a child is often correlated with the size of the child's fifth finger. This correlation takes into account the anatomy of a child's airway, allowing for a more accurate estimation of the appropriate tube size based on their age and physical development. The fifth finger, or little finger, is generally a good indicator of a child's airway diameter, helping to select a tube that is both appropriately sized for ventilation and minimizes trauma to the airway. By using the size of the child's little finger as a guide, medical professionals can better ensure that the endotracheal tube will provide effective airflow without being overly large, which could lead to complications such as airway obstruction or injury. This method emphasizes the importance of anatomical relevance when selecting equipment for pediatric care, ensuring that interventions are both safe and effective.

9. What age is classified as infants in pediatric assessments?

A. Under 6 months old

B. Under 1 year old

C. From 1 to 2 years old

D. From newborn to 1 year old

Infants in pediatric assessments are classified as being under 1 year old. This age range encompasses the critical developmental milestones that occur during infancy, including physical, cognitive, and social-emotional growth. Recognizing this age range is important in PALS, as it influences the clinical assessments, treatment protocols, and interventions that are appropriate for this young population. This classification helps healthcare providers tailor their approach to meet the unique needs of infants, as their physiology and responses to illness and treatment can differ significantly from older children and adults. Understanding these developmental stages ensures that pediatric care is both effective and age-appropriate.

SAMPLE

10. Which pharmacologic property of aspirin is important in the treatment of myocardial infarction?

A. Antithrombotic

B. Antipyretic

C. Analgesic

D. Anti-inflammatory

The pharmacologic property of aspirin that is crucial in the treatment of myocardial infarction is its antithrombotic effect. Aspirin works by inhibiting platelet aggregation, which reduces the ability of platelets to clump together and form blood clots. In the context of a myocardial infarction, where a clot can obstruct blood flow to the heart muscle, the antithrombotic action of aspirin plays a vital role in preventing further clot formation or extension of an existing clot. This can help restore blood flow and limit damage to the heart tissue. While aspirin does have antipyretic properties that reduce fever, analgesic effects for pain relief, and anti-inflammatory effects, these are not the primary reasons it's used in an acute myocardial infarction scenario. The focus during a myocardial infarction is to prevent clot formation and ensure blood flow, making the antithrombotic property the critical factor in its therapeutic role.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://pediatricadvancedlifesupport.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE