PEDIATRIC EMERGENCIES PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://pediatricemergencies.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Currant-jelly stools are characteristic of which condition, and what is the recommended initial evaluation step?

A. Appendicitis; ultrasound

B. Meckel diverticulum; barium study

C. Gastroenteritis; stool culture

D. Intussusception; ultrasound confirmation

2. Which finding is not typically associated with cardiogenic shock in a child?

A. Warm, well-perfused extremities

B. Mottled skin

C. Low urine output

D. Altered mental status

3. Bronchiolitis is best described as which of the following?

A. A viral infection of the lower airway that commonly affects infants and children younger than 2 years of age

B. A bacterial infection of the upper airway

C. A fungal infection of the lungs

D. An allergic inflammatory reaction of the airways

4. Caustic ingestion (drain cleaner) in a child: initial management?

A. Induce emesis to dilute the chemical.

B. Neutralize with an alkali immediately.

C. Call poison control and observe at home.

D. Do not induce emesis; rinse mouth; secure airway if needed; obtain pediatric toxicology input.

5. Early distributive shock in children is characterized by:**

A. Cold, clammy skin

B. Warm, flushed skin

C. Low heart rate

D. Bradycardia

SAMPLE

6. A 7-month-old infant presents with listlessness, pallor, and increased work of breathing and has a rapid heart rate that is unresponsive to oxygen; which cardiac rhythm is most likely on monitor?

A. Narrow QRS complex rhythm with absent P waves and heart rate greater than 220 beats/min.

B. Wide QRS complex rhythm with reentrant tachycardia

C. Sinus tachycardia at a moderate rate

D. Atrial flutter with regular rhythm

7. In contrast to adults, children:

A. Have proportionately larger heads.

B. Have proportionately smaller heads.

C. Have larger legs relative to arms.

D. Have less head-to-body proportion.

8. Which component is NOT used to distinguish sinus tachycardia from reentry supraventricular tachycardia?

A. P waves

B. QRS complex width

C. PR interval

D. Heart rate

9. If a 2-year-old child with a foreign body airway obstruction becomes unresponsive, you should:

A. Position the child prone and perform back blows

B. Shout for help and wait for EMS

C. Remove the foreign body with a finger sweep

D. Position supine and perform chest compressions

10. Dilated cardiomyopathy is best described as which of the following?

A. Thickened stiff walls

B. Normal pumping with dilation

C. Isolated atrial dilation

D. Weakened and enlarged, making it a less efficient pump

SAMPLE

Answers

SAMPLE

1. D
2. A
3. A
4. D
5. B
6. A
7. A
8. B
9. D
10. D

SAMPLE

Explanations

SAMPLE

1. Currant-jelly stools are characteristic of which condition, and what is the recommended initial evaluation step?

A. Appendicitis; ultrasound

B. Meckel diverticulum; barium study

C. Gastroenteritis; stool culture

D. Intussusception; ultrasound confirmation

Currant-jelly stools point to intussusception, where a segment of bowel slides into another, causing mucosal venous congestion and bleeding that creates the bloody, mucus-filled stool. The best first step to evaluate this suspected condition is an abdominal ultrasound. It's noninvasive and highly sensitive, and it typically shows the characteristic target (doughnut) sign in cross-section, confirming the diagnosis quickly. If the ultrasound confirms intussusception and the child is stable, nonoperative reduction with an air contrast enema is usually attempted.

2. Which finding is not typically associated with cardiogenic shock in a child?

A. Warm, well-perfused extremities

B. Mottled skin

C. Low urine output

D. Altered mental status

In cardiogenic shock, the heart fails to pump effectively, so cardiac output drops and tissues become underperfused. To preserve blood flow to vital organs, the body constricts peripheral vessels, leading to cool, sometimes mottled skin, decreased urine output from reduced renal perfusion, and possible altered mental status from brain hypoperfusion. Warm, well-perfused extremities would not fit this pattern; they suggest better peripheral perfusion or a shock type with vasodilation, such as distributive shock, rather than cardiogenic shock. So the finding of warm, well-perfused extremities is not typical of pediatric cardiogenic shock, whereas mottled skin, low urine output, and altered mental status align with this condition.

3. Bronchiolitis is best described as which of the following?

A. A viral infection of the lower airway that commonly affects infants and children younger than 2 years of age

B. A bacterial infection of the upper airway

C. A fungal infection of the lungs

D. An allergic inflammatory reaction of the airways

Bronchiolitis is a viral infection of the lower airways that commonly affects infants and children younger than two years. It is most often caused by RSV and involves inflammation of the bronchioles, leading to airway obstruction with wheeze, tachypnea, and increased work of breathing, sometimes with fever and nasal symptoms. This description fits because bronchiolitis is a small-airway (lower respiratory tract) viral illness in very young children, rather than a bacterial infection of the upper airway, a fungal lung infection, or an allergic inflammatory process. Management is typically supportive, focusing on hydration, nasal suctioning, and oxygen if needed, with antibiotics not routinely indicated unless a bacterial coinfection is suspected.

SAMPLE

4. Caustic ingestion (drain cleaner) in a child: initial management?

A. Induce emesis to dilute the chemical.

B. Neutralize with an alkali immediately.

C. Call poison control and observe at home.

D. Do not induce emesis; rinse mouth; secure airway if needed; obtain pediatric toxicology input.

Caustic ingestion can cause ongoing mucosal injury and potential airway compromise, so the priority is to protect the airway and limit further damage while obtaining expert guidance. Do not induce vomiting or use neutralizing agents because throwing up or reacting the chemical with acids or bases can worsen tissue injury and increase the risk of perforation or aspiration. Instead, rinse the mouth to remove residual chemical and assess the patient for signs of airway difficulty; secure the airway if there are symptoms such as drooling, stridor, hoarseness, or respiratory distress. Seek pediatric toxicology input promptly to determine the appropriate next steps, including need for endoscopy or admission. Home observation is unsafe because injuries can evolve and require professional evaluation and monitoring.

5. Early distributive shock in children is characterized by:**

A. Cold, clammy skin

B. Warm, flushed skin

C. Low heart rate

D. Bradycardia

Early distributive shock in children shows vasodilation with maintained or increased cardiac output, so the skin feels warm and appears flushed. This warm, flushed appearance reflects increased cutaneous blood flow despite overall circulatory instability. The body often responds with tachycardia to support perfusion, not with a slow heart rate. In contrast, cold, clammy skin points to late shock or a different pattern (such as hypovolemic or cardiogenic states), and a low heart rate or bradycardia is not typical of the early distributive phase. So warm, flushed skin best identifies the early distributive pattern.

6. A 7-month-old infant presents with listlessness, pallor, and increased work of breathing and has a rapid heart rate that is unresponsive to oxygen; which cardiac rhythm is most likely on monitor?

A. Narrow QRS complex rhythm with absent P waves and heart rate greater than 220 beats/min.

B. Wide QRS complex rhythm with reentrant tachycardia

C. Sinus tachycardia at a moderate rate

D. Atrial flutter with regular rhythm

In infants who are listless, pale, and working hard to breathe, a very fast, regular, narrow-complex tachycardia is most concerning and classically points to supraventricular tachycardia caused by a reentrant circuit. The rapid rate often exceeds 220 beats per minute in this age group, and because the atrial activity is occurring so quickly, distinct P waves are not seen on the monitor. The ventricles are still being activated through the normal conduction system, which is why the QRS complexes remain narrow. This pattern explains the hemodynamic instability: the heart isn't able to fill adequately between beats, reducing cardiac output and leading to poor perfusion despite supplemental oxygen. If you compared it to other rhythms, a wide QRS tachycardia would suggest ventricular involvement or aberrant conduction, which is less typical in this clinical scenario. Sinus tachycardia would show visible P waves and a more moderate rate, and atrial flutter would produce a sawtooth pattern with a different rhythm. The described narrow, rapid, P-wave–invisible tachycardia best fits infant SVT, the rhythm most likely responsible for these symptoms.

SAMPLE

7. In contrast to adults, children:

A. Have proportionately larger heads.

B. Have proportionately smaller heads.

C. Have larger legs relative to arms.

D. Have less head-to-body proportion.

Children have proportionally larger heads because brain growth drives rapid expansion of the cranium in early life, and the skull bones are not yet fully fused. This makes the head appear oversized relative to the rest of the body, a trait that gives infants and young children a top-heavy look. Over time, the trunk and limbs grow faster, and the body proportions shift toward the smaller head-to-body ratio seen in adults. That's why the statement that children have proportionately larger heads is the best description. The other ideas don't fit the typical growth pattern: the head isn't proportionally smaller, limb proportions aren't defined by a larger leg relative to the arm in this context, and saying there's less head-to-body proportion contradicts the observed early-childhood pattern.

8. Which component is NOT used to distinguish sinus tachycardia from reentry supraventricular tachycardia?

A. P waves

B. QRS complex width

C. PR interval

D. Heart rate

Distinguishing sinus tachycardia from reentry SVT relies on atrial activity and rate more than how wide the ventricular complex is. The QRS width is not a reliable discriminator because both conditions can have a normal, narrow QRS when there's no aberrant conduction. In sinus tachycardia you typically see visible P waves preceding each QRS with a normal PR interval and a rate that rises in response to physiologic stress. In reentry SVT, P waves may be hidden or appear abnormally related to the QRS, and the rhythm is usually very rapid and regular, with a PR interval that can be normal or difficult to measure. So the QRS width doesn't help differentiate these two tachycardias, while the presence and timing of P waves, PR interval, and overall rate do.

9. If a 2-year-old child with a foreign body airway obstruction becomes unresponsive, you should:

A. Position the child prone and perform back blows

B. Shout for help and wait for EMS

C. Remove the foreign body with a finger sweep

D. Position supine and perform chest compressions

When a child who is choking becomes unresponsive, the priority is immediate CPR to maintain circulation and oxygen delivery. Place the child on their back and start high-quality chest compressions right away. The chest compressions help perfuse the brain and heart and may help move the obstruction as air is pushed in during rescue breaths. Once unresponsive, the emphasis shifts from relief maneuvers to restoring circulation; back blows and chest thrusts are for a conscious choking child, not for an unresponsive one. A finger sweep should be avoided unless you can clearly see the object and can safely remove it without pushing it deeper. After starting compressions, provide rescue breaths as appropriate (for a single rescuer, use a 30:2 ratio). Continue CPR and call for help if you haven't already.

SAMPLE

10. Dilated cardiomyopathy is best described as which of the following?

A. Thickened stiff walls

B. Normal pumping with dilation

C. Isolated atrial dilation

D. Weakened and enlarged, making it a less efficient pump

Dilated cardiomyopathy describes a heart with enlarged chambers, especially the ventricles, and a weakened muscular contraction, so the heart can't pump as effectively. That combination—enlarged, underperforming pump—is precisely what "weakened and enlarged, making it a less efficient pump" conveys. Other options miss the key point: thickened, stiff walls describe hypertrophic cardiomyopathy; normal pumping with dilation contradicts the hallmark systolic dysfunction; isolated atrial dilation isn't the pattern seen in dilated cardiomyopathy.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://pediatricemergencies.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE