PERIOPERATIVE NURSING CARE PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://perioperativenursingcare.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the goal of reconstructive surgery?

A. Remove diseased tissue

B. Repair congenital defects

C. Restore appearance and function of damaged organs or tissues

D. Establish a diagnosis

2. What does evisceration refer to in a surgical context?

A. Protrusion of the abdominal viscera through the wound

B. Incomplete wound healing after surgery

C. A type of infection following a surgical procedure

D. Excessive bleeding from a surgical site

3. How long is an informed consent valid once signed by the patient?

A. 15 days

B. 30 days

C. 60 days

D. 90 days

4. What percentage of pneumonia cases are approximately caused by gram-bacilli?

A. About one-third

B. About one-half

C. Less than one-quarter

D. About three-quarters

5. Which of the following patients is at risk for urinary retention after surgery?

A. Patients who have had intense physical training

B. Patients who have undergone proctological surgery

C. Women with multiple pregnancies

D. Patients with no prior surgeries

SAMPLE

6. What is a common postoperative condition observed in patients with preexisting pulmonary dysfunction?

A. Hypoxia

B. Hypoventilation

C. Atelectasis

D. Hyperventilation

7. What is one way to educate patients about their surgical risks?

A. Providing clear verbal instructions and written materials

B. Using highly technical terms

C. Limiting discussion to the surgeon's expertise only

D. Assuming patients are aware of all risks

8. Which condition results from a critical reduction in functional residual capacity post-surgery?

A. Bronchospasm

B. Atelectasis

C. Pneumonia

D. Hypoxia

9. What is a potential complication of an over-distended bladder?

A. Acute kidney failure

B. Confusion and muscle relaxation issues

C. Severe abdominal pain

D. Increased urinary output

10. Why is preoperative fasting important?

A. To save time during surgery

B. To prevent nausea and vomiting post-surgery

C. To reduce the risk of aspiration during anesthesia

D. To avoid discomfort during recovery

SAMPLE

Answers

SAMPLE

1. C
2. A
3. B
4. B
5. C
6. B
7. A
8. B
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. What is the goal of reconstructive surgery?

A. Remove diseased tissue

B. Repair congenital defects

C. Restore appearance and function of damaged organs or tissues

D. Establish a diagnosis

The goal of reconstructive surgery primarily focuses on restoring both the appearance and the function of damaged organs or tissues. This type of surgery is often performed after trauma, infection, or surgery that has resulted in a loss of form or function. It encompasses a broad range of procedures that can address issues such as facial reconstruction after an accident, repairing damaged limbs, or correcting structural abnormalities from birth, thus aiming to improve both the aesthetics and functionality in patients' lives. Other goals of surgery, such as removing diseased tissue, repairing congenital defects, or establishing a diagnosis, do not encapsulate the comprehensive intent of reconstructive surgery. While some reconstructive procedures can involve addressing congenital defects, the overarching aim of these surgeries is to enable patients to regain functionality and improve their quality of life post-injury or surgery.

2. What does evisceration refer to in a surgical context?

A. Protrusion of the abdominal viscera through the wound

B. Incomplete wound healing after surgery

C. A type of infection following a surgical procedure

D. Excessive bleeding from a surgical site

Evisceration in a surgical context specifically refers to the protrusion of the abdominal organs or viscera through an opened or dehisced wound. This can occur as a serious complication following surgery, particularly abdominal surgeries, and may indicate that the surgical site has not healed adequately. Recognizing evisceration is crucial for timely intervention, as it can lead to increased risk of infection, further complications, and the need for surgical repair. The other options describe different conditions or complications: incomplete wound healing pertains to surgical wounds that do not close properly but do not necessarily involve the protrusion of organs; an infection related to a surgical procedure refers to the presence of pathogenic organisms causing inflammation; and excessive bleeding from a surgical site illustrates a hemorrhagic complication rather than an issue with wound integrity. Hence, the focus on the protrusion of organs makes the definition of evisceration clear and distinct in its significance and clinical implications.

3. How long is an informed consent valid once signed by the patient?

A. 15 days

B. 30 days

C. 60 days

D. 90 days

The validity of informed consent once signed by the patient is commonly recognized as being around 30 days. This timeframe acknowledges the necessity for patients to make informed decisions about their healthcare options while ensuring that the information they receive, along with their consent, remains relevant to their treatment and any potential surgical procedures. After 30 days, the nature of a patient's health condition or the specifics of the planned procedure may change, requiring renewed discussion and possibly a new consent process. This is particularly important in ensuring that the patient's understanding and consent reflect their current state and any new risks or benefits that may have arisen since the initial consent was obtained. While specific policies may vary by institution or local regulations, the 30-day guideline helps uphold ethical standards and patient autonomy in the perioperative setting.

SAMPLE

4. What percentage of pneumonia cases are approximately caused by gram-bacilli?

A. About one-third

B. About one-half

C. Less than one-quarter

D. About three-quarters

The correct answer reflects that approximately one-half of pneumonia cases are caused by gram-negative bacilli. This understanding is crucial in perioperative nursing care because it highlights the significance of recognizing the types of bacteria that can lead to pneumonia, particularly in patients undergoing surgery. Gram-negative bacilli, such as E. coli and Klebsiella pneumoniae, are known to be common pathogens associated with pneumonia, especially in hospitalized or immunocompromised patients. Awareness of this helps healthcare providers tailor antibiotic prophylaxis and treatment strategies effectively, ensuring better outcomes for surgical patients who may be at increased risk of developing pneumonia postoperatively. While the other choices present varying estimates, they do not accurately reflect the significant prevalence of gram-negative bacilli as causative agents in pneumonia cases, making it critical for nursing practice to focus on the correct understanding of these statistics for effective patient care.

5. Which of the following patients is at risk for urinary retention after surgery?

A. Patients who have had intense physical training

B. Patients who have undergone proctological surgery

C. Women with multiple pregnancies

D. Patients with no prior surgeries

Women with multiple pregnancies are at an increased risk for urinary retention after surgery primarily due to pelvic floor changes that occur during pregnancy and childbirth. These changes can lead to weakened pelvic muscles and potential nerve damage, both of which may affect the urinary system and obstruct normal bladder function. In the perioperative setting, anesthesia, particularly spinal or epidural anesthesia, can also contribute to urinary retention by inhibiting the bladder's ability to contract and empty effectively. Given that women who have gone through multiple pregnancies may already have compromised pelvic support, the addition of surgical intervention further elevates their risk for urinary retention. While other patient scenarios may involve specific risks related to surgical procedures, they do not have the same direct connection to urinary function as the physiological changes faced by women with multiple pregnancies.

SAMPLE

6. What is a common postoperative condition observed in patients with preexisting pulmonary dysfunction?

A. Hypoxia

B. Hypoventilation

C. Atelectasis

D. Hyperventilation

In patients with preexisting pulmonary dysfunction, hypoventilation is a common postoperative condition due to several factors associated with anesthesia and the surgical procedure itself. These patients may have impaired respiratory function before surgery, meaning their ability to ventilate effectively can be further compromised after undergoing anesthesia. During surgery, especially when general anesthesia is used, the respiratory drive can be decreased. This decreased drive, combined with the effects of medications and the possibility of postoperative pain, can lead to shallow breathing. The result is hypoventilation, where the patient does not breathe deeply enough or frequently enough to maintain adequate alveolar ventilation, leading to an increase in carbon dioxide levels and a decrease in oxygen levels. Understanding the implications of hypoventilation is critical in the postoperative care setting, especially for patients with existing pulmonary issues. This condition can exacerbate any preexisting respiratory problems, necessitating thorough monitoring and interventions to ensure adequate respiratory function and gas exchange in the immediate post-surgical period.

7. What is one way to educate patients about their surgical risks?

A. Providing clear verbal instructions and written materials

B. Using highly technical terms

C. Limiting discussion to the surgeon's expertise only

D. Assuming patients are aware of all risks

Providing clear verbal instructions and written materials is essential in educating patients about their surgical risks because it promotes understanding and retention of critical information. Clear communication helps to demystify complex medical terminology, ensuring that patients grasp the risks involved in their procedures. When patients receive straightforward verbal explanations alongside written materials, they have the opportunity to review the information at their own pace, which can enhance comprehension. This approach also allows patients to ask questions and seek clarification on any aspects they find confusing, leading to better-informed consent processes and overall satisfaction with their care. Utilizing highly technical terms can alienate patients who may not have a medical background, causing confusion rather than clarity. Limiting the discussion to the surgeon's expertise overlooks the importance of holistic patient education, which should include input from the entire healthcare team. Assuming patients are aware of all risks can lead to incomplete assessments of their understanding and leave them unprepared for potential complications, ultimately undermining the informed consent process.

SAMPLE

8. Which condition results from a critical reduction in functional residual capacity post-surgery?

A. Bronchospasm

B. Atelectasis

C. Pneumonia

D. Hypoxia

Atelectasis occurs when there is a critical reduction in functional residual capacity, particularly following surgery. This condition involves the collapse of alveoli in the lungs, which can prevent effective gas exchange and lead to decreased oxygenation of the blood. Postoperatively, factors such as shallow breathing, pain, and immobility may contribute to the reduced lung volume, increasing the likelihood of atelectasis. By understanding how functional residual capacity is compromised after surgical procedures, healthcare providers can implement preventative measures and interventions, such as encouraging deep breathing exercises, early mobilization, and the use of incentive spirometry. These strategies help to maintain lung function and prevent complications like atelectasis, which can significantly affect patient recovery. Other conditions listed, such as bronchospasm, pneumonia, and hypoxia, while potentially related to postoperative complications, do not directly stem from the specific reduction in functional residual capacity as seen in atelectasis. Each has its own etiology but is not primarily associated with the immediate impact of reduced lung volume following surgery.

9. What is a potential complication of an over-distended bladder?

A. Acute kidney failure

B. Confusion and muscle relaxation issues

C. Severe abdominal pain

D. Increased urinary output

Over-distension of the bladder can lead to a range of complications, and confusion along with muscle relaxation issues is one of them. When the bladder becomes overly full, it can stretch beyond its normal capacity, which might disrupt the normal signaling between the bladder and the nervous system. This disruption can result in confusion, particularly in vulnerable populations such as the elderly, as well as a decrease in the ability to control bladder contractions properly. In terms of the physiological response, an over-distended bladder can inhibit the bladder wall's ability to contract effectively, causing relaxation of the muscles involved in voiding. This can lead to symptoms such as urgency and increased discomfort, as well as cognitive changes like confusion, especially if the individual is unable to void. The potential for confusion arises from the body's response to discomfort and the impact of the condition on mental clarity. Further, if the bladder remains distended for a prolonged period, it can lead to more severe complications, including urinary retention and potential kidney issues, but the immediate manifestation in a patient may be confusion and lack of coordination due to muscle issues. Hence, recognizing the signs and symptoms associated with bladder over-distension is critical for timely intervention in perioperative nursing practice.

SAMPLE

10. Why is preoperative fasting important?

A. To save time during surgery

B. To prevent nausea and vomiting post-surgery

C. To reduce the risk of aspiration during anesthesia

D. To avoid discomfort during recovery

Preoperative fasting is crucial primarily to reduce the risk of aspiration during anesthesia. When a patient undergoes general anesthesia, their protective reflexes, including the gag reflex, can be impaired. If the stomach contains food or liquid, there is a heightened risk that these substances could be regurgitated and then aspirated into the lungs, leading to serious complications such as aspiration pneumonia or airway obstruction. Fasting helps ensure that the stomach is empty, significantly minimizing the possibility of aspiration. This practice aligns with the guidelines from anesthesiology and surgical safety protocols, which advocate for appropriate fasting times based on the type of food ingested. Clear liquids may have a shorter fasting period than solid foods, considering their quicker digestion. Other reasons for preoperative fasting, while relevant, are secondary to the primary concern of aspiration risk. For example, while preventing nausea and vomiting post-surgery and avoiding discomfort during recovery are important, they do not address the immediate and life-threatening risk of aspiration as critically as the need to maintain an empty stomach before anesthesia. The focus on safety and minimizing complications underscores the importance of adherence to preoperative fasting guidelines.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://perioperativenursingcare.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE