PHARMACOLOGY III – CNS MODULE PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://pharmacology3cns.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What effect does propofol have on the central nervous system?

A. Stimulates neurotransmitter release
B. Increases excitability
C. Induces sedation
D. Enhances cognitive function

2. What is a notable adverse effect associated with propofol administration?

A. Increased blood pressure
B. Microbial growth
C. Prolonged sedation
D. Cardiac arrest

3. Which disease is characterized by a deficiency in dopamine?

A. Alzheimer's disease
B. Parkinson's disease
C. Multiple sclerosis
D. Huntington's disease

4. What are contraindications of anticholinergics?

A. Severe dehydration, hypertension
B. Dementia, glaucoma
C. Sleep apnea, asthma
D. Kidney disease, liver disease

5. Which symptom is not classified under extrapyramidal symptoms?

A. Tardive dyskinesia
B. Hallucinations
C. Asynchrony
D. Acute dystonia

6. In terms of pharmacokinetics, what is a key feature of dopaminergic drugs?

A. They have high lipid solubility
B. They are poorly absorbed
C. They bind tightly to plasma proteins
D. They are not metabolized in the liver

SAMPLE

7. What component of carbidopa-levodopa inhibits the enzyme dopa decarboxylase?

A. Levodopa

B. Carbidopa

C. Ropinirole

D. Bromocriptine

8. What is the antidote for benzodiazepine overdose?

A. Activated charcoal

B. Flumazenil

C. Diphenhydramine

D. Rivastigmine

9. Which COMT inhibitor is often used with levodopa in advanced Parkinson's Disease and may cause liver damage?

A. Entacapone

B. Tolcapone

C. Selegiline

D. Amantadine

10. Which of the following is NOT true about MAO inhibitors?

A. They can cause hypertensive crisis with tyramine.

B. They are used to increase dopamine levels.

C. They selectively inhibit MAO-A only.

D. They can interact with a variety of medications.

SAMPLE

Answers

SAMPLE

1. C
2. B
3. B
4. B
5. B
6. A
7. B
8. B
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. What effect does propofol have on the central nervous system?

A. Stimulates neurotransmitter release

B. Increases excitability

C. Induces sedation

D. Enhances cognitive function

Propofol is primarily known for its sedative properties and is commonly used in clinical settings for sedation and anesthesia. Its mechanism of action involves enhancing the inhibitory effect of the neurotransmitter gamma-aminobutyric acid (GABA) at the GABA-A receptor. This results in increased inhibitory neurotransmission, leading to a calming effect on the central nervous system, which is characterized by sedation. When administered, propofol facilitates the transition to unconsciousness and reduces the perception of pain, making it an effective agent for procedures requiring sedation. Its rapid onset and short duration of action are particularly valued in both operating and procedural settings. In contrast, the other options describe effects that are not consistent with propofol's pharmacological profile. It does not stimulate neurotransmitter release or increase excitability, nor does it enhance cognitive function; rather, these options describe activities that align with excitatory neurotransmission or improvement in neural activity, which is contrary to the sedative effects produced by propofol.

2. What is a notable adverse effect associated with propofol administration?

A. Increased blood pressure

B. Microbial growth

C. Prolonged sedation

D. Cardiac arrest

The notable adverse effect associated with propofol administration is microbial growth. Propofol is formulated in a lipid emulsion that can provide an environment conducive to microbial proliferation if not handled or stored properly. This concern is particularly relevant because propofol is often administered in a setting where sterility is critical, such as in surgical or intensive care environments. Inappropriate handling, such as using outdated vials or leaving opened vials at room temperature, can lead to contamination, which may cause serious infections in patients receiving the drug. It's essential for healthcare providers to adhere strictly to guidelines for the storage and administration of propofol to mitigate this risk. Prolonged sedation and cardiac arrest can occur with propofol, but these are less specifically tied to its formulation and more a consequence of the pharmacodynamic effects of an anesthetic agent. Increased blood pressure is not typically a characteristic effect of propofol; in fact, it often causes hypotension. Therefore, understanding the risk of microbial growth highlights a specific safety concern that should always be considered during its administration.

SAMPLE

3. Which disease is characterized by a deficiency in dopamine?

A. Alzheimer's disease

B. Parkinson's disease

C. Multiple sclerosis

D. Huntington's disease

Parkinson's disease is characterized by a deficiency in dopamine, which is a crucial neurotransmitter involved in the regulation of movement and coordination. In Parkinson's disease, the dopaminergic neurons in the substantia nigra of the brain degenerate and die, leading to a significant decrease in dopamine levels. This deficiency results in the hallmark symptoms of Parkinson's disease, including tremors, stiffness, bradykinesia (slowness of movement), and postural instability. The role of dopamine in the central nervous system is vital for facilitating smooth and controlled movements, as it helps transmit signals within the basal ganglia, which are responsible for coordinating voluntary movements. A deficiency not only disrupts normal motor function but can also impact mood and cognitive functions, further complicating the clinical picture. Understanding the relationship between dopamine deficiency and Parkinson's disease is essential for recognizing the underlying mechanisms of the disorder, which ultimately informs treatment strategies that may involve dopamine replacement therapies or medications aimed at enhancing dopaminergic activity.

4. What are contraindications of anticholinergics?

A. Severe dehydration, hypertension

B. Dementia, glaucoma

C. Sleep apnea, asthma

D. Kidney disease, liver disease

Anticholinergics are a class of medications that block the action of acetylcholine, a neurotransmitter involved in various bodily functions. The correct answer highlights the contraindications of using anticholinergics, particularly in patients with dementia and glaucoma. In patients with dementia, anticholinergics can exacerbate cognitive impairment and confusion. Acetylcholine plays a critical role in memory and learning processes; thus, blocking its action can worsen the condition of individuals who already have compromised cognitive functions, leading to a decline in their overall mental status. Regarding glaucoma, anticholinergics can potentially increase intraocular pressure, which can lead to worsening of the condition. This is particularly relevant in closed-angle glaucoma, where the angle between the iris and cornea is already narrow. By inhibiting the outflow of aqueous humor due to pupil dilation (mydriasis), anticholinergics can precipitate a glaucoma attack, making these medications contraindicated for patients with this eye condition. Other choices provided — such as severe dehydration with hypertension, sleep apnea with asthma, and kidney or liver disease — are considerations for specific side effects or interactions but are not universally recognized as absolute contraindications for the use of anticholinergics like dementia

SAMPLE

5. Which symptom is not classified under extrapyramidal symptoms?

A. Tardive dyskinesia

B. Hallucinations

C. Asynchrony

D. Acute dystonia

Hallucinations are not classified as extrapyramidal symptoms; rather, they are a psychiatric symptom typically associated with various mental health disorders, including schizophrenia and other psychotic conditions. Extrapyramidal symptoms (EPS) are side effects that arise from the use of certain antipsychotic medications, particularly first-generation (typical) antipsychotics, and affect motor control. Extrapyramidal symptoms primarily involve movement disorders, which include tardive dyskinesia, characterized by involuntary, repetitive movements often of the face or tongue, and acute dystonia, which presents as sustained muscle contractions, abnormal postures, and sometimes painful spasms. Asynchrony, while less commonly referenced, can be associated with dyskinetic movements, hence also regarded within the spectrum of EPS. The distinction is important because understanding the different types of symptoms and their classifications aids in effective diagnosis and treatment of side effects associated with antipsychotic medications. Recognizing hallucinations as a separate entity further emphasizes the complex nature of treatment in psychiatric disorders where both pharmacological and psychological approaches may be necessary.

6. In terms of pharmacokinetics, what is a key feature of dopaminergic drugs?

A. They have high lipid solubility

B. They are poorly absorbed

C. They bind tightly to plasma proteins

D. They are not metabolized in the liver

Dopaminergic drugs often exhibit high lipid solubility, which is a significant feature in pharmacokinetics. This characteristic enhances their ability to cross the blood-brain barrier, facilitating their action on the central nervous system where dopamine receptors are located. The ability to readily enter the brain is crucial for achieving the therapeutic effects desired in treating conditions such as Parkinson's disease and other dopamine-related disorders. High lipid solubility also contributes to a rapid onset of action and allows for varied routes of administration, including oral and transdermal methods. This feature is relevant because it can influence drug stability, distribution, and overall bioavailability in the system, impacting how effectively these drugs can exert their pharmacological effects. Other choices do not accurately reflect key features inherent to dopaminergic drugs. While some dopaminergic drugs may be absorbed to varying degrees and may bind to plasma proteins, stating they are poorly absorbed or have a general binding tendency is not universally applicable. Additionally, most dopaminergic drugs do undergo hepatic metabolism, which is a crucial aspect of their pharmacokinetics, making the statement regarding their metabolism misleading. Therefore, high lipid solubility stands out as a defining characteristic among dopaminergic medications.

SAMPLE

7. What component of carbidopa-levodopa inhibits the enzyme dopa decarboxylase?

A. Levodopa

B. Carbidopa

C. Ropinirole

D. Bromocriptine

Carbidopa is the component of the combination carbidopa-levodopa that inhibits the enzyme dopa decarboxylase. This inhibition is crucial because dopa decarboxylase is responsible for converting levodopa into dopamine outside of the brain, particularly in the periphery. By blocking this enzyme, carbidopa ensures that more levodopa is available to cross the blood-brain barrier and be converted into dopamine within the central nervous system. This pharmacological strategy enhances the effectiveness of levodopa therapy in treating conditions such as Parkinson's disease while minimizing side effects associated with peripheral dopamine production, such as nausea and cardiovascular effects. The other options refer to different medications and their mechanisms of action. Ropinirole and bromocriptine are dopamine receptor agonists and do not inhibit dopa decarboxylase. Levodopa is the precursor to dopamine and does not have any effect on the decarboxylation process itself. Thus, carbidopa's role is essential in maximizing the therapeutic effects of levodopa in the treatment of Parkinson's disease.

8. What is the antidote for benzodiazepine overdose?

A. Activated charcoal

B. Flumazenil

C. Diphenhydramine

D. Rivastigmine

Flumazenil is specifically designed to counteract the effects of benzodiazepines, making it the appropriate antidote for a benzodiazepine overdose. It works as a competitive antagonist at the benzodiazepine binding site on the GABA-A receptor, which can reverse the sedation and respiratory depression caused by benzodiazepines. In clinical practice, flumazenil can be administered to patients who present with a benzodiazepine overdose to effectively restore normal levels of consciousness and respiratory function, provided that the patient does not have a history of seizures or is not on other drugs that may induce seizures, as flumazenil can precipitate seizures in sensitive individuals. Other options do not serve the same purpose; for instance, activated charcoal may help limit absorption if given soon after ingestion of the benzodiazepine, but it does not reverse the effects of the drug. Diphenhydramine is an antihistamine with sedative properties and does not counteract the effects of benzodiazepines. Rivastigmine, a drug used in the management of Alzheimer's disease, works differently by inhibiting acetylcholinesterase and also does not address benzodiazepine overdose. Thus, flumazenil is the targeted

SAMPLE

9. Which COMT inhibitor is often used with levodopa in advanced Parkinson's Disease and may cause liver damage?

A. Entacapone

B. Tolcapone

C. Selegiline

D. Amantadine

Tolcapone is the COMT inhibitor commonly used in conjunction with levodopa for patients with advanced Parkinson's disease. It functions by inhibiting catechol-O-methyltransferase (COMT), an enzyme that metabolizes levodopa in the periphery, thereby increasing the availability of levodopa for conversion to dopamine in the brain. This is particularly useful in addressing the fluctuations in motor control experienced by patients in later stages of Parkinson's. One significant concern with tolcapone is its potential for hepatotoxicity; it has been associated with elevated liver enzymes and liver damage in some patients. Because of this risk, patients receiving tolcapone require regular monitoring of liver function tests to catch any signs of liver injury early. This risk distinguishes tolcapone from other COMT inhibitors like entacapone, which is also used with levodopa but has a much lower risk of liver-related side effects. Other options, such as selegiline and amantadine, work through different mechanisms and do not primarily inhibit COMT, which further solidifies the specific role of tolcapone in the treatment of advanced Parkinson's disease.

10. Which of the following is NOT true about MAO inhibitors?

A. They can cause hypertensive crisis with tyramine.

B. They are used to increase dopamine levels.

C. They selectively inhibit MAO-A only.

D. They can interact with a variety of medications.

The statement that MAO inhibitors selectively inhibit MAO-A only is not true. While some MAO inhibitors are selective for MAO-A, there are others that can inhibit both MAO-A and MAO-B. Non-selective MAO inhibitors affect both subtypes, which is important for their clinical use and interactions. MAO-A is primarily responsible for the metabolism of norepinephrine and serotonin, while MAO-B is involved more with dopamine and phenylethylamine. Thus, non-selective MAO inhibitors can influence a broader range of neurotransmitter systems, leading to their varied therapeutic effects and potential side effects. The other statements about MAO inhibitors are accurate reflections of their properties. For instance, they can indeed lead to hypertensive crises if foods rich in tyramine are consumed due to the inhibition of tyramine metabolism. Additionally, these medications help increase dopamine levels indirectly by preventing its breakdown. They also have the potential to interact with various other medications, making consideration of drug interactions crucial in clinical settings.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://pharmacology3cns.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE