PHARMACOLOGY MATERNITY HESI FINAL (EVOLVE) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://pharmmaternityhesievolve.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What medication is appropriate for a lactating woman diagnosed with migraine headaches who wishes to continue breastfeeding?**

   A. Ibuprofen

   B. Sumatriptan

   C. Acetaminophen

   D. Ergotamine

2. **A nurse is reviewing a postmenopausal client's history, which reveals that the client previously received hormonal therapy (HT) as treatment for osteoporosis. The nurse recognizes that HT increases the risk of which condition?**

   A. Ovarian cancer

   B. Heart disease

   C. Breast cancer

   D. Stroke

3. **What baseline assessment is essential before initiating an intravenous magnesium sulfate infusion?**

   A. Blood pressure

   B. Heart rate

   C. Respiratory rate

   D. Oxygen saturation

4. **What is the essential monitoring for a pregnant client receiving antiepileptic drugs?**

   A. Blood glucose levels

   B. Electrolyte balance

   C. Fetal heart rate and development

   D. Fluid intake

5. **What is the commonly used medication for the treatment of postpartum hemorrhage?**

   A. Methylergometrine

   B. Oxytocin

   C. Carboprost

   D. Ergometrine

SAMPLE

6. What are the primary concerns with using nonsteroidal anti-inflammatory drugs (NSAIDs) in late pregnancy?

A. Increased chance of labor

B. Potential acute kidney injury

C. Premature closure of the ductus arteriosus and increased bleeding risk

D. Enhanced uterine contractions

7. What type of drug is terbutaline commonly classified as?

A. Pain reliever

B. Antibiotic

C. Tocolytic agent

D. Antidepressant

8. What is the function of beta-mimetics in maternity care?

A. To induce uterine contractions

B. To relax smooth muscle and decrease uterine contractions

C. To promote fetal movement

D. To increase uterine muscle tone

9. Which medication is known to positively impact fetal growth in hypertensive mothers?

A. Hydrochlorothiazide

B. Methyldopa

C. Lisinopril

D. Atorvastatin

10. What is a common side effect of opioid analgesics used during labor?

A. Increased heart rate in the mother

B. Nausea and vomiting

C. Respiratory depression in the newborn

D. Elevated blood pressure

SAMPLE

Answers

SAMPLE

1. B
2. C
3. C
4. C
5. A
6. C
7. C
8. B
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. What medication is appropriate for a lactating woman diagnosed with migraine headaches who wishes to continue breastfeeding?

A. Ibuprofen

B. Sumatriptan

C. Acetaminophen

D. Ergotamine

Sumatriptan is a medication that is considered safe for use in lactating women who experience migraine headaches. As a selective serotonin receptor agonist belonging to the triptan class, sumatriptan is effective for the acute treatment of migraines. Studies have shown that it has minimal levels in breast milk and that its use does not pose significant risks to a breastfeeding infant. In contrast, some other treatments for migraines can have potential risks when it comes to lactation. Ergotamine, for instance, can inhibit lactation and may have adverse effects on a nursing infant, making it an inappropriate choice. While ibuprofen and acetaminophen are generally safe for treating mild to moderate headaches and are often used in breastfeeding women, they may not provide adequate relief for the specific symptoms and intensity of migraine headaches compared to sumatriptan. Thus, sumatriptan stands out as the most suitable option for a lactating woman who needs effective migraine management while continuing to breastfeed.

2. A nurse is reviewing a postmenopausal client's history, which reveals that the client previously received hormonal therapy (HT) as treatment for osteoporosis. The nurse recognizes that HT increases the risk of which condition?

A. Ovarian cancer

B. Heart disease

C. Breast cancer

D. Stroke

Hormonal therapy (HT) has been associated with an increased risk of breast cancer, especially in postmenopausal women. This is primarily due to the estrogen component in HT, which can promote the growth of hormone-sensitive tumors. The relationship between estrogen exposure and breast cancer risk has been established through various studies, indicating that women who receive long-term hormonal therapy may have a higher incidence of breast cancer compared to those who do not. When evaluating other potential risks associated with HT, such as heart disease and stroke, research findings provide mixed results. While there is some evidence that hormonal therapy may have a protective effect on heart health when initiated close to the onset of menopause, the overall increased risk of breast cancer is more clearly defined. The relationship to ovarian cancer is less established in this context, and while heart disease is a consideration, it does not directly relate to the prominent association with breast cancer risk in patients receiving HT. Thus, recognizing that HT specifically heightens the risk for breast cancer is crucial for healthcare providers when discussing treatment options with postmenopausal clients.

SAMPLE

3. What baseline assessment is essential before initiating an intravenous magnesium sulfate infusion?

A. Blood pressure

B. Heart rate

C. Respiratory rate

D. Oxygen saturation

Before initiating an intravenous magnesium sulfate infusion, measuring the respiratory rate is essential. Magnesium sulfate has a known effect on neuromuscular transmission and can lead to respiratory depression, particularly at higher serum magnesium levels. Monitoring the respiratory rate helps ensure that the patient can maintain adequate ventilation and provides an early indication of potential adverse effects related to magnesium therapy. By establishing a baseline respiratory rate, healthcare providers can make informed decisions about continuing treatment or implementing interventions if a significant decrease in the patient's respiratory function is observed after the infusion begins. This precaution is crucial in a maternity setting, where patients may be at higher risk for complications during labor and delivery.

4. What is the essential monitoring for a pregnant client receiving antiepileptic drugs?

A. Blood glucose levels

B. Electrolyte balance

C. Fetal heart rate and development

D. Fluid intake

Monitoring fetal heart rate and development is essential for a pregnant client receiving antiepileptic drugs because these medications can have significant effects on both the mother and the developing fetus. Some antiepileptic drugs are known teratogens and can lead to potential fetal developmental issues if not monitored carefully. Regular assessment of the fetal heart rate helps ensure that the fetus is not experiencing distress, which could result from maternal medication use. In addition, monitoring fetal development through ultrasounds or other imaging techniques can help detect any potential congenital anomalies or developmental delays earlier, allowing for prompt management or support options as needed. While blood glucose levels, electrolyte balance, and fluid intake monitoring are important in various clinical scenarios, they are not specifically the focus when considering the impacts of antiepileptic drugs during pregnancy. Instead, the priority is to safeguard fetal well-being and assess the potential effects of the medications prescribed to the mother.

SAMPLE

5. What is the commonly used medication for the treatment of postpartum hemorrhage?

A. Methylergometrine

B. Oxytocin

C. Carboprost

D. Ergometrine

The commonly used medication for the treatment of postpartum hemorrhage is oxytocin. This medication is a synthetic form of the naturally occurring hormone oxytocin, which plays a critical role in promoting uterine contractions. During the postpartum period, the delivery of the placenta can lead to uterine atony, or lack of tone, which is a primary cause of postpartum hemorrhage. By stimulating uterine contractions, oxytocin effectively helps to reduce the risk of excessive bleeding by facilitating the uterus's ability to contract and expel any retained placenta. While methylergometrine, carboprost, and ergometrine can also be used in managing postpartum hemorrhage, they are typically employed in specific circumstances or as second-line agents after oxytocin. Methylergometrine and ergometrine work by causing more forceful uterine contractions but can have contraindications and side effects that must be carefully monitored. Carboprost, which is a prostaglandin, is effective primarily in cases of severe postpartum hemorrhage, particularly when other medications are ineffective. However, oxytocin remains the first-line treatment due to its effectiveness and safety profile, making it the most commonly used medication for this condition.

6. What are the primary concerns with using nonsteroidal anti-inflammatory drugs (NSAIDs) in late pregnancy?

A. Increased chance of labor

B. Potential acute kidney injury

C. Premature closure of the ductus arteriosus and increased bleeding risk

D. Enhanced uterine contractions

The primary concerns with using nonsteroidal anti-inflammatory drugs (NSAIDs) in late pregnancy are linked to the potential for premature closure of the ductus arteriosus and an increased risk of bleeding. The ductus arteriosus is a critical fetal blood vessel that connects the pulmonary artery to the aorta, allowing blood to bypass the lungs while in utero. NSAIDs can inhibit prostaglandin synthesis, which is necessary for keeping the ductus arteriosus open. If it closes too early, it can lead to significant fetal complications, such as pulmonary hypertension and impaired blood flow. Additionally, NSAIDs can impair platelet function, leading to an increased risk of bleeding during delivery or in the postpartum period. This heightened bleeding risk is particularly concerning in the context of both the mother's health and the safety of the newborn. While other options might hint at related issues, they do not represent the most significant concerns associated with NSAID use in the context of late pregnancy.

SAMPLE

7. What type of drug is terbutaline commonly classified as?

A. Pain reliever

B. Antibiotic

C. Tocolytic agent

D. Antidepressant

Terbutaline is classified as a tocolytic agent, which is a type of medication used to inhibit premature labor by relaxing the muscles of the uterus. Tocolytics are primarily employed in obstetrics to delay preterm labor, allowing time for other interventions, such as corticosteroids, to enhance fetal lung maturity or for transportation to a facility equipped to handle premature births. Terbutaline specifically functions as a beta-adrenergic agonist, mimicking the effects of adrenaline on the smooth muscle of the uterus. This action helps reduce contractions and can be critically important in managing preterm labor scenarios. Understanding terbutaline's role in this context is essential for providing appropriate care and intervention in maternity settings.

8. What is the function of beta-mimetics in maternity care?

A. To induce uterine contractions

B. To relax smooth muscle and decrease uterine contractions

C. To promote fetal movement

D. To increase uterine muscle tone

Beta-mimetics, also known as beta-agonists, primarily function to relax smooth muscle in the body, which includes the uterine muscle. In maternity care, they are often used as tocolytics, meaning they are prescribed to decrease uterine contractions, particularly in cases where preterm labor is a concern. By relaxing the uterine muscles, beta-mimetics help to delay labor, allowing for further fetal development and stabilization. This mechanism is crucial in managing preterm labor because it can provide added time for fetal maturation, particularly in critical areas such as lung development, which can significantly improve outcomes after birth. The use of beta-mimetics therefore reflects their therapeutic purpose in maternity care—helping to manage premature labor by inhibiting contractions rather than inducing them or increasing muscle tone. This understanding is key for healthcare professionals while treating expectant mothers facing the risk of early labor.

9. Which medication is known to positively impact fetal growth in hypertensive mothers?

A. Hydrochlorothiazide

B. Methyldopa

C. Lisinopril

D. Atorvastatin

Methyldopa is known to positively impact fetal growth in hypertensive mothers due to its effectiveness in managing hypertension while being one of the safer options during pregnancy. It acts as a centrally acting alpha-2 adrenergic agonist, leading to decreased vascular resistance and reduced blood pressure without compromising uteroplacental blood flow, which is crucial for fetal development. This medication has a long history of use in pregnant women and is associated with favorable outcomes when treating elevated blood pressure, helping to prevent complications such as intrauterine growth restriction. Maintaining appropriate blood pressure levels in hypertensive mothers is essential for ensuring adequate blood flow and nutrients to the fetus, thereby supporting healthy fetal growth. In contrast, some of the other medications listed have different implications during pregnancy. For instance, lisinopril is an ACE inhibitor that is contraindicated during pregnancy due to the risk of fetal harm, especially in the second and third trimesters. Hydrochlorothiazide, while used in some cases, may not have the same level of evidence supporting its safety and efficacy in promoting fetal growth. Atorvastatin, a statin used primarily for cholesterol management, is also not recommended during pregnancy as it poses risks to the fetus. Therefore, methyldopa

SAMPLE

10. What is a common side effect of opioid analgesics used during labor?

A. Increased heart rate in the mother

B. Nausea and vomiting

C. Respiratory depression in the newborn

D. Elevated blood pressure

Opioid analgesics are frequently used during labor to manage pain effectively. While these medications are beneficial for the mother, they also carry potential side effects that can affect both the mother and the newborn. One significant side effect associated with the use of opioids is the risk of respiratory depression, particularly in the newborn. Opioids cross the placental barrier, and if administered to the mother during labor, they may lead to decreased respiratory function in the newborn. This happens because the newborn's central nervous system is still developing and may be more sensitive to the effects of opioids. If the mother receives a higher dose or if the drug is given shortly before delivery, the newborn may experience respiratory depression, which can manifest as inadequate breathing effort or low oxygen levels at birth. This impact on the newborn is a critical consideration in managing labor pain with opioids, and healthcare providers often closely monitor both the mother and child during and after administration to mitigate risks associated with any potential side effects.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://pharmmaternityhesievolve.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE