
PHARMACY CONTINUING EDUCATION (CE) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://pharmacyce.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What factor contributes to 'inflammaging' in elderly individuals?

A. Increased stimulation of the adaptive immune system

B. Reduced integrity of the intestinal epithelial barrier

C. Decreased adipose tissue accumulation in the body

D. A shift to a chronic anti-inflammatory state with aging

2. What is the safest disposal recommendation for expired fentanyl patches?

A. Store them in a locked cabinet

B. Dispose by mixing in coffee grounds and throwing in the trash

C. Flushing them down the toilet

D. Wait for a drug take-back program

3. What was the percentage increase in the risk of all-cause mortality for patients with low health literacy compared to those with good health literacy?

A. 32% higher

B. 91% higher

C. 164% higher

D. Similar risk

4. For patients with persistent asthma symptoms, which item is NOT required before increasing therapy?

A. Check inhaler technique

B. Assure strict medication adherence

C. Recommend the addition of inhaled tiotropium to the provider

D. Inquire about allergen exposure and counsel on triggers

5. Which symptoms are categorized as typical for COVID-19?

A. Fever, cough, and shortness of breath

B. Shortness of breath, dizziness, and myalgia

C. Myalgia, cough, and sore throat

D. Sore throat, malaise, and fever

SAMPLE

6. What is a hallmark feature of fibrodysplasia ossificans progressiva?

A. Increased severity of muscle trauma

B. Loss of mobility

C. Heightened risk of intramuscular complications

D. Growth spurts leading to functional improvement

7. Which of the following is NOT an attribute of fibrodysplasia ossificans progressiva (FOP)?

A. Heterotopic ossification (HO)

B. Elongation of molar teeth

C. Malformation of the great toes

D. Progressive loss of mobility

8. Which medication is MOST concerning when starting tazemetostat for a patient also using verapamil?

A. QT prolongation risk

B. Lisinopril use

C. Proton pump inhibitor use

D. Verapamil use

9. What is the recommended product for dry eye symptoms in patients seeking relief?

A. Cyclosporine A eye drops

B. Tetrahydrozoline eye drops

C. Omega-3 fatty acid oral capsules

D. Polyethylene glycol eye drops

10. Which of the following is the first step for sexual harassment victims who wish to file a complaint?

A. Take legal action through EEOC

B. Speak to human resources

C. Confront the harasser directly

D. Collect and document evidence

SAMPLE

Answers

SAMPLE

1. B
2. C
3. B
4. C
5. A
6. B
7. B
8. D
9. D
10. D

SAMPLE

Explanations

SAMPLE

1. What factor contributes to 'inflammaging' in elderly individuals?

A. Increased stimulation of the adaptive immune system

B. Reduced integrity of the intestinal epithelial barrier

C. Decreased adipose tissue accumulation in the body

D. A shift to a chronic anti-inflammatory state with aging

The correct choice identifies reduced integrity of the intestinal epithelial barrier as a significant factor contributing to 'inflammaging' in elderly individuals. Inflammaging refers to the chronic low-grade inflammation that often accompanies aging, and the integrity of the intestinal barrier plays a crucial role in regulating systemic inflammation. As people age, the intestinal epithelial barrier can become compromised due to several factors, including changes in diet, gut microbiota, and an increase in the permeability of the gut lining. This disruption can lead to the translocation of bacteria and their products into the bloodstream, provoking an inflammatory response. Consequently, the immune system becomes activated, contributing to a sustained, low-grade state of inflammation. This chronic inflammatory state is associated with various age-related diseases, including cardiovascular disease, neurodegenerative disorders, and metabolic conditions. The other options do not adequately explain the phenomenon of inflammaging as it pertains specifically to aging. Increased stimulation of the adaptive immune system, for example, typically involves a heightened immune response rather than the chronic inflammation seen in aging. Similarly, decreased adipose tissue accumulation would often correlate with a reduction in inflammatory cytokines, rather than an increase. Lastly, a shift to a chronic anti-inflammatory state is counterproductive in the context of inflammaging because it describes a

2. What is the safest disposal recommendation for expired fentanyl patches?

A. Store them in a locked cabinet

B. Dispose by mixing in coffee grounds and throwing in the trash

C. Flushing them down the toilet

D. Wait for a drug take-back program

Flushing expired fentanyl patches down the toilet is not considered the safest disposal method. While some medications can be safely discarded in this manner, fentanyl patches are particularly potent opioids that pose significant risks of accidental exposure, overdose, and environmental contamination. The safest disposal recommendation for expired fentanyl patches is to wait for a drug take-back program. These programs are designed to allow individuals to return medication safely to designated collection sites, where it can be disposed of in an environmentally responsible manner. Drug take-back programs help prevent misuse and limit the potential for accidental exposure to potent medications, particularly in households with children or pets. Mixing the patches with coffee grounds and throwing them in the trash is often recommended for other medications but does not provide the same level of safety as a dedicated take-back program for high-risk substances such as fentanyl. Storing them in a locked cabinet, while it may prevent unauthorized access during the time the patches are still valid, does nothing to mitigate the risks associated with expired medications. Thus, it is critical to utilize drug take-back programs for the proper disposal of expired fentanyl patches.

SAMPLE

3. What was the percentage increase in the risk of all-cause mortality for patients with low health literacy compared to those with good health literacy?

A. 32% higher

B. 91% higher

C. 164% higher

D. Similar risk

The choice indicating a 91% higher risk of all-cause mortality for patients with low health literacy compared to those with good health literacy is supported by various studies demonstrating the significant impact of health literacy on health outcomes. Individuals with low health literacy often face challenges in understanding health information, navigating the healthcare system, and managing their health effectively. This complicates their ability to adhere to treatment plans, make informed decisions about their health, and recognize when to seek appropriate care. As a result, they may experience higher rates of complications, worsening chronic conditions, and ultimately increased mortality. Research has shown that patients with poor health literacy are more likely to have poorer health outcomes, which correlates with a higher risk of death. The 91% figure reflects a substantial body of evidence linking inadequate health literacy with negative health outcomes, emphasizing the importance of health communication and patient education in healthcare settings. By addressing literacy issues, healthcare providers can help mitigate these risks and improve overall patient health.

4. For patients with persistent asthma symptoms, which item is NOT required before increasing therapy?

A. Check inhaler technique

B. Assure strict medication adherence

C. Recommend the addition of inhaled tiotropium to the provider

D. Inquire about allergen exposure and counsel on triggers

Increasing therapy for persistent asthma symptoms typically involves a systematic approach to ensure that all current management strategies are being effectively utilized. To optimize asthma control, healthcare professionals must first assess several key factors concerning the patient's current treatment and environment. Checking inhaler technique is crucial as improper use can lead to insufficient drug delivery and suboptimal control of asthma symptoms. Ensuring that patients are using their inhalers correctly is a precondition before making any recommendations about therapy adjustments. Assuring strict medication adherence is also essential. Patients may not be experiencing adequate control of their asthma due to non-adherence to their prescribed medication regimen. It is necessary to confirm that patients are taking their medications as instructed before considering an increase in therapy. Inquiring about allergen exposure and counseling on triggers is a fundamental part of asthma management. Understanding and addressing potential environmental triggers can profoundly influence asthma control and guide appropriate therapy adjustments. Recommending the addition of inhaled tiotropium, however, does not fall under the required assessments before increasing therapy. While inhaled tiotropium can be beneficial for some patients as an add-on therapy, this particular recommendation can only be made after the initial evaluations and steps have been taken, including ensuring correct inhaler use, adherence to medications, and addressing environmental factors.

SAMPLE

5. Which symptoms are categorized as typical for COVID-19?

A. Fever, cough, and shortness of breath

B. Shortness of breath, dizziness, and myalgia

C. Myalgia, cough, and sore throat

D. Sore throat, malaise, and fever

The symptoms categorized as typical for COVID-19 primarily include fever, cough, and shortness of breath. These symptoms are commonly reported by individuals infected with the virus and have been recognized as key indicators in the early stages of the disease. Fever is a common response to infection, reflecting the body's natural defense mechanism. A cough, often dry, is another hallmark of COVID-19, representing respiratory involvement. Shortness of breath can indicate more severe respiratory compromise, which is frequently observed in patients as the disease progresses. Notably, these three symptoms encapsulate the core respiratory issues associated with the virus, making them critical for both diagnosis and initial triage when encountering a patient suspected of having COVID-19. In contrast, the other sets of symptoms listed may indeed be associated with COVID-19 but are not as specific or widely recognized as the classic trio of fever, cough, and shortness of breath. For instance, dizziness and myalgia can overlap with a range of other illnesses, making them less reliable as indicators for COVID-19 specifically. Hence, while individuals experiencing those symptoms could potentially have COVID-19, they do not represent the typical symptomatology as clearly as the first group.

6. What is a hallmark feature of fibrodysplasia ossificans progressiva?

A. Increased severity of muscle trauma

B. Loss of mobility

C. Heightened risk of intramuscular complications

D. Growth spurts leading to functional improvement

The hallmark feature of fibrodysplasia ossificans progressiva (FOP) is indeed a loss of mobility. FOP is a rare genetic condition characterized by the abnormal formation of bone in muscles, tendons, and other connective tissues, a process known as heterotopic ossification. This condition typically progresses over time, leading to a gradual restriction of movement. As new bone forms in areas where it should not, patients experience increasing stiffness and immobility, particularly affecting joints and limbs. Loss of mobility is a significant concern for individuals with FOP, resulting from the ossified tissues limiting their range of motion and ability to perform everyday activities. The condition can lead to severe disability, and managing mobility becomes a central focus of care as the disease progresses. Other options provide relevant information but do not encapsulate the primary characteristic of FOP as effectively. For example, while increased severity of muscle trauma may occur due to an increased risk of flare-ups or injuries, it is not as defining of the condition as the progressive loss of mobility. Heightened risk of intramuscular complications can be more related to the development of ossification and muscle behavior after injury, but still lacks the clarity offered by the primary mobility focus. Growth spurts

SAMPLE

7. Which of the following is NOT an attribute of fibrodysplasia ossificans progressiva (FOP)?

A. Heterotopic ossification (HO)

B. Elongation of molar teeth

C. Malformation of the great toes

D. Progressive loss of mobility

Fibrodysplasia ossificans progressiva (FOP) is a rare genetic disorder characterized primarily by the formation of heterotopic ossification (the abnormal growth of bone in soft tissues) following injury or spontaneously. Among the attributes associated with FOP, its hallmark feature is indeed heterotopic ossification, along with progressive loss of mobility as the condition progresses and muscle and connective tissues gradually transform into bone. The elongation of molar teeth is not recognized as an attribute of FOP. While patients may experience various dental developments or issues over time, elongation of molar teeth is not a characteristic symptom or feature of this condition. Instead, malformation of the great toes is a noted physical trait in individuals with FOP, often presenting as an abnormality in the shape or orientation of the toes. Understanding these attributes allows for better identification and management of this complex condition, enabling healthcare professionals to support affected patients effectively by focusing on their unique needs throughout the progression of FOP.

8. Which medication is MOST concerning when starting tazemetostat for a patient also using verapamil?

A. QT prolongation risk

B. Lisinopril use

C. Proton pump inhibitor use

D. Verapamil use

When considering the use of tazemetostat in conjunction with verapamil, the concern primarily arises from the pharmacokinetic interactions between the two medications. Verapamil is a calcium channel blocker that is known to inhibit certain cytochrome P450 enzymes, particularly CYP3A4. Tazemetostat, an EZH2 inhibitor used in certain cancers, is extensively metabolized by CYP3A4. When a patient is already on verapamil, the use of tazemetostat can lead to increased serum levels of the latter due to the inhibition of its metabolism. This can increase the risk of adverse effects associated with tazemetostat, potentially leading to toxicity. Therefore, monitoring and consideration of dose adjustment or alternative treatment strategies are essential when initiating tazemetostat in a patient who is taking verapamil. The potential for QT prolongation, the use of lisinopril, and proton pump inhibitors, while relevant in their own contexts, do not present the same level of concern in this scenario as the interaction between tazemetostat and verapamil. Thus, careful evaluation of the patient's medication regimen, especially concerning verapamil, is vital when starting tazemetostat therapy.

SAMPLE

9. What is the recommended product for dry eye symptoms in patients seeking relief?

A. Cyclosporine A eye drops

B. Tetrahydrozoline eye drops

C. Omega-3 fatty acid oral capsules

D. Polyethylene glycol eye drops

Polyethylene glycol eye drops are recommended for dry eye symptoms because they act as an effective lubricant for the eyes. This class of eye drops helps to moisturize the ocular surface, providing immediate relief from dryness and irritation by maintaining tear film stability. Polyethylene glycol is particularly beneficial as it can create a soothing effect, reduce friction, and support overall eye comfort. In contrast, cyclosporine A eye drops are typically used in cases of moderate to severe dry eye, particularly when inflammation is a significant factor, and are not the first-line treatment for mild symptoms. Tetrahydrozoline eye drops are an antihistamine and decongestant that may provide temporary relief of redness but do not address the underlying dryness. Omega-3 fatty acid oral capsules can support eye health and may improve dry eye symptoms over time but are not an immediate topical treatment for dryness. Thus, polyethylene glycol eye drops are the most suitable option for patients seeking immediate relief from dry eye symptoms.

10. Which of the following is the first step for sexual harassment victims who wish to file a complaint?

A. Take legal action through EEOC

B. Speak to human resources

C. Confront the harasser directly

D. Collect and document evidence

The first step for sexual harassment victims who wish to file a complaint involves collecting and documenting evidence. This is a crucial initial action because it provides a solid foundation for any allegations made against the harasser. Detailed documentation can include records of incidents, dates, times, locations, witnesses, and any communication related to the harassment. This evidence not only strengthens the victim's claim but also helps establish a pattern of behavior if there are multiple incidents. By gathering this information first, victims can present a well-organized account of their experiences when they decide to escalate the complaint process, whether that involves speaking to human resources, taking legal action, or confronting the harasser. Proper documentation also helps ensure that the complaint is taken seriously and can lead to a more effective resolution of the situation.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://pharmacyce.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE