PHYSICAL THERAPY ASSISTANT PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://pta-exam.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **During the Neer impingement test, what is the primary source of pain if the test is positive?**

   A. Compression of the greater tuberosity against the anterior acromion

   B. Stretching of the rotator cuff muscles

   C. Inflammation of the biceps tendon

   D. Bruising of the acromion process

2. **Which phenomenon describes alternating muscle contractions and relaxations in response to a sustained stretch?**

   A. Muscle spasticity

   B. Myokymia

   C. Clonus

   D. Muscle cramping

3. **Diminished deep tendon reflexes (hyporeflexia) are characteristic of which type of disorders?**

   A. Upper motor neuron disorders

   B. Lower motor neuron disorders

   C. Cerebellar disorders

   D. Peripheral nerve disorders

4. **Which laboratory test is most appropriate to confirm suspected hypercapnia in a patient?**

   A. Pulmonary function tests

   B. Arterial blood gas analysis

   C. Chest X-ray

   D. Sputum culture

5. **What characterizes Morton's neuroma?**

   A. An inflammation of the ankle joint

   B. A benign tumor affecting the first metatarsal

   C. A benign neuroma affecting the third interdigital nerve

   D. An injury to the Achilles tendon

SAMPLE

6. Which condition is most likely to result post-operatively if a patient's hemoglobin level is low after coronary artery bypass graft surgery?

A. Hypertension

B. Hyperglycemia

C. Hypoxia

D. Hypotension

7. What is a possible cause for early knee flexion from midstance through preswing in a patient with a transtibial amputation?

A. The socket is aligned in excessive flexion

B. The prosthetic foot is too dorsiflexed

C. The patient is lacking hip extension strength

D. The knee mechanism is locked

8. Which of the following is true about a cystocele?

A. It involves the urinary bladder.

B. It commonly occurs post-menopause.

C. It is less common than uterine prolapse.

D. It primarily affects men.

9. Where is the most appropriate location to attach an electrode for neuromuscular electrical stimulation of the peroneus longus?

A. Posteriorly on the calf

B. On the anterolateral surface of the lower leg

C. On the medial aspect of the thigh

D. Over the foot's dorsum

10. Which grade characterizes a patient completing more than half the range of motion but having insufficient strength to maintain the test position?

A. Good

B. Fair

C. Fair minus

D. Good minus

SAMPLE

Answers

SAMPLE

1. A
2. C
3. B
4. B
5. C
6. C
7. A
8. A
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. During the Neer impingement test, what is the primary source of pain if the test is positive?

A. Compression of the greater tuberosity against the anterior acromion

B. Stretching of the rotator cuff muscles

C. Inflammation of the biceps tendon

D. Bruising of the acromion process

The Neer impingement test is designed to assess the presence of shoulder impingement, often related to conditions affecting the rotator cuff or subacromial space. When this test is positive, the primary source of pain is due to the compression of the greater tuberosity against the anterior acromion. This occurs during shoulder flexion with internal rotation, which narrows the space between the acromion and the greater tuberosity of the humeral head. If there are inflamed tendons or structures within the subacromial space, such as the supraspinatus tendon or the subacromial bursa, this compression will precipitate pain. The positive test indicates that either structural changes or inflammation is leading to this mechanical compression during the movement. While stretching of the rotator cuff muscles and inflammation of the biceps tendon can be related to shoulder pain, they are not the primary mechanisms assessed by the Neer test. Bruising of the acromion process is less relevant as the test focuses on the dynamics of movement between the greater tuberosity and the acromion rather than direct injuries or contusions. Therefore, understanding the mechanism of compression helps clarify why this source of pain is the focal point

2. Which phenomenon describes alternating muscle contractions and relaxations in response to a sustained stretch?

A. Muscle spasticity

B. Myokymia

C. Clonus

D. Muscle cramping

The phenomenon that describes alternating muscle contractions and relaxations in response to a sustained stretch is clonus. Clonus is characterized by rhythmic, repetitive contractions that occur as a response to an external stimulus, such as a stretch. This involuntary neuromuscular response signifies an upper motor neuron lesion and can be observed in conditions such as multiple sclerosis, spinal cord injuries, or after a stroke. In contrast, muscle spasticity refers to increased muscle tone and exaggerated deep tendon reflexes, leading to stiff and jerky movements, but it does not describe the rhythmic alternating contractions seen in clonus. Myokymia is the term used for involuntary muscle contractions that appear as slow, wave-like movements in muscle fibers, which differ from the rhythmic patterns associated with clonus. Muscle cramping involves sustained involuntary contractions but does not involve the alternating pattern characteristic of clonus; instead, cramps often result in pain and a prolonged tightness in the muscle involved. Therefore, the correct identification of clonus aligns with its definitive pattern of alternation during sustained stretch responses.

SAMPLE

3. Diminished deep tendon reflexes (hyporeflexia) are characteristic of which type of disorders?

A. Upper motor neuron disorders

B. Lower motor neuron disorders

C. Cerebellar disorders

D. Peripheral nerve disorders

Diminished deep tendon reflexes, or hyporeflexia, are indeed characteristic of lower motor neuron disorders. This phenomenon occurs because lower motor neurons are responsible for transmitting signals from the spinal cord to the muscles. When these neurons are damaged or dysfunctional, either due to injury, disease, or degeneration, their ability to effectively convey reflexive movements is impaired. As a result, reflex actions, which are typically swift and involuntary, become less pronounced or absent altogether. In contrast, upper motor neuron disorders typically present with hyperreflexia, which means increased reflex responses. This is due to the disruption of the descending pathways that normally modulate reflex activity, leading to exaggerated reflexes. Cerebellar disorders do not primarily affect deep tendon reflexes, as cerebellar function is more related to coordination and balance than to the reflex arcs being assessed. Peripheral nerve disorders can lead to hyporeflexia as well, but they are not exclusively characterized by it; they can also manifest symptoms such as pain or muscle weakness. Lower motor neuron involvement is more definitive for the specific presentation of diminished reflexes, which is why it is identified as the correct response.

4. Which laboratory test is most appropriate to confirm suspected hypercapnia in a patient?

A. Pulmonary function tests

B. Arterial blood gas analysis

C. Chest X-ray

D. Sputum culture

To confirm suspected hypercapnia, arterial blood gas analysis is the most appropriate laboratory test. Hypercapnia refers to an elevated level of carbon dioxide (CO_2) in the blood, which can occur in various respiratory conditions. Arterial blood gas analysis provides precise measurements of oxygen (O_2), carbon dioxide (CO_2), pH, and bicarbonate levels, allowing healthcare providers to assess the respiratory function and metabolic status of the patient. By evaluating the levels of CO_2 in the arterial blood, this test can effectively indicate whether a patient is experiencing hypercapnia and to what extent. Other tests, while useful in different contexts, do not directly measure CO_2 levels. For example, pulmonary function tests assess lung capacity and airflow, but they do not provide information about blood gas levels. A chest X-ray can reveal structural issues in the lungs but cannot measure gas exchange or CO_2 levels. Lastly, sputum culture is used to identify infections or pathogens in the respiratory tract and does not relate to the measurement of carbon dioxide in the blood. Thus, arterial blood gas analysis stands out as the definitive test for confirming hypercapnia in a clinical setting.

SAMPLE

5. What characterizes Morton's neuroma?

A. An inflammation of the ankle joint

B. A benign tumor affecting the first metatarsal

C. A benign neuroma affecting the third interdigital nerve

D. An injury to the Achilles tendon

Morton's neuroma is characterized as a benign neuroma that primarily affects the third interdigital nerve, which is located between the third and fourth toes of the foot. This condition arises due to the thickening of the tissue surrounding the nerve, often resulting from compression or irritation. Patients typically experience symptoms such as pain, tingling, or a burning sensation in the affected area, which can often be exacerbated by wearing tight shoes or engaging in activities that put pressure on the forefoot. The choice that describes a benign tumor affecting the first metatarsal does not accurately represent Morton's neuroma, as it specifically involves the interdigital nerve rather than a tumor on the metatarsal itself. Additionally, an inflammation of the ankle joint and an injury to the Achilles tendon pertain to entirely different conditions unrelated to Morton's neuroma, which focuses on a specific nerve issue within the forefoot. The specificity of Morton's neuroma to the third interdigital nerve helps clarify the nature of the condition and emphasizes its distinct presentation in contrast to other foot-related ailments.

6. Which condition is most likely to result post-operatively if a patient's hemoglobin level is low after coronary artery bypass graft surgery?

A. Hypertension

B. Hyperglycemia

C. Hypoxia

D. Hypotension

A low hemoglobin level after coronary artery bypass graft surgery can significantly impact oxygen delivery to tissues throughout the body. Hemoglobin, which is found in red blood cells, plays a crucial role in transporting oxygen from the lungs to various tissues and returning carbon dioxide from the tissues back to the lungs. When hemoglobin levels are low, the oxygen-carrying capacity of the blood is reduced, leading to a condition known as hypoxia, where tissues do not receive adequate oxygen for proper function. Hypoxia can result in a variety of clinical issues, including fatigue, confusion, and even organ dysfunction, as the body struggles to meet its metabolic demands without sufficient oxygen. This is particularly critical in the post-operative period following a major surgery such as coronary artery bypass grafting, where the body's demand for oxygen may increase due to surgery stress and recovery processes. In contrast, hypertension, hyperglycemia, and hypotension do not directly correlate to low hemoglobin levels in the same manner as hypoxia. Hypertension involves high blood pressure, which is generally unrelated to hemoglobin levels, and hyperglycemia refers to elevated glucose levels, often associated with conditions like diabetes, rather than being tied directly to hemoglobin. Hypotension, while it may occur post-surgery

SAMPLE

7. What is a possible cause for early knee flexion from midstance through preswing in a patient with a transtibial amputation?

A. The socket is aligned in excessive flexion

B. The prosthetic foot is too dorsiflexed

C. The patient is lacking hip extension strength

D. The knee mechanism is locked

The alignment of the socket in excessive flexion is a significant factor that can lead to early knee flexion during the midstance through preswing phases for individuals with a transtibial amputation. When the socket is positioned too far forward in relation to the foot, it can create a mechanical disadvantage that encourages the knee to flex prematurely as the individual begins to transition their weight forward. This atypical gait pattern can contribute to instability and an inefficient walking style, as the prosthesis is not functioning optimally due to its alignment. In contrast, if a prosthetic foot is too dorsiflexed or if the patient lacks hip extension strength, these conditions may lead to different gait compensations rather than early knee flexion. A locked knee mechanism would prevent any knee motion altogether rather than causing early flexion. Therefore, socket alignment plays a crucial role in optimizing the gait mechanics and overall function of a transtibial prosthesis, making it an essential consideration in rehabilitation and prosthetic fitting.

8. Which of the following is true about a cystocele?

A. It involves the urinary bladder.

B. It commonly occurs post-menopause.

C. It is less common than uterine prolapse.

D. It primarily affects men.

A cystocele is a condition that specifically involves the urinary bladder. It occurs when the wall between the bladder and the vagina weakens, causing the bladder to bulge into the vaginal wall. This can lead to discomfort, urinary issues, and other complications. Understanding the nature of a cystocele is crucial for recognizing its symptoms and effects on urinary function, particularly in women, as it is often associated with childbirth and conditions that increase intra-abdominal pressure. While it is also true that a cystocele frequently occurs in post-menopausal women due to hormonal changes that affect connective tissue strength, the defining characteristic of a cystocele is its direct involvement with the urinary bladder itself.

SAMPLE

9. Where is the most appropriate location to attach an electrode for neuromuscular electrical stimulation of the peroneus longus?

A. Posteriorly on the calf

B. On the anterolateral surface of the lower leg

C. On the medial aspect of the thigh

D. Over the foot's dorsum

The most appropriate location to attach an electrode for neuromuscular electrical stimulation (NMES) of the peroneus longus is on the anterolateral surface of the lower leg. This muscle plays a crucial role in foot eversion and plantar flexion, and its anatomical position allows for effective stimulation when electrodes are placed in this area. Attaching the electrode to the anterolateral surface allows the electrical currents to effectively target the peroneus longus due to its proximity to the surface and its path along the lateral aspect of the lower leg. This placement maximizes the stimulation of the motor points associated with the muscle, ensuring optimal activation and resulting contraction. In contrast, other locations such as the posterior calf would primarily target the gastrocnemius and soleus muscles. The medial aspect of the thigh is related to adductor muscles, not the peroneus longus, and the dorsum of the foot would be too distal to effectively stimulate the peroneus longus, as its primary action is coordinated through the lower leg. Therefore, positioning the electrode on the anterolateral surface is essential for achieving the desired therapeutic effects of NMES on the peroneus longus.

10. Which grade characterizes a patient completing more than half the range of motion but having insufficient strength to maintain the test position?

A. Good

B. Fair

C. Fair minus

D. Good minus

The classification pertains to the grading of muscle strength and function, particularly in how well a patient can perform tasks related to their physical capabilities. When a patient can complete more than half of the range of motion but lacks the necessary strength to maintain that position, this indicates that their strength is below what is typically expected for a fair grade. "Fair" strength indicates that a patient can complete full range of motion against gravity but has some degree of weakness. When a patient surpasses half the range of motion but cannot sustain the test position, it reflects a level of strength that is insufficient for the full range, distinguishing it as "Fair minus." This grade captures the patient's ability to perform a majority of the motion but denotes a notable weakness, as they lack the strength to hold the position through the entire range against resistance or gravity. Grading scales provide a structured way to assess a patient's functional capability. In this scenario, identifying the subtleties in muscle strength gradings is essential for appropriate treatment planning and progress tracking.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://pta-exam.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE