
PHYSICIAN ASSISTANT CLINICAL KNOWLEDGE RATING AND ASSESSMENT TOOL (PACKRAT) 3 PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://packrat3.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **A 60-year-old male presents with a one-centimeter firm nodule at the right temple with heavy keratinization. There is no fluctuance. Which is the most likely diagnosis?**

   A. Actinic keratosis

   B. Squamous cell cancer

   C. Granuloma annulare

   D. Merkel cell carcinoma

2. **A non-displaced mid-shaft ulnar fracture on imaging is best treated with which splint?**

   A. Thumb spica

   B. Sugar tong

   C. Cock-up wrist

   D. Short arm gutter

3. **In acute pericarditis, which complication is most commonly observed?**

   A. Pericardial effusion

   B. Left ventricular failure

   C. Superior vena cava syndrome

   D. Subclavian steal syndrome

4. **A 53-year-old female massage therapist presents with sudden swelling over the tip of the elbow. The exam shows a 4 cm fluid-filled mass tender to palpation with no erythema or warmth. What is the most appropriate initial management?**

   A. Rest and NSAIDs

   B. Surgical excision

   C. Incision and drainage

   D. Aspiration and corticosteroid injection

5. **Why are monoamine oxidase inhibitors not considered first-line therapy for major depression?**

   A. Hypertensive crisis with tyramine-containing foods

   B. They are less safe than SSRIs

   C. They have no drug interactions

   D. They have lower efficacy

SAMPLE

6. Which finding would most strongly argue against active tuberculosis in a patient with suspected TB?

A. Negative PPD

B. Weight Loss

C. Night Sweats

D. Cough with Hemoptysis

7. A patient with sudden chest pain and dyspnea has tracheal deviation to the left, faint breath sounds on the right, and the right chest is hyperresonant. The most appropriate initial intervention is which?

A. Tracheostomy

B. Chest tube with underwater seal, left 2nd intercostal space

C. Needle thoracostomy right 2nd intercostal space

D. Lung scan for pulmonary embolus and begin heparin therapy

8. What is the most common cause of portal hypertension leading to esophageal varices?

A. Cirrhosis

B. Hepatic abscess

C. Hepatic adenoma

D. Alcoholic hepatitis

9. Hard exudates in the retina are described as which of the following?

A. Drusen

B. Cotton-wool patches

C. Yellowish bright lesions with well-defined borders

D. Preretinal hemorrhages

10. Which radiographic finding is commonly seen in bronchiectasis?

A. Normal

B. Platelike atelectasis with tram lines

C. Cavitation

D. Pleural effusion

SAMPLE

Answers

SAMPLE

1. B
2. A
3. A
4. A
5. A
6. A
7. C
8. A
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. A 60-year-old male presents with a one-centimeter firm nodule at the right temple with heavy keratinization. There is no fluctuance. Which is the most likely diagnosis?

A. Actinic keratosis

B. Squamous cell cancer

C. Granuloma annulare

D. Merkel cell carcinoma

In sun-exposed skin of an older adult, a firm, keratin-filled nodule is most suggestive of a squamous cell carcinoma arising from epidermal keratinocytes. The heavy keratinization reflects the tumor's keratin production, a hallmark of SCC, which often presents as a firm, possibly crusted or hyperkeratotic nodule on the face, ears, or temple. Actinic keratosis, by contrast, is a premalignant lesion that typically looks like a rough, scaly patch or plaque rather than a true firm nodule. Merkel cell carcinoma tends to present as a rapidly enlarging, painless, firm nodule that is usually flesh-colored to red or violaceous and not keratinized. Granuloma annulare appears as benign, annular patches or plaques rather than a solitary keratinized nodule. The absence of fluctuance argues against an abscess or infected cyst, reinforcing a neoplastic process rather than an infectious one. Given the patient's age, sun exposure, location, and the lesion's characteristics, squamous cell carcinoma is the most likely diagnosis. A biopsy would be indicated to confirm and guide treatment.

2. A non-displaced mid-shaft ulnar fracture on imaging is best treated with which splint?

A. Thumb spica

B. Sugar tong

C. Cock-up wrist

D. Short arm gutter

When a forearm fracture is non-displaced and confined to the mid-shaft of the ulna, the aim is to restrict movement at the fracture site enough to allow cast or splint healing while avoiding unnecessary immobilization of the entire arm. A thumb spica splint accomplishes this by stabilizing the wrist and thumb, which helps limit forearm rotation and bending motions that could disturb the ulna fracture, yet it keeps the elbow free to move and allows easy access for neurovascular checks and comfort. This makes it a practical and effective initial immobilization in the ED for a stable, non-displaced ulna fracture. The other splints don't provide the same targeted stabilization: a cock-up wrist splint mainly immobilizes the wrist with little forearm control; a short arm gutter mainly supports distal forearm injuries; a sugar-tong splint offers more comprehensive forearm immobilization but is bulkier and less comfortable in some cases.

SAMPLE

3. In acute pericarditis, which complication is most commonly observed?

A. Pericardial effusion

B. Left ventricular failure

C. Superior vena cava syndrome

D. Subclavian steal syndrome

The concept being tested is that acute pericarditis most often leads to a build-up of fluid in the pericardial space. Inflammation of the pericardium increases vascular permeability and drives fluid exudation into the surrounding cavity, so a pericardial effusion is the most frequent complication seen with acute pericarditis. This effusion can be small and asymptomatic, or large enough to cause symptoms or even tamponade if it accumulates rapidly and restricts the heart's ability to fill. The other choices describe problems that aren't typical direct complications of pericardial inflammation: left ventricular failure is a primary pumping problem rather than a pericardial complication; superior vena cava syndrome involves obstruction of the SVC, usually from other mediastinal processes; and subclavian steal syndrome arises from proximal arterial flow issues, not the pericardium.

4. A 53-year-old female massage therapist presents with sudden swelling over the tip of the elbow. The exam shows a 4 cm fluid-filled mass tender to palpation with no erythema or warmth. What is the most appropriate initial management?

A. Rest and NSAIDs

B. Surgical excision

C. Incision and drainage

D. Aspiration and corticosteroid injection

This scenario most fits aseptic olecranon bursitis from repetitive elbow pressure. A fluid-filled bump over the tip of the elbow that is tender but without warmth, erythema, or systemic signs points away from infection and toward inflammatory bursitis. The best first step is conservative management: rest the elbow, use NSAIDs for pain and inflammation, and avoid activities or pressures that irritate the bursa. This approach often reduces swelling and discomfort without the risks of invasive procedures. Infection would change the plan—septic bursitis would present with warmth, redness, fever, or rapid progression, and then aspiration with culture and antibiotics would be appropriate. Procedures like incision and drainage or corticosteroid injections are reserved for complicated, recurrent, or septic cases, not as initial management for a presumed aseptic bursitis. Surgical excision is typically a later option for chronic or recurrent bursitis when conservative therapy fails.

SAMPLE

5. Why are monoamine oxidase inhibitors not considered first-line therapy for major depression?

A. Hypertensive crisis with tyramine-containing foods

B. They are less safe than SSRIs

C. They have no drug interactions

D. They have lower efficacy

Monoamine oxidase inhibitors are avoided as first-line therapy because they carry a high risk of dangerous interactions, especially with tyramine-containing foods. MAO enzymes normally break down tyramine; when this enzyme is inhibited, tyramine can cause a rapid release of norepinephrine, leading to a hypertensive crisis. This dietary interaction—often called the "cheese reaction" when foods like aged cheeses, cured meats, and certain wines are involved—creates a serious, potentially life-threatening safety concern that makes these drugs far less convenient and riskier to use than newer antidepressants like SSRIs or SNRIs. In addition, MAO inhibitors have numerous drug interactions and require strict dietary restrictions and monitoring, further limiting their practicality as first-line therapy. While they can be effective in some patients, these safety and management challenges explain why they aren't chosen initially.

6. Which finding would most strongly argue against active tuberculosis in a patient with suspected TB?

A. Negative PPD

B. Weight Loss

C. Night Sweats

D. Cough with Hemoptysis

Tuberculin skin testing shows whether the immune system has been sensitized to TB. A negative test would argue against TB infection, making active disease less likely in this scenario. However, the test isn't perfect: false negatives can occur if the person is anergic or immunocompromised, in very early infection, or in very young or elderly patients. So while a negative PPD lowers the probability of active TB, it does not completely rule it out. The other findings—weight loss, night sweats, and coughing with hemoptysis—are classic symptoms that support TB and would favor continuing evaluation for active disease. If suspicion remains high despite a negative PPD, proceed with imaging and microbiologic testing (sputum smear, culture, and NAAT).

SAMPLE

7. A patient with sudden chest pain and dyspnea has tracheal deviation to the left, faint breath sounds on the right, and the right chest is hyperresonant. The most appropriate initial intervention is which?

A. Tracheostomy

B. Chest tube with underwater seal, left 2nd intercostal space

C. Needle thoracostomy right 2nd intercostal space

D. Lung scan for pulmonary embolus and begin heparin therapy

This scenario shows a tension pneumothorax. The signs—tracheal deviation away from the affected side, diminished or absent breath sounds on the involved side, and a hyperresonant chest—reflect air trapped under pressure that is shifting the mediastinum and impairing venous return. The immediate priority is to rapidly relieve that pressure to restore circulation and oxygenation. The fastest, life-saving step is needle thoracostomy in the right second intercostal space (midclavicular line) to decompress the pleural space. This provides immediate relief of the pressure, allows the lung to re-expand, and buys time before a definitive chest tube is placed. After decompression, a chest tube is typically inserted on the same side to provide ongoing drainage and re-expansion. Why the other options aren't appropriate right now: a tracheostomy would not address the underlying air buildup in the pleural space and isn't indicated emergently. A chest tube on the left side would not treat the problem on the right. Looking for a pulmonary embolism with a lung scan and starting heparin delays crucial decompression and is not appropriate in a suspected tension pneumothorax.

8. What is the most common cause of portal hypertension leading to esophageal varices?

A. Cirrhosis

B. Hepatic abscess

C. Hepatic adenoma

D. Alcoholic hepatitis

Cirrhosis is the most common cause of portal hypertension leading to esophageal varices because the fibrotic scarring and regenerative nodules distort hepatic architecture and markedly increase resistance to portal blood flow. This elevated portal venous pressure drives blood to form collateral vessels, including those in the esophagus, resulting in varices that can rupture and bleed. The other conditions listed don't typically cause sustained portal hypertension: a hepatic abscess is an acute infection that doesn't chronically obstruct portal inflow; a hepatic adenoma is a benign tumor without major impact on portal pressure; alcoholic hepatitis can contribute to cirrhosis, but the persistent portal hypertension arises most often once cirrhosis has developed.

9. Hard exudates in the retina are described as which of the following?

A. Drusen

B. Cotton-wool patches

C. Yellowish bright lesions with well-defined borders

D. Preretinal hemorrhages

Hard exudates are lipid-rich deposits that leak from damaged retinal capillaries, commonly seen in conditions like diabetic or hypertensive retinopathy. They appear as yellowish, bright, well-defined, refractile spots in the retina, often near the macula, which is why the description "yellowish bright lesions with well-defined borders" fits best. Other signs have different appearances: drusen are extracellular lipid deposits between the RPE and Bruch's membrane and tend to have softer, fuzzier borders; cotton-wool patches are fluffy white nerve fiber layer infarcts; preretinal hemorrhages are blood, giving boat-shaped or flame-shaped patterns rather than yellowish spots.

SAMPLE

10. Which radiographic finding is commonly seen in bronchiectasis?

A. Normal

B. Platelike atelectasis with tram lines

C. Cavitation

D. Pleural effusion

Bronchiectasis causes chronic dilation and thickening of the airways from repeated infections and inflammation. On a chest radiograph, the most characteristic finding is plate-like (subsegmental) atelectasis accompanied by tram-line markings. The plate-like atelectasis comes from mucus plugging and localized collapse in dependent lung regions, while the tram lines are the parallel, thickened bronchial walls that appear as linear densities. Cavitation and pleural effusion are not typical hallmarks of bronchiectasis on a plain film, and a completely normal radiograph is unlikely once the disease is established.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://packrat3.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE