PHYSICIAN ASSISTANT CLINICAL KNOWLEDGE RATING AND ASSESSMENT TOOL (PACKRAT) 4 PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://packrat4.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. A 26-year-old male presents with fatigue, night sweats, and a painless neck mass; what is the next best step in evaluation?

A. Bone marrow biopsy and aspiration

B. Lymph node biopsy

C. Chest CT scan

D. Reexamine in 2-4 weeks

2. A very anxious patient presents with dizziness, perioral and extremity paresthesias, and tachypnea. This presentation is most consistent with:

A. Bronchial Asthma

B. Hyperventilation Syndrome

C. Spontaneous Pneumothorax

D. Emphysema

3. Which diagnosis is most likely in a 16-day-old infant with very forceful vomiting after feeds, persistent hunger, and a small epigastric mass?

A. Achalasia

B. Tracheoesophageal Fistula

C. Pyloric Stenosis

D. Meckel's Diverticulum

4. In the acute evaluation of head injury, skull radiographs primarily detect which finding?

A. Skull fracture

B. Intracranial hemorrhage

C. Cerebral edema

D. Meningeal enhancement

5. The treatment of choice for syphilis is which of the following?

A. Benzathine penicillin G

B. Valacyclovir

C. Metronidazole

D. Ceftriaxone

SAMPLE

6. Which statement best describes the presentation of viral gastroenteritis in children?

A. Afebrile with loose, watery, non-bloody diarrhea

B. Febrile with bloody diarrhea

C. Chronic diarrhea with weight loss

D. Severe dehydration with mucous in stool

7. In a patient with dysuria, vulvovaginal pruritus, and a frothy discharge, which finding would be expected on a saline preparation?

A. Clue cells visible on saline preparation

B. Hyphae and budding yeast on KOH preparation

C. Intracellular Gram-negative diplococci on Gram stain

D. Motile, flagellated protozoa on saline preparation

8. Which laboratory abnormality is most commonly noted in bulimia nervosa?

A. Glycosuria

B. Hypokalemia

C. Metabolic Acidosis

D. Hyperalbuminemia

9. Postoperative oliguria with an increased BUN:creatinine ratio and a fractional excretion of sodium less than 1% is most consistent with which diagnosis?

A. Prerenal azotemia

B. Acute tubular necrosis

C. Acute glomerulonephritis

D. Obstructive uropathy

10. Which skin rash is pathognomonic of rheumatic fever?

A. Erythema marginatum

B. Erythema nodosum

C. Diffuse petechiae

D. Purulent bullae

SAMPLE

Answers

SAMPLE

1. B
2. B
3. C
4. A
5. A
6. A
7. D
8. B
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. A 26-year-old male presents with fatigue, night sweats, and a painless neck mass; what is the next best step in evaluation?

A. Bone marrow biopsy and aspiration

B. Lymph node biopsy

C. Chest CT scan

D. Reexamine in 2-4 weeks

This presentation raises strong suspicion for lymphoma in a young adult with a painless neck mass and systemic symptoms like fatigue and night sweats. The key step is to obtain a tissue diagnosis from the enlarged lymph node to confirm lymphoma and determine the subtype, which drives treatment. An excisional lymph node biopsy is the best option because it provides full nodal architecture, allowing pathologists to identify malignant cells such as Reed-Sternberg cells and to perform necessary immunophenotyping. This architectural detail is essential for distinguishing lymphoma from benign reactive hyperplasia or infection and for accurate subtyping, which guides therapy. Other tests have roles after diagnosis: imaging like chest CT helps stage disease and assess mediastinal involvement, and bone marrow biopsy can be indicated if there is suspicion of marrow involvement or certain subtypes, but they do not establish the initial diagnosis. Waiting or rechecking in a few weeks would delay needed evaluation and treatment.

2. A very anxious patient presents with dizziness, perioral and extremity paresthesias, and tachypnea. This presentation is most consistent with:

A. Bronchial Asthma

B. Hyperventilation Syndrome

C. Spontaneous Pneumothorax

D. Emphysema

The main idea is hyperventilation syndrome from anxiety. When someone breathes rapidly, they blow off CO2 faster than the body can produce it, leading to respiratory alkalosis. This causes dizziness from cerebral vasoconstriction and tingling around the mouth and in the extremities due to the rise in pH and the resulting shift in calcium binding (lower ionized calcium). Tachypnea fits with an anxious, hyperventilating patient, making this the most consistent explanation for the symptoms. Bronchial asthma and spontaneous pneumothorax would typically present with different signs—wheezing or unilateral chest pain with decreased breath sounds, respectively—rather than the characteristic paresthesias and dizziness from low CO2. Emphysema is a chronic condition with progressive dyspnea, not an acute anxious presentation with perioral tingling. In short, rapid, anxiety-driven breathing lowers CO2, produces respiratory alkalosis, causes paresthesias and dizziness, and fits the clinical picture best. Management focuses on calming the patient and re-establishing normal breathing patterns.

SAMPLE

3. Which diagnosis is most likely in a 16-day-old infant with very forceful vomiting after feeds, persistent hunger, and a small epigastric mass?

A. Achalasia

B. Tracheoesophageal Fistula

C. Pyloric Stenosis

D. Meckel's Diverticulum

This vignette highlights hypertrophic pyloric stenosis. When the pyloric muscle thickens, the gastric outlet becomes narrowed, causing nonbilious, forceful vomiting after feeds. The infant often stays hungry because the stomach empties quickly, triggering a cycle of feeding and vomiting. A small, firm epigastric mass—the thickened pylorus, sometimes described as an "olive"—is a classic finding in this condition, and presentation in the first weeks of life (around two to six weeks) fits a 16-day-old well. Other conditions don't align with this pattern: achalasia can cause trouble with swallowing and regurgitation but not a palpable pyloric mass in the epigastrium; tracheoesophageal fistula presents with coughing, choking, and drooling during feeds and respiratory distress very early in life; Meckel's diverticulum typically causes painless lower GI bleeding rather than projectile vomiting or a gastric outlet obstruction with a pyloric mass.

4. In the acute evaluation of head injury, skull radiographs primarily detect which finding?

A. Skull fracture

B. Intracranial hemorrhage

C. Cerebral edema

D. Meningeal enhancement

In acute head trauma, the main thing skull X rays are used to identify is bone injury. They're best at showing fracture lines, displacement, or other disruptions in the bony skull. They don't reliably reveal soft tissue injuries inside the brain, such as intracranial hemorrhage or edema, and they aren't used to assess meningeal conditions. Detecting blood within the brain or swelling is more accurately done with CT (or MRI with contrast for certain details). So the finding skull radiographs are most likely to reveal is a skull fracture.

5. The treatment of choice for syphilis is which of the following?

A. Benzathine penicillin G

B. Valacyclovir

C. Metronidazole

D. Ceftriaxone

Benzathine penicillin G is the go-to treatment for syphilis because it delivers a prolonged, treponemicidal level of penicillin in the bloodstream with a single injection for early disease (and a course of injections for later stages), ensuring sustained exposure of Treponema pallidum to the antibiotic. This long-acting formulation is specifically effective against the organism and has stood the test of time as the first-line therapy across stages of syphilis, except for neurosyphilis, which requires IV penicillin. The other drugs listed don't reliably treat syphilis—valacyclovir targets herpesviruses, metronidazole targets anaerobes and certain parasites, and ceftriaxone can be used as an alternative in some penicillin-allergic cases but is not the preferred first-line option.

SAMPLE

6. Which statement best describes the presentation of viral gastroenteritis in children?

A. Afebrile with loose, watery, non-bloody diarrhea

B. Febrile with bloody diarrhea

C. Chronic diarrhea with weight loss

D. Severe dehydration with mucous in stool

Viral gastroenteritis in children usually presents with a sudden onset of vomiting followed by loose, watery stools that are not bloody, and children are often afebrile or have only a low fever. The absence of blood in the stool helps distinguish it from bacterial causes, which more often bring fever and hematochezia. Chronic diarrhea with weight loss points to other conditions like malabsorption or inflammatory diseases, not an acute viral illness. Dehydration can occur with any gastroenteritis, but mucus in stool is not a defining feature of a viral infection. So the pattern of afebrile, loose, watery, non-bloody diarrhea aligns best with viral gastroenteritis in kids.

7. In a patient with dysuria, vulvovaginal pruritus, and a frothy discharge, which finding would be expected on a saline preparation?

A. Clue cells visible on saline preparation

B. Hyphae and budding yeast on KOH preparation

C. Intracellular Gram-negative diplococci on Gram stain

D. Motile, flagellated protozoa on saline preparation

Dysuria with vulvovaginal pruritus and a frothy discharge points to infection with Trichomonas vaginalis, a flagellated protozoan. On a saline saline wet mount, the hallmark finding is motile, pear-shaped trophozoites that have flagella and move actively. This motility is what sets trichomonas apart from other vaginal infections. Clue cells seen on saline indicate bacterial vaginosis, not a motile protozoan. Hyphae and budding yeast on a KOH prep point to vulvovaginal candidiasis. Intracellular Gram-negative diplococci on Gram stain would suggest gonorrheal infection. The presence of motile, flagellated protozoa on saline directly corresponds to Trichomonas infection, which matches the clinical picture.

8. Which laboratory abnormality is most commonly noted in bulimia nervosa?

A. Glycosuria

B. Hypokalemia

C. Metabolic Acidosis

D. Hyperalbuminemia

Frequent purging in bulimia causes loss of potassium along with gastric hydrogen ions. Vomiting and laxative use deplete potassium stores, and the accompanying volume depletion stimulates aldosterone, promoting further potassium loss in the kidneys. The net result is a low serum potassium level, making hypokalemia the most common laboratory finding in bulimia. Glycosuria isn't a typical finding unless another condition like diabetes is present. Metabolic acidosis would be unlikely here because vomiting leads to loss of HCl and typically causes metabolic alkalosis, not acidosis. Hyperalbuminemia isn't expected from bulimia-related electrolyte disturbances.

SAMPLE

9. Postoperative oliguria with an increased BUN:creatinine ratio and a fractional excretion of sodium less than 1% is most consistent with which diagnosis?

A. Prerenal azotemia

B. Acute tubular necrosis

C. Acute glomerulonephritis

D. Obstructive uropathy

When the kidney is underperfused but the tubules are still intact, it conserves water and sodium to preserve circulating volume. This produces an elevated BUN relative to creatinine because urea is reabsorbed more avidly in the proximal tubule during low flow, while creatinine is not. The kidneys also actively reabsorb sodium, so the fractional excretion of sodium stays very low, typically less than 1%. In the postoperative setting, this pattern points to reduced renal perfusion or volume depletion causing prerenal azotemia. This differs from intrinsic kidney injury, where the tubules are damaged and cannot reclaim sodium effectively, leading to a higher fractional excretion of sodium (often >2%). Glomerular disease would present with different urine findings such as proteinuria or RBC casts, and obstructive (postrenal) causes usually involve signs of obstruction or impaired urine flow with variable sodium handling as the condition evolves.

10. Which skin rash is pathognomonic of rheumatic fever?

A. Erythema marginatum

B. Erythema nodosum

C. Diffuse petechiae

D. Purulent bullae

Erythema marginatum is the rash most closely associated with rheumatic fever. It appears as a ring-shaped, nonpruritic eruption with a pale center, often on the trunk and proximal limbs, and it tends to come and go. Its appearance is highly specific for rheumatic fever and is one of the major Jones criteria used to support the diagnosis. Other skin findings listed—erythema nodosum (tender nodules on the shins), diffuse petechiae (tiny red spots from bleeding), and purulent bullae (blistering infections)—do not have the same diagnostic specificity for rheumatic fever, so they don't serve as the definitive clue that erythema marginatum provides.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://packrat4.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE