POSTERIOR SEGMENT BLOCK 15 – AMD AND OTHER CAUSES OF CNV PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://posteriorsegmentblock15.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1
Table of Contents 2
Introduction 3
How to Use This Guide 4
Questions 5
Answers 8
Explanations 10
Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What type of visual symptoms may result from idiopathic choroidal neovascularization?

A. Metamorphopsia or blurry vision

B. Night blindness

C. Peripheral vision loss

D. Sudden complete vision loss

2. Which factor is a significant inhibitor of angiogenesis?

A. VEGF

B. Pigment epithelium derived factor (PEDF)

C. Angiopoietin-1

D. TGF-α

3. What characterizes the AREDS Advanced AMD grading stage?

A. Geographic atrophy not involving fovea

B. Presence of geographic atrophy involving the foveal center

C. Only small drusen present

D. No signs of AMD

4. What structure does the inner collagenous zone of Bruch's membrane separate?

A. RPE plasma membrane from the retinal pigment epithelium

B. Basal laminar deposits from the RPE basement membrane

C. RPE basement membrane from the choroidal vessels

D. Choroidal stroma from Bruch's outer collagenous zone

5. Which type of CNV is characterized as occurring in the subretinal space?

A. Type 1

B. Type 2

C. Type 3

D. Type 4

SAMPLE

6. High doses of zinc in AREDS1 are potentially associated with which kind of problems?

A. Cardiovascular problems

B. Neurological problems

C. Genitourinary tract problems

D. Respiratory problems

7. Choroidal neovascularization (CNV) extends through which layer from the choriocapillaris?

A. Retinal nerve fiber layer

B. Bruch's membrane

C. Outer nuclear layer

D. Photoreceptor layer

8. What is the recommended dosing regimen for Eylea?

A. One injection every month

B. One injection every 2 months

C. One injection every 6 weeks

D. One injection every 3 months

9. What demographic is most likely to experience toxic retinopathies as a differential of dry non-neovascular AMD?

A. Individuals with diabetes

B. Young males

C. Older women

D. Individuals with a history of substance use

10. Which type of diet intake is associated with a higher risk of advanced AMD?

A. High Carbohydrate

B. High Protein

C. Fatty

D. Low-Calorie

SAMPLE

Answers

SAMPLE

1. A
2. B
3. B
4. C
5. B
6. C
7. B
8. B
9. D
10. C

SAMPLE

Explanations

SAMPLE

1. What type of visual symptoms may result from idiopathic choroidal neovascularization?

A. Metamorphopsia or blurry vision

B. Night blindness

C. Peripheral vision loss

D. Sudden complete vision loss

A. Metamorphopsia or blurry vision is the correct response because idiopathic choroidal neovascularization (CNV) often affects the macula, which is the central part of the retina responsible for sharp, detailed vision. When abnormal blood vessels grow in the choroid layer of the eye and leak fluid or blood, this can distort visual perception. Patients commonly report symptoms like metamorphopsia, where straight lines appear wavy or distorted, and blurry vision due to the disruption in the normal retinal architecture. These visual changes are often among the first indicators of CNV, making them characteristic symptoms of this condition. Other visual symptoms listed, while they can occur in various ocular diseases, are less specifically associated with idiopathic CNV. Night blindness typically relates more to rod photoreceptor function and is not a primary symptom of CNV. Peripheral vision loss might suggest other retinal or optic nerve issues rather than localized changes from CNV. Sudden complete vision loss is more indicative of retinal detachment or severe retinal ischemia rather than the gradual deterioration often seen with choroidal neovascularization.

2. Which factor is a significant inhibitor of angiogenesis?

A. VEGF

B. Pigment epithelium derived factor (PEDF)

C. Angiopoietin-1

D. TGF-α

Pigment epithelium-derived factor (PEDF) plays a crucial role as a significant inhibitor of angiogenesis. It is a neurotrophic factor that is primarily produced by retinal pigment epithelial cells and has been shown to possess strong anti-angiogenic properties. PEDF acts by counteracting the effects of pro-angiogenic factors such as vascular endothelial growth factor (VEGF), thereby helping to regulate and maintain normal blood vessel formation. In contexts where inappropriate angiogenesis contributes to conditions like age-related macular degeneration (AMD), the presence of PEDF can help prevent the abnormal growth of blood vessels which is often associated with the progression of these diseases. This makes PEDF a vital component in ocular health and a potential therapeutic target for diseases driven by neovascularization. The other factors mentioned, such as VEGF, angiopoietin-1, and TGF-α, are known for promoting angiogenesis in various physiological and pathological processes, highlighting the distinct role of PEDF as an inhibitor in this context.

SAMPLE

3. What characterizes the AREDS Advanced AMD grading stage?

A. Geographic atrophy not involving fovea

B. Presence of geographic atrophy involving the foveal center

C. Only small drusen present

D. No signs of AMD

The AREDS Advanced AMD grading stage is characterized by the presence of geographic atrophy that involves the foveal center. Geographic atrophy represents a significant form of macular degeneration that results in the loss of retinal pigment epithelium and photoreceptors, leading to visual impairment. When this atrophy encroaches on the fovea, the central part of the retina responsible for sharp, detailed vision, it indicates a more advanced stage of age-related macular degeneration (AMD). This condition can severely affect a person's central vision, making it critical to identify and understand this stage in the context of AMD management and treatment strategies. In contrast, the other options either describe earlier stages of AMD or do not indicate any disease presence at all. Therefore, understanding the distinction of geographic atrophy involving the fovea is crucial for proper grading and subsequent clinical considerations.

4. What structure does the inner collagenous zone of Bruch's membrane separate?

A. RPE plasma membrane from the retinal pigment epithelium

B. Basal laminar deposits from the RPE basement membrane

C. RPE basement membrane from the choroidal vessels

D. Choroidal stroma from Bruch's outer collagenous zone

The inner collagenous zone of Bruch's membrane serves a crucial role in the structure of the eye, particularly in separating the retinal pigment epithelium (RPE) basement membrane from the choroidal vessels. This separation is significant because it helps maintain the integrity of the blood-retinal barrier and facilitates the transport of nutrients and waste products between the choroidal circulation and the retina. Bruch's membrane itself is a thin barrier that lies between the RPE and the choroid, and its inner collagenous zone specifically acts as a supportive structure that anchors the RPE to the deeper vascular layer, ensuring proper metabolic function. This functional alignment is vital for the health of the retina and impacts conditions such as age-related macular degeneration (AMD) when disruptions occur. The context around the other options highlights why they do not accurately represent the role of the inner collagenous zone of Bruch's membrane. The first choice erroneously implies a separation that does not involve the deeper layers of Bruch's membrane. The second option incorrectly states a relationship involving basal laminar deposits, which are associated but not separated by this particular zone. The fourth choice misrepresents the layer's position and its relevant structures, which do not correlate with the role

SAMPLE

5. Which type of CNV is characterized as occurring in the subretinal space?

A. Type 1

B. Type 2

C. Type 3

D. Type 4

Type 2 choroidal neovascularization (CNV) is specifically characterized by the presence of new blood vessels that develop in the subretinal space. This type of CNV is often associated with conditions such as age-related macular degeneration (AMD), where the new vessels breach the retinal pigment epithelium (RPE) and invade the neuroretinal layers. Understanding the subretinal location is critical, as it influences the clinical presentation and potential treatment strategies for the condition. In contrast, other types of CNV, such as Type 1 and Type 3, are associated with different anatomical relationships concerning the retinal layers and the RPE, which can affect their management and outcomes. Hence, identifying Type 2 as the variant occurring specifically in the subretinal space is essential for clinical diagnostics and therapeutic interventions.

6. High doses of zinc in AREDS1 are potentially associated with which kind of problems?

A. Cardiovascular problems

B. Neurological problems

C. Genitourinary tract problems

D. Respiratory problems

High doses of zinc, as studied in the Age-Related Eye Disease Study (AREDS1), are primarily associated with genitourinary tract problems, specifically urinary issues. Patients taking high doses of zinc may experience an increased risk of developing urinary tract infections and other related concerns. Zinc is an essential mineral, and while it plays a crucial role in various bodily functions, excessive intake can lead to side effects. The AREDS1 study aimed to assess the effectiveness of antioxidant vitamins and minerals in slowing the progression of age-related macular degeneration (AMD), but it also highlighted potential risks associated with high doses, including those related to the urinary system. Understanding the implications of high zinc intake is important for clinical practice, as healthcare providers need to monitor patients for adverse effects when recommending supplements based on the findings from such studies.

SAMPLE

7. Choroidal neovascularization (CNV) extends through which layer from the choriocapillaris?

A. Retinal nerve fiber layer

B. Bruch's membrane

C. Outer nuclear layer

D. Photoreceptor layer

The correct answer highlights the pathway and structural significance of choroidal neovascularization (CNV) as it extends from the choriocapillaris, which is primarily composed of small blood vessels. When CNV begins, it typically breaches the Bruch's membrane, a critical barrier situated between the choroidal circulation and the retinal layers. Bruch's membrane serves as a supportive matrix and plays a crucial role in the transportation of nutrients and waste between the retina and the underlying choroidal blood supply. When pathological conditions, such as age-related macular degeneration (AMD), lead to the formation of CNV, abnormal blood vessels can proliferate and invade through this membrane. This invasion is pivotal because it can disrupt retinal function and contribute to vision loss. Understanding this process is essential for recognizing the implications of CNV on vision, as its progression can lead to hemorrhages, fluid accumulation, and scarring that impair the retina. The other layers mentioned, like the retinal nerve fiber layer, outer nuclear layer, and photoreceptor layer, are involved in visual processing but do not play the same role in the initial pathway of CNV formation from the choriocapillaris. Thus, identifying Bruch's membrane as the

8. What is the recommended dosing regimen for Eylea?

A. One injection every month

B. One injection every 2 months

C. One injection every 6 weeks

D. One injection every 3 months

The recommended dosing regimen for Eylea (aflibercept) in the treatment of conditions like age-related macular degeneration (AMD) typically begins with a loading phase that involves one injection per month for the first three months, after which the interval can generally be extended. Following this initial loading phase, the common practice is to administer one injection every two months as a maintenance dose. This regimen is based on clinical studies showing that extending the interval to every two months allows for effective management of the condition while also providing flexibility for the patient's treatment plan. The transition from monthly to bi-monthly dosing enhances adherence and helps to reduce treatment burden without compromising efficacy. The knowledge of this regimen is essential for healthcare professionals to provide appropriate patient care in managing AMD and related conditions effectively.

SAMPLE

9. What demographic is most likely to experience toxic retinopathies as a differential of dry non-neovascular AMD?

A. Individuals with diabetes

B. Young males

C. Older women

D. Individuals with a history of substance use

Toxic retinopathies are conditions that can lead to retinal damage due to exposure to certain substances, including some medications and recreational drugs. Individuals with a history of substance use are particularly at risk because they may have been exposed to harmful agents that can adversely affect the retina. Substances like hydroxychloroquine and certain recreational drugs can lead to visual disturbances and retinal toxicity, manifesting in a way that can be confused with dry non-neovascular age-related macular degeneration (AMD). While older women and individuals with diabetes are common demographics for various retinal issues, including AMD, these populations are less associated with toxic retinopathies specifically. Young males, while they can be affected by such conditions, do not represent the broader demographic trend seen with toxic retinopathies. Thus, individuals with a history of substance use stand out as the group most likely to experience these toxic effects that can mimic or complicate the presentation of dry AMD.

10. Which type of diet intake is associated with a higher risk of advanced AMD?

A. High Carbohydrate

B. High Protein

C. Fatty

D. Low-Calorie

A diet high in fatty foods is associated with a higher risk of advanced age-related macular degeneration (AMD) due to its impact on overall vascular health and inflammation. Diets rich in saturated fats can contribute to systemic inflammation, which is thought to play a role in the progression of AMD. Additionally, certain types of fats, particularly those found in processed and fried foods, may promote oxidative stress and damage to retinal tissues over time. In contrast, other types of diets, such as those high in carbohydrates or proteins, do not share the same strong association with advanced AMD. Low-calorie diets, when properly balanced and nutritious, may even be protective against various age-related conditions, including AMD, by reducing overall inflammation and promoting better metabolic health. Consequently, a diet high in unhealthy fats is more closely linked to the increased risk of AMD, making this the correct choice.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://posteriorsegmentblock15.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE