
PREHOSPITAL TRAUMA LIFE SUPPORT (PHTLS) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://prehospitaltraumalifesupport.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **Deterioration of ventilation after inflation of a PASG following high-pressure abdominal compression suggests which injury?**

   A. Abdominal aortic aneurysm

   B. Ruptured diaphragm

   C. Ruptured esophagus

   D. "Paper bag" syndrome of the lungs

2. **What considerations should be made in handling a pregnant trauma patient?**

   A. Only focus on maternal care, ignoring fetal viability

   B. Assess for fetal viability and provide appropriate care

   C. Transport immediately without assessment

   D. Only check for maternal injuries without considering pregnancy

3. **What is the main priority in the initial assessment of a trauma patient?**

   A. Ensuring airway patency

   B. Assessing for any external bleeding

   C. Performing a detailed neurological examination

   D. Administering pain relief medications

4. **In the event of a traumatic brain injury, which sign is most indicative of increased intracranial pressure?**

   A. Confusion or disorientation

   B. Unequal pupil size

   C. Persistent vomiting

   D. Lateralizing motor weakness

5. **Why is it vital to keep trauma patients warm?**

   A. To prevent hypothermia, which can worsen coagulopathy and shock

   B. To maintain normal heart rates in recovery

   C. To promote faster wound healing

   D. To ensure comfort during transportation

SAMPLE

6. When assessing a trauma patient, what is the importance of noting a history of allergies?

A. To ensure proper medication management during treatment

B. It is not necessary

C. To assess the patient's general health

D. To determine past traumatic experiences

7. What role does shock play in trauma care?

A. It provides a clear understanding of the patient's pain level

B. It indicates the body's response to severe injury

C. It shows the effectiveness of airway management

D. It is a sign that medication is required immediately

8. What should be suspected in a trauma patient presenting with vomiting blood?

A. Possible abdominal injuries or upper gastrointestinal bleeding

B. Chronic liver disease

C. Internal bleeding from the lungs

D. Esophageal varices

9. What is the most appropriate initial action for a deep laceration with significant bleeding?

A. Apply a tourniquet

B. Apply direct pressure

C. Initiate rapid transport

D. Restore blood volume

10. In trauma care, what is a critical indicator of a patient's need for immediate attention?

A. Presence of pain in the chest area

B. Uncontrolled external bleeding

C. Minor abrasions on the skin

D. Patient's age

SAMPLE

Answers

SAMPLE

1. B
2. B
3. A
4. B
5. A
6. A
7. B
8. A
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. Deterioration of ventilation after inflation of a PASG following high-pressure abdominal compression suggests which injury?

A. Abdominal aortic aneurysm

B. Ruptured diaphragm

C. Ruptured esophagus

D. "Paper bag" syndrome of the lungs

The correct answer is linked to the concept of thoracic and abdominal compartment syndrome, particularly in the case of a ruptured diaphragm. When high pressures are applied to the abdomen through the use of a Pneumatic Antishock Garment (PASG), it can severely affect diaphragm function, especially if there is an existing injury to the diaphragm. A ruptured diaphragm would allow abdominal contents, including the stomach and intestines, to move into the thoracic cavity, which would compromise lung function and ventilation. This shift can lead to reduced chest expansion and ventilatory effort, manifesting as deterioration in ventilation after the application of the PASG. This situation highlights how mechanical effects from abdominal pressure can exacerbate an already compromised respiratory system due to the injury. In contrast, other injuries like an abdominal aortic aneurysm, a ruptured esophagus, or "paper bag" syndrome don't directly lead to a significant deterioration in ventilation specifically caused by inflation of a PASG. An abdominal aortic aneurysm primarily presents with vascular symptoms; a ruptured esophagus may lead to mediastinal complications but not directly from high abdominal pressure; and "paper bag" syndrome refers to lung injury from asphyxiation but is not a direct result of abdominal compression impacts.

2. What considerations should be made in handling a pregnant trauma patient?

A. Only focus on maternal care, ignoring fetal viability

B. Assess for fetal viability and provide appropriate care

C. Transport immediately without assessment

D. Only check for maternal injuries without considering pregnancy

When handling a pregnant trauma patient, it is crucial to assess for fetal viability and provide appropriate care. This is because the well-being of both the mother and the fetus is intertwined; trauma can have significant implications for both. Evaluating the fetal condition includes monitoring for signs of life, such as fetal heart tones, and understanding how maternal injuries may affect the fetus, especially in later stages of pregnancy. Proper assessment leads to better-informed decision-making regarding treatment and transport options. Providing care that considers the needs of both the mother and the fetus is essential for optimizing outcomes for both. This holistic approach acknowledges the potential risks trauma poses to the fetus and emphasizes the importance of maternal stabilization, which can directly impact fetal health. In contrast, focusing solely on maternal care while ignoring the fetus would risk neglecting vital aspects of care that could save the life of the baby. Transporting immediately without assessment would bypass critical evaluations that could identify injuries or complications affecting both the mother and fetus. Lastly, only checking for maternal injuries without taking the pregnancy into account could lead to inadequate care, as certain findings may suggest the need for an urgent intervention to protect both lives.

SAMPLE

3. What is the main priority in the initial assessment of a trauma patient?

A. Ensuring airway patency

B. Assessing for any external bleeding

C. Performing a detailed neurological examination

D. Administering pain relief medications

The main priority in the initial assessment of a trauma patient is ensuring airway patency. This is critical because a compromised airway can lead to inadequate oxygenation and, ultimately, can be life-threatening if not immediately addressed. In trauma situations, patients may have obstacles such as blood, vomit, or facial injuries that can obstruct the airway. Therefore, securing the airway is vital to allow for effective ventilation and oxygen delivery, which are essential for survival. While assessing for external bleeding and performing a detailed neurological examination are important components of the overall trauma assessment, they come after addressing the airway in the hierarchy of priorities. Ensuring that the patient can breathe effectively is the first step in the assessment process. Administering pain relief is also important, but it should not take precedence over securing the airway or ensuring that the patient can oxygenate properly.

4. In the event of a traumatic brain injury, which sign is most indicative of increased intracranial pressure?

A. Confusion or disorientation

B. Unequal pupil size

C. Persistent vomiting

D. Lateralizing motor weakness

Unequal pupil size is a significant indicator of increased intracranial pressure (ICP) following a traumatic brain injury. This condition, known as anisocoria, suggests that there may be pressure on the optic nerve or other structures within the cranial cavity. When ICP rises, it can lead to brain swelling or hemorrhage, impacting the surrounding nerves, including those that control pupil reaction and size. Changes in pupil size reflect underlying issues within the brain. For instance, if one pupil is dilated (mydriasis) and the other is constricted (miosis), this can indicate a hematoma, brain herniation, or other forms of injury that influence ICP. Monitoring these signs is crucial in the management of traumatic brain injuries, as they can lead to rapid deterioration if not addressed promptly. While confusion or disorientation, persistent vomiting, and lateralizing motor weakness can all suggest brain injury or other neurological impairments, they are not as specific to increased ICP as unequal pupil size. Unequal pupils are a more direct reflection of pressure dynamics within the cranial cavity, making them a critical component in initial assessments and interventions for patients with head trauma.

SAMPLE

5. Why is it vital to keep trauma patients warm?

A. To prevent hypothermia, which can worsen coagulopathy and shock

B. To maintain normal heart rates in recovery

C. To promote faster wound healing

D. To ensure comfort during transportation

Keeping trauma patients warm is essential primarily to prevent hypothermia, which can significantly exacerbate conditions such as coagulopathy and shock. When a patient experiences trauma, their body is already under stress, and the introduction of hypothermia can lead to a vicious cycle. Hypothermia can impair platelet function and clotting factors, worsening coagulopathy, which is the inability of blood to clot properly. This increased bleeding can compound the challenges involved in managing shock, as it not only reduces blood volume but also affects the body's ability to compensate for the loss. Maintaining normothermia is thus crucial in a trauma setting to support hemostatic functions and improve overall outcomes. While maintaining normal heart rates, promoting wound healing, and ensuring comfort can also be important aspects of patient care, they are secondary to the more immediate need to prevent life-threatening complications that arise from hypothermia. Ensuring that trauma patients are kept warm directly addresses a critical factor in their stabilization and recovery.

6. When assessing a trauma patient, what is the importance of noting a history of allergies?

A. To ensure proper medication management during treatment

B. It is not necessary

C. To assess the patient's general health

D. To determine past traumatic experiences

Noting a history of allergies is critical in trauma patient assessment primarily for ensuring proper medication management during treatment. Patients with known allergies may react adversely to certain medications or treatments that are otherwise standard practice. For instance, if a patient has a documented allergy to a particular antibiotic, administering that drug could lead to severe allergic reactions such as anaphylaxis, which not only complicates the patient's condition but can also endanger their life. In trauma situations, time is often of the essence, and knowing a patient's allergy history allows responders to make informed, rapid decisions about medication administration while avoiding potentially harmful substances. Additionally, this knowledge helps in determining alternative options that are safe for the patient. The other options do not align with the practical implications of allergy history during acute medical situations. General health assessments may provide insight into the patient's overall well-being, but they do not address immediate treatment needs. Similarly, while past traumatic experiences may be relevant in certain contexts, they typically do not influence the urgent management of medical care in trauma scenarios. Noting allergies directly affects patient safety and treatment efficacy, making it a primary concern in any critical care situation.

SAMPLE

7. What role does shock play in trauma care?

A. It provides a clear understanding of the patient's pain level

B. It indicates the body's response to severe injury

C. It shows the effectiveness of airway management

D. It is a sign that medication is required immediately

Shock plays a crucial role in trauma care as it serves as an indication of the body's response to severe injury. When a patient experiences trauma, the body can react with a state of shock, which is characterized by inadequate blood flow and oxygen delivery to vital organs. This physiological response occurs due to various factors, such as blood loss, fluid shifts, or severe pain, and is often a sign of underlying issues like hypovolemia or significant internal injuries. Recognizing the signs of shock is essential in trauma care, as it allows healthcare providers to prioritize treatment and implement interventions quickly to stabilize the patient. Immediate actions may involve administering fluids, controlling any active bleeding, or addressing airway concerns to restore adequate perfusion and protect organ function. While other aspects of trauma care, such as pain management and airway effectiveness, are important, they do not directly reflect the broader systemic response to injury that shock encompasses. Therefore, understanding and recognizing shock as a response to trauma is vital for determining the urgency and nature of treatment in critically injured patients.

8. What should be suspected in a trauma patient presenting with vomiting blood?

A. Possible abdominal injuries or upper gastrointestinal bleeding

B. Chronic liver disease

C. Internal bleeding from the lungs

D. Esophageal varices

In a trauma patient who presents with vomiting blood, the primary suspicion should be for possible abdominal injuries or upper gastrointestinal bleeding. Trauma can lead to various types of injuries that affect the abdominal organs, such as the liver, spleen, and stomach, which are susceptible to lacerations or contusions that can result in internal bleeding. Additionally, upper gastrointestinal bleeding can stem from several causes, including gastric or duodenal ulcers or tears in the lining of the esophagus, particularly after significant force or injury. Although chronic liver disease, internal bleeding from the lungs, and esophageal varices could also lead to hematemesis (vomiting blood), these conditions are typically not the immediate concern in the context of an acute trauma scenario. Trauma mechanisms often directly impact abdominal structures, making abdominal injuries a likely source of bleeding. Prompt evaluation is critical as it allows for appropriate management and intervention to address potentially life-threatening conditions.

SAMPLE

9. What is the most appropriate initial action for a deep laceration with significant bleeding?

A. Apply a tourniquet

B. Apply direct pressure

C. Initiate rapid transport

D. Restore blood volume

Applying direct pressure to a deep laceration with significant bleeding is the most appropriate initial action. This method addresses the primary goal of controlling bleeding effectively. Direct pressure helps to compress the blood vessels at the site of the injury, which can significantly reduce blood loss. It is a fundamental technique taught in trauma care, emphasizing that controlling hemorrhage is a crucial step before considering further interventions. This approach should be applied immediately while assessing the patient's overall condition, which may include checking responsiveness, airway, breathing, and circulation. If direct pressure is not successful in controlling the bleeding within a reasonable time, other measures like a tourniquet can be considered, especially for limbs. However, initiating rapid transport or restoring blood volume can be critical in managing trauma patients, those actions are typically secondary to and contingent upon the effective control of bleeding at the source. Thus, starting with direct pressure is a foundational principle in trauma management.

10. In trauma care, what is a critical indicator of a patient's need for immediate attention?

A. Presence of pain in the chest area

B. Uncontrolled external bleeding

C. Minor abrasions on the skin

D. Patient's age

Uncontrolled external bleeding is a critical indicator of a patient's need for immediate attention because it represents a medical emergency that can lead to rapid deterioration and potentially life-threatening consequences. When a patient experiences uncontrolled bleeding, the body loses blood volume, which can result in shock. Shock impairs the delivery of oxygen to vital organs, increasing the risk of organ failure and death if not addressed promptly. In trauma care, the priority is to stabilize the patient and prevent blood loss. Immediate intervention, such as direct pressure on the wound, elevation of the injured area, or the use of a tourniquet if applicable, is crucial to manage uncontrolled bleeding. This immediate response can significantly improve the patient's chances of survival and recovery. Other indicators, such as the presence of pain in the chest area or a patient's age, may also signify a need for attention, but they do not convey the same level of urgency as uncontrolled external bleeding. Minor abrasions on the skin typically do not require immediate intervention compared to the serious implications of uncontrolled bleeding.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://prehospitaltraumalifesupport.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE