PROPHECY EMERGENCY DEPARTMENT REGISTERED NURSE (RN) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://prophecyemergencydeptrn.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What is the immediate response for a patient who has ingested a toxic substance?**

   A. Call for emergency assistance

   B. Assess airway and breathing, then call Poison Control

   C. Administer activated charcoal

   D. Induce vomiting immediately

2. **In a scenario involving a 79-year-old male with severe diarrhea, what nursing action is essential during patient education?**

   A. Instruct on dietary modifications

   B. Emphasize the importance of hydration

   C. Discuss medication options

   D. Advise on hygiene practices

3. **What is the first line of treatment for a patient with multiple sclerosis presenting with weakened lower extremities?**

   A. Intravenous fluids

   B. Methylprednisolone 1 g IV daily

   C. Steroid injections

   D. Physical therapy

4. **What is one of the first actions a nurse should take in an emergency situation?**

   A. Document the incident for records

   B. Call for back-up immediately

   C. Assess the environment for hazards

   D. Begin immediate treatment

5. **In a pediatric emergency, what is the first step in a primary assessment?**

   A. Assessing the circulation

   B. Assessing the airway

   C. Assessing breathing

   D. Assessing the patient's history

SAMPLE

6. What is the most effective way to assess circulation in an unconscious adult patient?

A. Checking the radial pulse

B. Checking for a pulse at the carotid artery

C. Assessing capillary refill

D. Checking for a pulse at the femoral artery

7. A patient with pneumonia has a penicillin allergy. Which medication would you question if ordered?

A. Ceftriaxone

B. Azithromycin

C. Clindamycin

D. Levofloxacin

8. What does a "flat" ECG line indicate?

A. Normal sinus rhythm

B. Bradycardia

C. Asystole or absence of electrical activity in the heart

D. Atrial fibrillation

9. An elderly woman shows fatigue and shortness of breath similar to previous urinary tract infection symptoms. What diagnostic test is crucial to obtain during triage?

A. X-ray of the chest

B. Complete blood count

C. ECG

D. Urinalysis

10. What is the first-line treatment for a patient experiencing a myocardial infarction?

A. Administration of oxygen

B. Administration of aspirin

C. Insertion of a stent

D. Placement on telemetry monitoring

SAMPLE

Answers

SAMPLE

1. B
2. D
3. B
4. C
5. B
6. B
7. A
8. C
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. What is the immediate response for a patient who has ingested a toxic substance?

A. Call for emergency assistance

B. Assess airway and breathing, then call Poison Control

C. Administer activated charcoal

D. Induce vomiting immediately

The immediate response for a patient who has ingested a toxic substance involves assessing the patient's airway and breathing first to ensure that they are stable. This is a critical step because airway compromise or respiratory distress can lead to life-threatening situations. Once the patient's airway and breathing have been evaluated, it is appropriate to contact Poison Control. This organization can provide specialized guidance based on the specific toxic substance involved, including recommendations for further treatment. While getting emergency assistance is certainly important in a potential poisoning case, the priority should be on ensuring the patient's safety first through assessment of their airway and breathing. Administering activated charcoal may be indicated in some poisonings, but it should not be done until a healthcare professional advises it, as it is not appropriate for all toxic substances and may not be effective or safe if the patient is not alert or if certain agents were ingested. Inducing vomiting is generally no longer recommended in many cases of poisoning, as it can pose risks such as aspiration. Therefore, the most effective immediate response prioritizes patient stabilization and seeking expert guidance.

2. In a scenario involving a 79-year-old male with severe diarrhea, what nursing action is essential during patient education?

A. Instruct on dietary modifications

B. Emphasize the importance of hydration

C. Discuss medication options

D. Advise on hygiene practices

In the context of patient education for a 79-year-old male with severe diarrhea, emphasizing the importance of hydration is crucial. Severe diarrhea can lead to significant fluid and electrolyte loss, particularly in older adults who may already be at higher risk for dehydration. Therefore, educating the patient on the need to maintain adequate fluid intake helps to prevent dehydration and its associated complications, such as kidney injury or electrolyte imbalances. While discussing hygiene practices, dietary modifications, and medication options are also important components of care in managing diarrhea, they do not take precedence over the immediate need for hydration. Ensuring that the patient understands how to rehydrate and the signs of dehydration to watch for is essential for their safety and recovery. This educational focus on hydration empowers the patient to take charge of their health while minimizing the risks associated with severe diarrhea.

SAMPLE

3. What is the first line of treatment for a patient with multiple sclerosis presenting with weakened lower extremities?

A. Intravenous fluids

B. Methylprednisolone 1 g IV daily

C. Steroid injections

D. Physical therapy

The first line of treatment for a patient with multiple sclerosis (MS) who presents with weakened lower extremities is methylprednisolone 1 g intravenously daily. This corticosteroid treatment is specifically used to reduce inflammation and manage acute exacerbations of multiple sclerosis. Corticosteroids work by suppressing the immune system's response, which is crucial since MS is characterized by episodes of neurologic dysfunction due to demyelination. The use of high-dose steroids helps to shorten the duration of exacerbations and promote recovery of function. Other treatments such as intravenous fluids do not address the underlying inflammation and neurological symptoms of MS. Steroid injections can be used but are generally not as effective as high-dose intravenous methylprednisolone for acute exacerbations. While physical therapy is essential for rehabilitation and improving mobility in MS, it is not the first line of treatment during an acute episode characterized by significant weakness. Consequently, the administration of high-dose intravenous steroids is the most appropriate initial approach in such scenarios.

4. What is one of the first actions a nurse should take in an emergency situation?

A. Document the incident for records

B. Call for back-up immediately

C. Assess the environment for hazards

D. Begin immediate treatment

In an emergency situation, one of the first critical actions a nurse should take is to assess the environment for hazards. This is essential to ensure both the safety of the patient and the healthcare team. By evaluating the immediate environment, the nurse can identify potential dangers such as spills, electrical hazards, or the presence of aggressive individuals that could compromise the care being provided or the safety of everyone involved. This assessment allows the nurse to create a safe space for delivering care, preventing accidents or injuries that could further complicate the situation. Once the environment is deemed safe, the nurse can then move forward with appropriate interventions, call for additional personnel if needed, or document the incident accurately after managing the emergency. Prioritizing safety through environmental assessment lays the groundwork for effective and efficient emergency care.

SAMPLE

5. In a pediatric emergency, what is the first step in a primary assessment?

A. Assessing the circulation

B. Assessing the airway

C. Assessing breathing

D. Assessing the patient's history

The first step in a primary assessment, particularly in a pediatric emergency, is to assess the airway. This step is critical because the airway must be clear for effective breathing to occur. In children, the anatomy and physiology differ from adults, making them more susceptible to airway obstruction. By quickly evaluating whether the airway is patent, the nurse can determine if immediate interventions are needed, such as repositioning the head, suctioning, or administering supplemental oxygen. Ensuring that the airway is clear is the foundation for further assessment and treatment, as any compromise in this area can significantly impact the pediatric patient's ability to breathe and oxygenate their body effectively. Following the assessment of the airway, the nurse would typically assess breathing and circulation to continue the primary survey effectively, ensuring a comprehensive evaluation of the patient's condition.

6. What is the most effective way to assess circulation in an unconscious adult patient?

A. Checking the radial pulse

B. Checking for a pulse at the carotid artery

C. Assessing capillary refill

D. Checking for a pulse at the femoral artery

The most effective way to assess circulation in an unconscious adult patient is by checking for a pulse at the carotid artery. The carotid artery is centrally located in the neck and is a major vessel that provides blood to the brain. In an unconscious patient, it is crucial to quickly ascertain whether the heart is still beating effectively to ensure that vital organs, especially the brain, are receiving adequate blood flow and oxygen. Using the carotid pulse allows the healthcare provider to efficiently determine circulatory status without having to rely on potential muscle tone or other factors that might obscure peripheral circulation, which could be less reliable in an unconscious individual. This is particularly critical in emergency situations where time is of the essence, and immediate interventions may be necessary. In comparison, checking the radial pulse or femoral artery, while also valid options, may not provide as rapid an assessment of brain perfusion. Capillary refill could be misleading in unconscious patients who may not have normal peripheral perfusion regardless of their cardiac status. Therefore, assessing the carotid pulse is the preferred method in this context.

SAMPLE

7. A patient with pneumonia has a penicillin allergy. Which medication would you question if ordered?

A. Ceftriaxone

B. Azithromycin

C. Clindamycin

D. Levofloxacin

Ceftriaxone is a cephalosporin antibiotic, which shares a structure similar to penicillin. In patients with a documented penicillin allergy, there is a potential risk for cross-reactivity with cephalosporins, although the risk is relatively low, especially with those in later generations like ceftriaxone. However, the caution stems from the potential for serious allergic reactions, which could occur if the patient is indeed hypersensitive to penicillin. In contrast, azithromycin, clindamycin, and levofloxacin are not structurally related to penicillin and do not typically present this cross-reactivity risk, making them safer alternatives for treating pneumonia in patients with a penicillin allergy. Selecting a medication that avoids possible allergic reactions is crucial in providing safe and effective care. Thus, questioning the use of ceftriaxone in a patient with a known penicillin allergy is a prudent approach to avoid potential complications.

8. What does a "flat" ECG line indicate?

A. Normal sinus rhythm

B. Bradycardia

C. Asystole or absence of electrical activity in the heart

D. Atrial fibrillation

A "flat" ECG line, also known as asystole, indicates the absence of electrical activity in the heart. This condition signifies that there is no effective cardiac output, resulting in a lack of blood circulation to vital organs. In a flatline situation, the heart is not generating any impulses that would typically coordinate contractions of the heart muscle. Asystole is a critical emergency, often necessitating immediate intervention, such as cardiopulmonary resuscitation (CPR) and possible advanced cardiac life support (ACLS). In contrast, normal sinus rhythm reflects a healthy heart rhythm with organized electrical activity, bradycardia refers to an unusually slow heart rate with underlying electrical activity, and atrial fibrillation shows a chaotic rhythm with irregular heartbeats. Therefore, it is clear that a flat ECG line specifically signifies asystole, marking a serious medical condition requiring urgent care.

SAMPLE

9. An elderly woman shows fatigue and shortness of breath similar to previous urinary tract infection symptoms. What diagnostic test is crucial to obtain during triage?

A. X-ray of the chest

B. Complete blood count

C. ECG

D. Urinalysis

When an elderly woman presents with fatigue and shortness of breath that mirrors symptoms she has experienced in the context of previous urinary tract infections, a critical diagnostic test to perform during triage is a urinalysis. This assessment is particularly vital because urinary tract infections (UTIs) can lead to systemic symptoms, including changes in mental status, fatigue, and respiratory distress, especially in older adults. A urinalysis helps determine whether a new or recurring UTI may be the underlying cause of her current symptoms. This test can identify the presence of bacteria, white blood cells, and other indicators of infection. Given the close relationship between urinary infections and the presentation of fatigue or respiratory symptoms in this age group, it is essential to evaluate for an active infection first. While the other tests listed may provide valuable information in other contexts (such as ruling out cardiac issues with an ECG or assessing overall health with a complete blood count), the immediate need in this scenario is to ascertain whether the fatigue and shortness of breath are associated with a urinary infection, making urinalysis the most relevant and urgent diagnostic test during triage.

10. What is the first-line treatment for a patient experiencing a myocardial infarction?

A. Administration of oxygen

B. Administration of aspirin

C. Insertion of a stent

D. Placement on telemetry monitoring

The first-line treatment for a patient experiencing a myocardial infarction is the administration of aspirin. Aspirin plays a critical role in the management of acute coronary syndrome, including myocardial infarction, by inhibiting platelet aggregation. This action reduces the likelihood of further clot formation at the site of a ruptured atherosclerotic plaque, which is often the underlying cause of the infarction. Early administration can significantly improve outcomes by limiting the extent of the myocardial damage and improving blood flow to the affected area. In addition to aspirin, other interventions such as oxygen administration and telemetry monitoring are important components of the overall management plan in a clinical setting, but they do not have the same immediate impact on minimizing the damage caused by the infarction as aspirin does. Insertion of a stent is a more advanced intervention typically reserved for further along in the treatment pathway after initial stabilization and assessment. Thus, while all these interventions are vital in managing a myocardial infarction, the administration of aspirin is pivotal due to its rapid effect on clot prevention and stabilization.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://prophecyemergencydeptrn.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE