PROTECT EMT PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://protectemt.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. An 85-year-old patient presents with a forehead cut. Which question is important to ask during assessment?

A. Do you take blood thinners?

B. Do you have any allergies?

C. Do you take daily vitamins?

D. Do you have a fever?

2. Which of the following is part of SLUGEM?

A. Salivation

B. Fever

C. Weight loss

D. Dyspnea

3. Which statement best defines a transient ischemic attack (TIA)?

A. A transient blockage in a cerebral artery

B. A permanent stroke

C. An aneurysm in the brain

D. A spinal cord injury

4. Which best describes epinephrine's mechanism of action?

A. Vasodilation and bronchoconstriction

B. Vasoconstriction and bronchodilation

C. No effect on vessels or airways

D. Increases mucus production

5. Which item is part of Cushing's triad, a sign of potential brain herniation?

A. Unequal pupils

B. Altered mental status

C. CSF leakage

D. Cushing's triad

SAMPLE

6. In a trauma patient in shock with adequate breathing, what is the first action?

A. Stop the bleed

B. Open the airway

C. Place a blanket

D. Administer high-flow oxygen

7. In a patient with anaphylaxis who is unconscious, what is the top priority?

A. Breathing

B. Circulation

C. Airway

D. Defibrillation

8. After a fall with possible spinal injury, unconsciousness, neck and back pain, and weakness in arms and legs, what should you do first?

A. Check breathing and pulse

B. Obtain vital signs

C. Spinal motion restriction

D. Call for a higher level of care

9. In multisystem trauma, which step follows the primary survey and C-spine assessment during transport?

A. Secondary survey

B. Rest of care in route

C. Tertiary survey

D. Recheck of airway

10. What should you do after every call?

A. File a report only

B. Clean the clinic

C. Restock the ambulance

D. Schedule the next shift

SAMPLE

Answers

SAMPLE

1. A
2. A
3. A
4. B
5. D
6. A
7. C
8. C
9. A
10. C

SAMPLE

Explanations

SAMPLE

1. An 85-year-old patient presents with a forehead cut. Which question is important to ask during assessment?

A. Do you take blood thinners?

B. Do you have any allergies?

C. Do you take daily vitamins?

D. Do you have a fever?

When evaluating a forehead cut in an elderly patient, asking about blood thinners is critical because these medications significantly increase bleeding risk. If a patient is on anticoagulants or antiplatelets, even a small laceration can bleed more and longer, making hemostasis harder and hastening the need for rapid transport to a facility for assessment and potential reversal of anticoagulation. Knowing this upfront guides how you manage the wound—how long you apply direct pressure, whether you need additional measures to control bleeding, and how you communicate with the receiving team about possible complications. While asking about allergies, vitamins, or fever can be useful in other contexts, they don't directly change the immediate assessment and bleeding-control plan the way anticoagulant use does.

2. Which of the following is part of SLUGEM?

A. Salivation

B. Fever

C. Weight loss

D. Dyspnea

SLUGEM lists the muscarinic effects seen with excess acetylcholine, focusing on glandular secretions and related parasympathetic signs. The S in SLUGEM stands for Salivation, which is exactly what this option describes. Fever, weight loss, and dyspnea are not part of these cholinergic secretory signs and don't fit the mnemonic. In situations like organophosphate exposure, you'd expect increased secretions such as saliva, tears, urine, GI upset, vomiting, and pupil constriction; salivation is the specific sign highlighted by this option.

3. Which statement best defines a transient ischemic attack (TIA)?

A. A transient blockage in a cerebral artery

B. A permanent stroke

C. An aneurysm in the brain

D. A spinal cord injury

A transient ischemic attack happens when blood flow to a part of the brain is briefly interrupted, causing stroke-like symptoms that disappear once the flow returns. The key here is that the deficits resolve completely and quickly, usually within minutes to a few hours, leaving no lasting damage. Because it's a temporary event, a TIA serves as a warning sign that a more serious stroke could occur if not addressed promptly, so urgent medical evaluation is essential. In contrast, a real stroke involves permanent brain tissue injury and lasting deficits; an aneurysm is a bulging vessel that can rupture and cause bleeding in the brain; a spinal cord injury affects the spinal cord rather than the brain and produces different symptoms.

SAMPLE

4. Which best describes epinephrine's mechanism of action?

A. Vasodilation and bronchoconstriction

B. Vasoconstriction and bronchodilation

C. No effect on vessels or airways

D. Increases mucus production

Epinephrine works by stimulating adrenergic receptors to produce vasoconstriction in the blood vessels and bronchodilation in the airways. Activation of alpha-1 receptors on vascular smooth muscle causes vasoconstriction, helping reduce edema and support blood pressure. Activation of beta-2 receptors on bronchial smooth muscle causes bronchodilation, opening the airways for easier breathing. This combination explains why vasoconstriction and bronchodilation is the best description of its mechanism. It does not cause bronchoconstriction, it does cause vasoconstriction rather than dilation in the systemic circulation, and it does not increase mucus production.

5. Which item is part of Cushing's triad, a sign of potential brain herniation?

A. Unequal pupils

B. Altered mental status

C. CSF leakage

D. Cushing's triad

Cushing's triad is a late warning sign of rising intracranial pressure and potential brain herniation, reflecting brainstem compromise. It consists of three coordinated changes: elevated systolic blood pressure with a widened pulse pressure, slow heart rate (bradycardia), and irregular or slowed breathing. This combination shows the body's attempt to preserve cerebral perfusion while the brainstem is stressed. Among the options, the item that directly represents this specific triad is the triad itself—it's the defining cluster of signs indicating critical intracranial pressure. The other options aren't part of the classic triad: unequal pupils can occur with brain injury but not as a component of Cushing's triad; altered mental status is common with brain injury but not the triad's distinct combination; CSF leakage indicates a breach in containment rather than the triad.

6. In a trauma patient in shock with adequate breathing, what is the first action?

A. Stop the bleed

B. Open the airway

C. Place a blanket

D. Administer high-flow oxygen

In this scenario, the most important thing is to stop life-threatening blood loss. When a trauma patient is in shock, the immediate threat to survival is hemorrhage, which rapidly reduces circulating blood volume and impairs perfusion to vital organs. If you address the bleeding first—by applying direct pressure, using a tourniquet for severe limb wounds, and rapidly controlling hemorrhage—you tackle the root cause of the shock. Breathing is already adequate, so airway procedures aren't the priority right now, and while supplying oxygen is important, it doesn't reverse the rapid blood loss if bleeding continues. A blanket can help with warmth, but it doesn't save the patient from the consequences of ongoing hemorrhage. After bleeding is controlled, reassess the airway and breathing and continue supportive care.

SAMPLE

7. In a patient with anaphylaxis who is unconscious, what is the top priority?

A. Breathing

B. Circulation

C. Airway

D. Defibrillation

Securing the airway is the immediate priority. In anaphylaxis, swelling of the upper airway and bronchospasm can rapidly block airflow, and an unconscious patient cannot protect or clear the airway on their own. If the airway isn't open, you can't effectively ventilate or oxygenate, and the risk of aspiration is high. So the first actions focus on airway protection and oxygen delivery—positioning for best airflow, suctioning secretions, and preparing for airway adjuncts or advanced airway management as needed—while you promptly treat the underlying reaction with epinephrine. Breathing and circulation are also important, but they depend on having a patent airway, and defibrillation isn't the immediate need unless a cardiac arrest with a shockable rhythm occurs.

8. After a fall with possible spinal injury, unconsciousness, neck and back pain, and weakness in arms and legs, what should you do first?

A. Check breathing and pulse

B. Obtain vital signs

C. Spinal motion restriction

D. Call for a higher level of care

When there's a fall with possible spinal injury and signs like unconsciousness, neck and back pain, and weakness in the limbs, the priority is to prevent any movement of the spine. Immobilizing the spine right away—stabilizing the head and neck, maintaining alignment, and using a collar or long spine board as available—reduces the risk of worsening a spinal cord injury from movement. Once the spine is immobilized, you can then address airway and breathing and call for higher care. The danger of movement here is the main reason this is the best first action.

9. In multisystem trauma, which step follows the primary survey and C-spine assessment during transport?

A. Secondary survey

B. Rest of care in route

C. Tertiary survey

D. Recheck of airway

After the initial life threats are addressed in the primary survey and the C-spine is secured, the next step is the secondary survey. This is a thorough head-to-toe assessment that looks for injuries not immediately life-threatening but still critical to identify and manage. It includes repeating vital signs, a detailed examination of all body regions, and obtaining a focused history (such as what happened, medications, allergies, past medical history, and last meal). The goal is to uncover hidden injuries, guide ongoing care during transport, and plan definitive treatment at the receiving facility. Rechecking airway is an ongoing part of care but not a distinct next phase after the primary survey; airway assessment and management continue as needed. The tertiary survey refers to a later re-evaluation in the hospital to catch injuries missed earlier, not the immediate next step.

SAMPLE

10. What should you do after every call?

A. File a report only

B. Clean the clinic

C. Restock the ambulance

D. Schedule the next shift

Restocking the ambulance is the action that keeps you ready for the next patient. After a call, replenishing supplies used, reloading medications (as allowed by policy), replacing worn or used items, and ensuring equipment is clean, charged, and in working order ensures you can start the next run without delay. This readiness directly supports rapid, continuous patient care and safety for the next encounter. Documentation and decontamination are important, but restocking targets the immediate preparedness of the unit. Filing a report is essential for records, cleaning the clinic isn't typically a field duty, and shifting schedules is handled by supervisors rather than the crew in the field.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://protectemt.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE