PSYCHIATRIC TECHNICIAN INTERVIEW PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://psychtechinterview.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which of the following is a protective factor for suicide risk?

A. Previous suicide attempts

B. Hopelessness

C. Access to means

D. Strong social support

2. When escalating assaultive behavior, which action is correct regarding restraints?

A. Apply restraints immediately without order

B. Obtain a physician's order before seclusion/restraint if necessary

C. Use chemical restraint as first option

D. Use only verbal de-escalation and ignore safety

3. In addressing stigma concerns, which statement best reflects an appropriate initial response?

A. Dismiss concerns.

B. Insist on immediate acceptance.

C. Validate feelings.

D. Blame the patient.

4. Which of the following is NOT an active listening technique?

A. Open-ended questions

B. Reflective listening

C. Empathic statements

D. Providing direct orders

5. Which statement best describes panic disorder?

A. Worry about multiple domains lasting more than six months.

B. Recurrent unexpected panic attacks with fear of recurrence and avoidance.

C. Delusions and disorganized thinking.

D. Persistent sadness and anhedonia.

SAMPLE

6. Which symptom is listed among Risperdal side effects including increased thirst and urination?

A. Increased thirst

B. Hair loss

C. Ear pain

D. Joint swelling

7. Which technique is recommended for de-escalating an agitated patient?

A. Maintain a non-threatening posture and safe distance

B. Acknowledge emotions

C. Speak in a calm voice with short sentences

D. Provide choices and time to respond

8. In a patient who reports dizziness, hunger, and sweating, what is the most appropriate immediate action?

A. I would assist him to a chair and call for help from other staff including HSS and unit or RN

B. Notify shift supervisor and doctor or a doctor on call

C. I would give him juice or any sweet beverage and ask cafeteria staff to bring his food stat

D. Well the food is on its way I would assess the client and determine if he had a history of hypoglycemia since the symptoms are a classic sign of hypoglycemia but dizzy and sweating are also symptoms of heat exhaustion

9. When would you stop performing CPR?

A. When you are exhausted and cannot continue

B. When trained personnel take over

C. When the patient is resuscitated and breathing and circulation has resumed

D. When a physician orders you to stop

10. Which item is not typically included in a crisis plan for self-harm risk?

A. Warning signs

B. Coping strategies

C. Supportive contacts

D. Medical diagnosis label

SAMPLE

Answers

SAMPLE

1. D
2. B
3. C
4. D
5. B
6. A
7. C
8. C
9. C
10. D

SAMPLE

Explanations

SAMPLE

1. Which of the following is a protective factor for suicide risk?

A. Previous suicide attempts

B. Hopelessness

C. Access to means

D. Strong social support

Strong social support helps reduce suicide risk by boosting coping and resilience during difficult times. When people have close connections—family, friends, peers, or community ties—they receive emotional comfort, practical help, and a sense of belonging. This social buffer makes it easier to manage stress, decreases feelings of isolation, and increases the likelihood of seeking help and sticking with treatment plans. In practice, building and maintaining these supportive relationships is a key way to protect against suicidal crises. By contrast, a history of suicide attempts, persistent hopelessness, and easy access to means all heighten risk because they reflect greater vulnerability, ongoing distress, and greater opportunity to act on suicidal impulses.

2. When escalating assaultive behavior, which action is correct regarding restraints?

A. Apply restraints immediately without order

B. Obtain a physician's order before seclusion/restraint if necessary

C. Use chemical restraint as first option

D. Use only verbal de-escalation and ignore safety

Restraints are a last-resort safety measure and must be used only with proper authorization. In most settings, you cannot apply seclusion or physical restraints without a physician's order. If there is immediate danger, you may take necessary steps to protect the patient and others, but you must obtain the order as soon as possible and ensure continuous monitoring and documentation. This requirement protects patient rights, supports safety, and ensures that restraints are used only when truly needed and in a controlled, reviewable way. Understanding the workflow helps: de-escalation and other less restrictive options should be attempted first, and restraints are reserved for situations where those efforts fail to prevent imminent harm. A chemical restraint also requires a physician's order and is not a first option; it carries its own monitoring and risk considerations. Conversely, relying solely on verbal de-escalation while ignoring safety breaches is not appropriate, because protecting people from harm is a priority and restraints (when used) must be justified and supervised. So the best practice is to obtain a physician's order before using seclusion or restraints, applying the intervention only when necessary, and following through with proper monitoring, documentation, and reassessment.

SAMPLE

3. In addressing stigma concerns, which statement best reflects an appropriate initial response?

A. Dismiss concerns.

B. Insist on immediate acceptance.

C. Validate feelings.

D. Blame the patient.

The main skill here is empathetic validation when someone expresses concerns about stigma. The best initial response is to acknowledge and validate their feelings, which shows respect, reduces defensiveness, and builds trust. For example, you might say, "I hear how hard this has been for you, and it makes sense you'd feel upset given what you've faced." This normalizes their experience and creates a safe space for further conversation, making it easier to address stigma and discuss coping strategies or support. Dismissing concerns shuts down the conversation and signals that their experience isn't valued, which can deepen feelings of isolation. Insisting on immediate acceptance can pressure the person and overlook their readiness to process the emotions, potentially causing resistance. Blaming the patient reinforces stigma and damages the therapeutic relationship, making it harder to engage in any constructive discussion afterward. So, validating feelings first lays the groundwork for open communication, trust, and collaborative strategies to cope with stigma.

4. Which of the following is NOT an active listening technique?

A. Open-ended questions

B. Reflective listening

C. Empathic statements

D. Providing direct orders

Active listening focuses on techniques that encourage the person to talk and feel understood. Open-ended questions invite more detail and avoid simple yes/no answers. Reflective listening shows you understand by restating what was said and mirroring the person's feelings. Empathic statements acknowledge the person's emotions and help build trust. Providing direct orders is not an active listening technique because it directs behavior rather than inviting dialogue or exploring the person's experience; it can feel controlling and shut down communication. In urgent safety moments, clear instructions may be necessary, but that action isn't part of active listening.

5. Which statement best describes panic disorder?

A. Worry about multiple domains lasting more than six months.

B. Recurrent unexpected panic attacks with fear of recurrence and avoidance.

C. Delusions and disorganized thinking.

D. Persistent sadness and anhedonia.

Panic disorder is defined by recurrent, unexpected panic attacks and the ongoing fear of having more attacks, often leading to avoidance of situations that might trigger them. The statement that describes attacks happening repeatedly and without warning, plus persistent concern about recurrence and avoidance, fits this diagnosis best. The other descriptions map to different conditions: widespread worry about many areas for a long time points to generalized anxiety; delusions and disorganized thinking suggest a psychotic disorder; and persistent sadness with loss of interest indicates depression.

SAMPLE

6. Which symptom is listed among Risperdal side effects including increased thirst and urination?

A. Increased thirst

B. Hair loss

C. Ear pain

D. Joint swelling

Risperdal can affect metabolic processes and, in some people, raise blood glucose. When glucose is high, the body may respond with increased thirst and increased urination. Among the given options, increased thirst fits this known side effect pattern of risperidone. Hair loss, ear pain, and joint swelling are not typically linked to Risperdal as common adverse effects. If a patient reports thirst and urination, it's important to assess for hyperglycemia or diabetes and consider medical evaluation or testing, with possible medication adjustment if needed.

7. Which technique is recommended for de-escalating an agitated patient?

A. Maintain a non-threatening posture and safe distance

B. Acknowledge emotions

C. Speak in a calm voice with short sentences

D. Provide choices and time to respond

Centering the interaction on how you communicate is key. When someone is agitated, their brain is in a heightened state, and messages that come out in a calm voice with short sentences help reduce that arousal. Short, uncomplicated statements are easier to process, leave less room for misinterpretation, and provide clear, actionable instructions that the person can follow without feeling overwhelmed. This approach signals safety and control, which helps the patient settle more quickly and co-create a safer space for dialogue. Other strategies matter and fit into a broader de-escalation plan—non-threatening posture and safe distance set the stage, and acknowledging emotions validates the person's experience. Providing choices and time to respond supports autonomy, but these are most effective when paired with the calm, concise way of speaking that lowers arousal and clarifies expectations from the start.

SAMPLE

8. In a patient who reports dizziness, hunger, and sweating, what is the most appropriate immediate action?

A. I would assist him to a chair and call for help from other staff including HSS and unit or RN

B. Notify shift supervisor and doctor or a doctor on call

C. I would give him juice or any sweet beverage and ask cafeteria staff to bring his food stat

D. Well the food is on its way I would assess the client and determine if he had a history of hypoglycemia since the symptoms are a classic sign of hypoglycemia but dizzy and sweating are also symptoms of heat exhaustion

When dizziness, hunger, and sweating occur, the most likely issue is hypoglycemia, and the priority is to raise the blood glucose quickly. Providing a fast-acting carbohydrate right away—such as juice or another sweet beverage—addresses the root problem immediately and can reverse symptoms before they worsen. After giving the juice, you can arrange for a more substantial snack or meal to maintain glucose levels and reassess the patient's status. If possible, check blood glucose to confirm improvement and continue to monitor. If the patient cannot swallow or becomes unconscious, oral sugar is not safe, and the next steps would be to use IV dextrose or intramuscular glucagon per protocol and call for help. Other actions that focus on notifying others or waiting for meals do not treat the immediate danger of low blood sugar.

9. When would you stop performing CPR?

A. When you are exhausted and cannot continue

B. When trained personnel take over

C. When the patient is resuscitated and breathing and circulation has resumed

D. When a physician orders you to stop

The main idea is that CPR is continued to support circulation and breathing until the person's own heart and lungs can resume functioning. The clearest, most objective stopping point is when spontaneous circulation and breathing return—that is, the patient is breathing on their own and a pulse is present. At that moment, external chest compressions and rescue efforts are no longer needed, and care should shift to monitoring and definitive treatment. While you may stop if trained responders take over or a physician orders you to stop, those are situational and depend on the context; they're not as definitive as the return of spontaneous circulation.

10. Which item is not typically included in a crisis plan for self-harm risk?

A. Warning signs

B. Coping strategies

C. Supportive contacts

D. Medical diagnosis label

A crisis plan is designed to guide someone through a self-harm risk moment with practical, action-oriented steps. Warning signs are included so you can spot when risk is rising and intervene early. Coping strategies provide concrete tools to manage urges in the moment without harming yourself. Supportive contacts list trusted people or professionals to reach out to and where to seek help quickly. The medical diagnosis label, while important for treatment records, doesn't offer immediate, actionable guidance for staying safe during a crisis and can feel irrelevant or private. It's not typically part of the crisis-plan content because the focus is on what you can do right now and who can help, rather than on medical labeling.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://psychtechinterview.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE