
PSYCHOLOGY STATISTICS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://psychologystats.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. If a set of exam scores forms a negatively skewed distribution, what can you conclude about the students' scores?

A. Most of the students had relatively high scores

B. Most of the students had relatively low scores

C. About 50% had high scores and the rest low

D. It is not possible to draw any conclusions about students' scores

2. Which expression represents the sum of the X values squared?

A. ΣX^2

B. (ΣX)^2

C. Σ(X^2)

D. ΣX

3. If the scores are 3, 7, and 5, what is ΣX?

A. 13

B. 15

C. 25

D. 30

4. Which type of statistics summarizes or describes the main features of a dataset?

A. Descriptive statistics

B. Inferential statistics

C. Descriptive and inferential

D. Experimental statistics

5. When the goal is to determine the average number of hours American adolescents spend watching television, the resulting value describes the population and is known as a _____.

A. Parameter

B. Statistic

C. Population

D. Sample

SAMPLE

6. What is a p-value?

A. The probability of obtaining data as extreme as observed (or more) assuming the null hypothesis is true.

B. The probability that the null hypothesis is true.

C. The observed effect size.

D. The probability of observing data under the alternative hypothesis.

7. What are the real limits for a score of X = 42, given measurements are recorded to the nearest half unit?

A. 41.75 and 42.25

B. 41.5 and 42.0

C. 41.75 and 42.0

D. 41.5 and 42.25

8. In regression analysis, what does a standardized beta coefficient indicate?

A. Indicates the expected change in the dependent variable in standard deviations for a one SD change in the predictor, holding other predictors constant.

B. Shows the raw units change in Y per unit change in X.

C. Represents the correlation between X and Y.

D. Measures the proportion of variance explained by the predictor.

9. What is the shape of the distribution for the data: 1, 2, 3, 3, 4, 4, 4, 5, 5, 5, 5, 6?

A. Symmetrical

B. Positively skewed

C. Negatively skewed

D. Normal

10. Which is an example of a discrete variable?

A. The amount of weight gained for each freshman at a local college

B. The gender of each student in a psychology class

C. The age of each student in a psychology class

D. The amount of time to solve a problem

SAMPLE

Answers

SAMPLE

1. A
2. A
3. B
4. A
5. A
6. A
7. A
8. A
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. If a set of exam scores forms a negatively skewed distribution, what can you conclude about the students' scores?

A. Most of the students had relatively high scores

B. Most of the students had relatively low scores

C. About 50% had high scores and the rest low

D. It is not possible to draw any conclusions about students' scores

When a distribution is negatively skewed, the long tail appears on the left with a cluster of scores toward the high end. This means most students earned relatively high scores, while a few very low scores pull the average down. So the best takeaway is that most students did well. The other ideas don't fit: a left-skew pattern doesn't indicate most scores are low, and a roughly even split would imply symmetry rather than skewness. Additionally, in this pattern the mean tends to be lower than the median due to those low outliers, reinforcing that the bulk of scores are high.

2. Which expression represents the sum of the X values squared?

A. ΣX^2

B. (ΣX)^2

C. Σ(X^2)

D. ΣX

Sum of squares means you square each data point first, then add them all up. So for observations X1, X2, ..., Xn, the sum of the squares is X1^2 + X2^2 + ... + Xn^2, which is what the notation Σ X^2 (or Σ(X^2)) expresses. This is different from squaring the total sum, (Σ X)^2, which would give extra cross terms from expanding the square of a sum (like 2·X1·X2, etc.). Thus the expression with the square on each X inside the summation is the correct representation of the sum of the X values squared.

3. If the scores are 3, 7, and 5, what is ΣX?

A. 13

B. 15

C. 25

D. 30

ΣX is the total of all observed values. Add the given scores: 3 + 7 + 5 equals 15. So ΣX is fifteen. If you later needed the average, you'd divide by the number of scores (3), which would give an average of 5.

SAMPLE

4. Which type of statistics summarizes or describes the main features of a dataset?

A. Descriptive statistics

B. Inferential statistics

C. Descriptive and inferential

D. Experimental statistics

Descriptive statistics focus on summarizing what the data look like. They provide a concise picture of a dataset through measures of central tendency (like the mean or median), variability (such as the standard deviation or range), and the overall distribution (histograms or frequency tables). Since the question asks for statistics that summarize or describe the main features of a dataset, descriptive statistics fit best because they directly describe the data at hand. Inferential statistics, by contrast, are about drawing conclusions beyond the data to a larger population, which isn't about describing the dataset itself. The option that combines descriptive and inferential isn't as precise for this prompt, and experimental statistics isn't a standard category.

5. When the goal is to determine the average number of hours American adolescents spend watching television, the resulting value describes the population and is known as a _____.

A. Parameter

B. Statistic

C. Population

D. Sample

This item is about the difference between a parameter and a statistic. A parameter is a numerical summary that describes a whole population—the entire group you're interested in. In this scenario, if you could measure every American adolescent and compute their average hours of TV watching, that average would be a population parameter. If you instead collected data from just a subset and computed its average, that value would be a statistic—an estimate of the population parameter that can vary from sample to sample. The population is the group of interest, while a sample is just a part of that group.

6. What is a p-value?

A. The probability of obtaining data as extreme as observed (or more) assuming the null hypothesis is true.

B. The probability that the null hypothesis is true.

C. The observed effect size.

D. The probability of observing data under the alternative hypothesis.

The p-value measures how surprising your data would be if the null hypothesis were true. It is the probability of obtaining data as extreme as the observed data (or more extreme) under that assumption. In other words, it tells you how compatible your data are with the notion that there is no real effect. This is not the probability that the null hypothesis is true itself, which is a common misinterpretation. The p-value is a statement about the data given the null, not about the truth of the null. It also does not reflect the size of the effect—that's the role of the observed (or estimated) effect size. And it isn't the probability of the data under the alternative hypothesis. So, a small p-value means the observed result would be unlikely if the null were true, which is why researchers often consider it evidence against the null at a chosen significance level.

SAMPLE

7. What are the real limits for a score of X = 42, given measurements are recorded to the nearest half unit?

A. 41.75 and 42.25

B. 41.5 and 42.0

C. 41.75 and 42.0

D. 41.5 and 42.25

Rounding to the nearest 0.5 sets clear cutoffs around the observed value. The true score that would still be recorded as 42 must lie between halfway marks to the neighboring recorded values. The midpoint between 41.5 and 42.0 is 41.75, and the midpoint between 42.0 and 42.5 is 42.25. So any true score from 41.75 up to 42.25 would be rounded to 42 (with the lower bound included and the upper bound effectively excluded). This gives the real-limit interval of 41.75 to 42.25, which is why the correct bounds are 41.75 and 42.25.

8. In regression analysis, what does a standardized beta coefficient indicate?

A. Indicates the expected change in the dependent variable in standard deviations for a one SD change in the predictor, holding other predictors constant.

B. Shows the raw units change in Y per unit change in X.

C. Represents the correlation between X and Y.

D. Measures the proportion of variance explained by the predictor.

Standardized beta is about effect size in standardized units. When all variables are transformed to z-scores, the regression slope tells you how many standard deviation units the dependent variable would change for a one standard deviation increase in the predictor, while holding the other predictors in the model constant. This makes the sizes of different predictors directly comparable, since they're all measured on the same scale. So, the coefficient conveys the unique, adjusted impact of that predictor on the outcome, independent of the original measurement units. It's not giving a raw unit change in Y, and it's not simply the correlation between X and Y (which doesn't account for other predictors in the model). In simple (one-predictor) regression, the standardized beta and the correlation can align, but in multiple regression they reflect the predictor's effect after controlling for others, expressed in standard deviation units.

9. What is the shape of the distribution for the data: 1, 2, 3, 3, 4, 4, 4, 5, 5, 5, 5, 6?

A. Symmetrical

B. Positively skewed

C. Negatively skewed

D. Normal

Skewness is about where the tail of a distribution extends. In this set, most values cluster toward the higher end (4s and 5s), while only a couple of values are very small (1 and 2). That creates a longer tail to the left, toward smaller numbers. The mean is slightly less than the median (mean about 3.92, median 4), which also signals a left, or negative, skew. So the data are best described as negatively skewed.

SAMPLE

10. Which is an example of a discrete variable?

A. The amount of weight gained for each freshman at a local college

B. The gender of each student in a psychology class

C. The age of each student in a psychology class

D. The amount of time to solve a problem

The main idea is distinguishing discrete versus continuous variables. A discrete variable takes on distinct, separate values, often categories or countable numbers, rather than a smooth range. Gender is a discrete variable because it represents a finite set of categories (such as male, female, nonbinary, etc.) rather than a measured amount on a continuous scale. The other options involve quantities that can vary fluidly and take on many values within a range: weight gained can be any real number on a scale, age is measured as a precise value (often with months and days, not just whole numbers), and time to solve a problem is a continuous measure that can be expressed with fractions of a second. So gender stands out as the discrete example.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://psychologystats.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE