PULMONARY EMERGENCIES PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://pulmonaryemergencies.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which finding supports an intermediate-risk pulmonary embolism diagnosis in a normotensive patient?

A. Hypotension

B. Normal troponin and BNP

C. RV dysfunction on echo or CT

D. Stable respiratory status

2. Which condition is a must-not-miss differential diagnosis for hemoptysis?

A. Pulmonary embolism

B. Common cold

C. Sinusitis

D. Bronchitis

3. What would the ABG/VBG typically show in COPD patients at baseline?

A. Elevated PCO2 with metabolic compensation

B. Low PCO2

C. Normal PCO2

D. Decreased bicarbonate

4. Which imaging test is most commonly used to definitively diagnose PE in contemporary practice?

A. V/Q scan

B. Pulmonary angiography

C. CXR

D. CT Angiography

5. Abrupt dyspnea, tachycardia, pleuritic chest pain, hypoxia, hemoptysis, and lower-extremity pain are classic for which condition?

A. Pneumonia

B. Pulmonary embolism

C. Acute bronchitis

D. Pneumothorax

SAMPLE

6. What patient position is recommended for suspected pulmonary edema?

A. Supine

B. Left lateral decubitus

C. Prone

D. Upright in high Fowler

7. Which of the following is a risk factor for pulmonary embolism?

A. Regular exercise

B. Smoking

C. Prolonged immobility

D. Low-fat diet

8. Which disease is most likely associated with nocturnal symptoms and reversible airway obstruction?

A. Asthma

B. COPD

C. Pneumonia

D. Pulmonary edema

9. Among the outpatient CAP options, which choice represents a macrolide antibiotic?

A. Macrolide

B. Amoxicillin

C. Doxycycline

D. Levofloxacin

10. A pleural effusion is typically detectable on chest radiograph when the pleural fluid volume reaches approximately how many milliliters?

A. 100 mL

B. 500 mL

C. 1000 mL

D. 250 mL

SAMPLE

Answers

SAMPLE

1. C
2. C
3. D
4. D
5. B
6. D
7. C
8. A
9. A
10. D

SAMPLE

Explanations

SAMPLE

1. Which finding supports an intermediate-risk pulmonary embolism diagnosis in a normotensive patient?

A. Hypotension

B. Normal troponin and BNP

C. RV dysfunction on echo or CT

D. Stable respiratory status

In pulmonary embolism, how a patient looks in terms of blood pressure guides risk: those who are unstable with hypotension are high risk (massive PE). Normotensive patients can still be at higher risk if the right ventricle is under strain. Finding right ventricular dysfunction on imaging shows the heart is struggling to cope with the sudden blockage in the lungs, meaning there is myocardial stress even though blood pressure is normal. This pattern places the patient in an intermediate (submassive) risk category because the right ventricle's failure to handle the increased afterload signals a higher chance of deterioration, guiding closer monitoring and consideration of more aggressive management in select cases. Hypotension would push you toward high-risk classification because it reflects hemodynamic collapse. Normal troponin and BNP suggest no myocardial injury, which lowers concern for risk, not increases it. Stable respiratory status doesn't specifically indicate right heart strain, so it doesn't pinpoint the risk level the same way RV dysfunction does.

2. Which condition is a must-not-miss differential diagnosis for hemoptysis?

A. Pulmonary embolism

B. Common cold

C. Sinusitis

D. Bronchitis

When assessing hemoptysis, you have to consider whether the bleeding is actually coming from the lower airways or from the upper airway. Blood from the nose or sinuses can drain into the throat and be coughed up as blood-tinged sputum, so sinusitis should be included as a must-not-miss possibility. If sinusitis is suspected, you'd typically see ENT-type symptoms such as facial pressure or pain, purulent nasal discharge, or sinus tenderness, and the chest may appear normal. This distinction matters because the management shifts from pulmonary-directed workup to treating the sinus infection and addressing the upper airway source. The other options are less likely to explain true hemoptysis on their own: a common cold or bronchitis mainly cause cough with mucus but not typically frank blood, and pulmonary embolism, while dangerous and important to consider, is a different clinical pathway that would usually present with more acute chest symptoms and embolic risk features.

3. What would the ABG/VBG typically show in COPD patients at baseline?

A. Elevated PCO2 with metabolic compensation

B. Low PCO2

C. Normal PCO2

D. Decreased bicarbonate

Chronic COPD leads to long standing CO2 retention, so the body compensates by the kidneys increasing bicarbonate. This means the baseline ABG/VBG typically shows elevated PCO2 with an elevated bicarbonate level, and the pH is often near normal or only mildly acidic because the metabolic compensation buffers the respiratory acidosis. In numbers, you might see PCO2 appreciably above normal (often 45–60 mmHg) with bicarbonate raised (roughly 28–34 mEq/L). Decreased bicarbonate would point to a primary metabolic acidosis—not what COPD baseline usually presents. A low or normal PCO2 would not fit the chronic CO2-retaining state of COPD.

SAMPLE

4. Which imaging test is most commonly used to definitively diagnose PE in contemporary practice?

A. V/Q scan

B. Pulmonary angiography

C. CXR

D. CT Angiography

CT pulmonary angiography is the imaging study most commonly used today to definitively diagnose pulmonary embolism. It directly visualizes clots within the pulmonary arteries and can also reveal alternative causes of symptoms, all in a rapid, widely available, and noninvasive test. Modern multidetector CTA has high sensitivity and specificity, making it the go-to test to confirm or exclude PE in most patients without needing invasive procedures. V/Q scanning can be helpful when contrast is a concern (such as iodine allergy or pregnancy), but its results are more often nondiagnostic or indeterminate, which makes it less definitive in routine practice. Pulmonary angiography is the historical gold standard but is invasive and carries more risk, so CTA has largely replaced it as the initial diagnostic tool. Chest X-ray, while useful to rule out other causes, does not diagnose PE.

5. Abrupt dyspnea, tachycardia, pleuritic chest pain, hypoxia, hemoptysis, and lower-extremity pain are classic for which condition?

A. Pneumonia

B. Pulmonary embolism

C. Acute bronchitis

D. Pneumothorax

Abrupt dyspnea with pleuritic chest pain and hypoxia, together with hemoptysis and lower-extremity pain, points to a pulmonary embolism. The scenario fits because a clot usually forms in the deep veins of the leg (deep vein thrombosis), breaks loose, and travels to the lungs, suddenly blocking pulmonary arteries. This acute obstruction triggers rapid breathing, chest pain that worsens with breaths, and reduced oxygen levels. Hemoptysis can occur when lung tissue becomes damaged or infarcted from the blocked blood supply. The leg pain supports the idea that the clot originated in the leg. Other conditions can cause chest pain or breathing difficulty, but they don't typically present with this combination of sudden dyspnea, pleuritic pain, hypoxemia, hemoptysis, and leg symptoms all together. Pneumonia usually brings fever and productive cough; acute bronchitis mainly causes cough without ongoing hypoxemia or leg-clot clues; pneumothorax involves sudden chest pain and unilateral findings like decreased breath sounds, but hemoptysis and leg DVT signs are not defining features.

SAMPLE

6. What patient position is recommended for suspected pulmonary edema?

A. Supine

B. Left lateral decubitus

C. Prone

D. Upright in high Fowler

Placing a patient with suspected pulmonary edema in an upright position helps relieve breathing difficulty quickly. When the head of the bed is raised (high Fowler, roughly 60 degrees or more), chest wall mechanics improve and the diaphragm can move more effectively, making it easier to take deep breaths. Upright positioning also reduces preload because gravity lowers venous return to the heart, which decreases left atrial pressure and the hydrostatic pressure in the pulmonary capillaries. Less pressure in the lungs means less fluid leakage into the alveoli and better oxygenation. Lying flat (supine) tends to worsen dyspnea in pulmonary edema by increasing venous return and worsening pulmonary congestion. The left lateral decubitus position doesn't provide the same relief and isn't specifically beneficial for pulmonary edema. Prone can help in certain severe respiratory conditions like ARDS, but it's not the standard initial position for edema-related dyspnea.

7. Which of the following is a risk factor for pulmonary embolism?

A. Regular exercise

B. Smoking

C. Prolonged immobility

D. Low-fat diet

Prolonged immobility promotes venous stasis, a key trigger for clot formation in the deep veins. When a clot forms in these veins, it can break loose and travel through the bloodstream to the lungs, where it blocks branches of the pulmonary artery—this is a pulmonary embolism. So this factor directly increases the likelihood of a clot forming and then embolizing to the lungs. Regular exercise tends to improve venous return and reduce stasis, lowering the risk of PE. A low-fat diet doesn't have a direct, established link to clot formation, and while smoking is a risk factor for many cardiovascular conditions, it is not as direct or as strong a contributor to venous thromboembolism as immobility-induced stasis.

8. Which disease is most likely associated with nocturnal symptoms and reversible airway obstruction?

A. Asthma

B. COPD

C. Pneumonia

D. Pulmonary edema

Nocturnal symptoms with reversible airway obstruction point to asthma due to airway hyperresponsiveness and inflammatory airways that fluctuate over time. In asthma, the airways tighten in response to triggers and inflammation, causing wheeze, coughing, and shortness of breath that often worsen at night or in the early morning. Importantly, this obstruction is largely reversible with bronchodilators and anti-inflammatory therapies, so lung function improves after treatment or with time. In contrast, COPD typically shows less reversibility, especially in longtime smokers, and does not characteristically present with nighttime symptom spikes. Pneumonia presents with infection-related signs such as fever and productive cough, not a primary, reversible obstructive pattern. Pulmonary edema causes dyspnea from fluid overload and cardiogenic signs, not reversible airway obstruction tied to bronchial hyperreactivity.

SAMPLE

9. Among the outpatient CAP options, which choice represents a macrolide antibiotic?

A. Macrolide

B. Amoxicillin

C. Doxycycline

D. Levofloxacin

Macrolides are a class of antibiotics that work by binding to the 50S ribosomal subunit to inhibit protein synthesis. They're commonly used in outpatient therapy for community-acquired pneumonia to cover atypical pathogens such as Mycoplasma and Chlamydophila. Among the options, the one that represents a macrolide antibiotic is the label that literally says "Macrolide," since the others belong to different classes: amoxicillin is a beta-lactam (penicillin-type), doxycycline is a tetracycline, and levofloxacin is a fluoroquinolone. So recognizing the class name points you to the macrolide. In practice, macrolides like azithromycin or clarithromycin may be chosen for outpatient CAP when atypical coverage is desired and there are no contraindications.

10. A pleural effusion is typically detectable on chest radiograph when the pleural fluid volume reaches approximately how many milliliters?

A. 100 mL

B. 500 mL

C. 1000 mL

D. 250 mL

When a pleural effusion becomes visible on a standard upright chest radiograph, it's because enough fluid has layered to blunt the costophrenic sulcus. The amount needed for detection is on the order of a few hundred milliliters, with about 250 mL being a common practical threshold for seeing a change on a routine film. Smaller amounts, around 100 mL, often don't produce a visible sign on a single view. Much larger volumes, like 500 mL or 1000 mL, would be readily apparent and can shift mediastinal structures or cause obvious symptoms. So, 250 mL is the best approximation for radiographic detectability on a typical chest X-ray. If more sensitivity is needed, ultrasound can detect much smaller volumes.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://pulmonaryemergencies.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE