RADIATION THERAPY BOARD PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://radiationtherapyboard.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **When treating the maxillary antrum, which organ is at the greatest risk for damage?**

   A. Pituitary

   B. Brain

   C. Lens

   D. Sphenoid

2. **Which type of radiation is most commonly used in external beam radiation therapy?**

   A. Alpha particles

   B. Beta particles

   C. X-rays

   D. Gamma rays

3. **Which of the following is not a likely side effect of cytotoxic chemotherapy drugs?**

   A. Nausea

   B. Cardiac toxicity

   C. Myelosuppression

   D. Stroke

4. **What type of radiation detector is used primarily for measuring high levels of radiation?**

   A. Geiger-Muller counter

   B. Pocket dosimeter

   C. Thermoluminescent dosimeter

   D. Ionization chamber

5. **Cells of malignant growth are commonly characterized as?**

   A. Anaplastic

   B. Well differentiated

   C. Slow growing

   D. Poorly oxygenated

SAMPLE

6. At which lumbar vertebra number is the iliac crest located?

A. 1

B. 2

C. 4

D. 5

7. What is the tolerance for daily check of linac output constancy?

A. 5%

B. 1%

C. 3%

D. 2%

8. Standard wedges vary in angle from which of the following degrees?

A. 15 to 75

B. 30 to 60

C. 45 to 75

D. 15 to 60

9. Most thyroid cancers develop from which cell type?

A. papillary

B. follicular

C. anaplastic

D. serous

10. Which sign of inflammation is characterized by warmth in the affected area?

A. Pain

B. Heat

C. Redness

D. Swelling

SAMPLE

Answers

SAMPLE

1. C
2. C
3. D
4. D
5. A
6. C
7. C
8. D
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. When treating the maxillary antrum, which organ is at the greatest risk for damage?

A. Pituitary

B. Brain

C. Lens

D. Sphenoid

The lens of the eye is considered to be at the greatest risk for damage when treating the maxillary antrum. This is primarily due to the anatomical proximity of the maxillary sinus to the orbit, where the lens is located. Radiation therapy directed towards the maxillary antrum can inadvertently expose the lens to radiation dose, which may lead to cataract formation or other ocular complications. While the other organs listed are relevant in the context of head and neck radiation therapy, they are generally located either further away or are not as susceptible to the specific field of treatment when the maxillary antrum is involved. The pituitary gland and the brain, although important structures, are positioned at a greater distance from the target area than the lens. The sphenoid sinus is also quite close, but it does not carry the same risk of direct radiation-induced damage as the lens due to the critical nature of vision and the established relationship between radiation exposure and cataract development. Thus, the lens is recognized as the organ at the highest risk during such treatments.

2. Which type of radiation is most commonly used in external beam radiation therapy?

A. Alpha particles

B. Beta particles

C. X-rays

D. Gamma rays

In external beam radiation therapy, the most commonly used type of radiation is X-rays. This is primarily due to their ability to penetrate tissue effectively, making them ideal for targeting tumors while sparing surrounding healthy tissue. X-rays can be generated with linear accelerators, which allow for precise dosing and the ability to create various treatment plans that align with patient-specific anatomy and the location of the tumor. Additionally, X-rays provide a therapeutic dose that can be finely tuned, allowing clinicians to deliver a focused beam to malignant cells while minimizing exposure to adjacent normal tissues. The versatility and effectiveness of X-rays make them the standard choice for many cancer treatments in external beam radiation therapy. In contrast, alpha particles are primarily used in targeted alpha therapy and have limited penetration, making them unsuitable for external beam applications. Beta particles also have limited penetration and are typically used for different treatment modalities. Gamma rays, while used in some radiation therapies (such as certain types of brachytherapy), are less common in external beam radiation therapy compared to X-rays.

SAMPLE

3. Which of the following is not a likely side effect of cytotoxic chemotherapy drugs?

A. Nausea

B. Cardiac toxicity

C. Myelosuppression

D. Stroke

Cytotoxic chemotherapy drugs primarily target rapidly dividing cells, which can lead to a number of well-documented side effects. Nausea is one of the most common side effects due to the impact of these drugs on the gastrointestinal tract and the brain's vomiting center. Myelosuppression is another significant side effect, as these drugs can impair the bone marrow's ability to produce blood cells, leading to conditions like anemia, thrombocytopenia, and leukopenia. Cardiac toxicity can occur with certain chemotherapeutic agents, particularly those known to affect cardiac function, such as doxorubicin. In contrast, while strokes are serious medical events, they are not directly caused by cytotoxic chemotherapy drugs. The occurrence of a stroke involves a complex interplay of factors such as vascular health, blood pressure, and clotting disorders, which are not directly related to the action of chemotherapy. Therefore, it is less likely to be associated with chemotherapy compared to the other options listed. This distinction is crucial as it helps in understanding the risk profiles of different treatment modalities in oncology.

4. What type of radiation detector is used primarily for measuring high levels of radiation?

A. Geiger-Muller counter

B. Pocket dosimeter

C. Thermoluminescent dosimeter

D. Ionization chamber

The ionization chamber is the correct choice for measuring high levels of radiation due to its ability to provide accurate readings across a wide range of exposure levels. This type of detector works by measuring the ion pairs created when radiation passes through a gas within the chamber. The gas becomes ionized, and the resulting current is proportional to the amount of radiation that has interacted with the gas, allowing for precise quantification of high radiation levels. Ionization chambers are particularly suited for applications where high radiation levels are encountered, such as in radiotherapy settings or during radiation safety assessments, because they can withstand and accurately measure these high exposures without saturating or damaging the detector. They can measure both photon and particle radiation effectively and are capable of providing both instantaneous and integrated dose measurements. In contrast, a Geiger-Muller counter, while useful for detecting low to moderate levels of radiation, can become saturated and provide inaccurate readings in high radiation environments. Pocket dosimeters are more suited for personal dosimetry and measuring lower levels of radiation exposure over time, making them less ideal for high-range measurements. Thermoluminescent dosimeters (TLDs) are primarily used for measuring accumulated doses over longer periods and are not designed for immediate high radiation readings. Thus, the unique

SAMPLE

5. Cells of malignant growth are commonly characterized as?

A. Anaplastic

B. Well differentiated

C. Slow growing

D. Poorly oxygenated

Cells of malignant growth are often characterized as anaplastic. Anaplastic cells, which are typically observed in malignant tumors, show a loss of differentiation and display abnormal shapes, sizes, and features that are markedly different from normal cells. This lack of differentiation is a hallmark of cancerous cells, indicating their invasive and aggressive nature. Anaplastic cells frequently exhibit rapid proliferation and tend to invade surrounding tissues, which is part of what makes them distinct from well-differentiated cells that resemble normal tissues and tend to grow more slowly. While poorly oxygenated cells can indeed be found in tumors because of their rapid growth outpacing their blood supply, this characteristic alone does not define malignant cells in the same way that anaplasia does. Thus, anaplasia serves as a critical indicator of the aggressive and disorganized behavior associated with malignant growth.

6. At which lumbar vertebra number is the iliac crest located?

A. 1

B. 2

C. 4

D. 5

The iliac crest is generally located at the level of the fourth lumbar vertebra (L4). This anatomical landmark is significant in various medical procedures, including lumbar punctures and spinal anesthesia, where identifying the right vertebral level is crucial for successful interventions. The iliac crest serves as a palpable landmark that helps clinicians identify the corresponding vertebral level and ensures accurate positioning during procedures involving the lower back. In clinical practice, understanding the relationship between the iliac crest and lumbar vertebrae aids in achieving proper patient positioning and can help avoid complications during interventions. Positions for imaging or surgical procedures often reference the iliac crest, further emphasizing its importance in anatomy and procedural accuracy.

7. What is the tolerance for daily check of linac output constancy?

A. 5%

B. 1%

C. 3%

D. 2%

The tolerance for daily checks of linear accelerator (linac) output constancy is indeed 3%. This percentage ensures that the output of the linac remains consistent and within acceptable limits on a daily basis, which is vital for delivering precise doses to patients in radiation therapy. Maintaining this level of accuracy is critical for the safety and effectiveness of treatment, as variations outside of this range could lead to underdosing or overdosing patients, impacting treatment outcomes. Regular checks of this magnitude support quality assurance protocols, enabling the detection of any potential drift or malfunction in the linac's performance. The 3% tolerance provides a balance between allowing for minor fluctuations due to environmental factors or equipment wear and ensuring that any significant issues are addressed promptly to maintain treatment integrity. In contrast, other percentages such as 1%, 2%, or 5% do not reflect the standard practice for daily checks of output constancy in radiation therapy, where a tighter tolerance of 1% may be applied for more critical situations or equipment, while 5% is generally considered too lenient for daily outputs, failing to adequately safeguard patient treatment accuracy.

SAMPLE

8. Standard wedges vary in angle from which of the following degrees?

A. 15 to 75

B. 30 to 60

C. 45 to 75

D. 15 to 60

Standard wedges used in radiation therapy typically have angles that range from 15 to 60 degrees. This range allows for the modification of the radiation beam profile, helping to achieve better dose distribution in the target area while minimizing exposure to surrounding healthy tissues. Wedges can be used to effectively create a more uniform dose distribution across the treatment volume, especially when dealing with irregularly shaped tumors or differing tissue densities. Using standard wedges within this specified range allows clinicians to tailor the therapy to individual patient needs, optimizing treatment outcomes. Wedges with angles less than 15 degrees or greater than 60 degrees are not standard and would not be commonly utilized or available in the typical clinical setting.

9. Most thyroid cancers develop from which cell type?

A. papillary

B. follicular

C. anaplastic

D. serous

Thyroid cancers primarily develop from follicular epithelial cells, which can differentiate into various types of cancer. Among these, papillary thyroid carcinoma is the most common subtype, accounting for approximately 80% of all thyroid cancers. This type of cancer originates from the follicular cells of the thyroid gland, which are responsible for the production of thyroid hormones. Papillary thyroid carcinoma is characterized by its distinct histological features, such as psammoma bodies and nuclear grooves, which help in its identification. The tumor typically has a good prognosis, as it tends to grow slowly and is less likely to metastasize aggressively compared to other thyroid cancer types. Understanding the cellular origin and behavior of papillary thyroid carcinoma is crucial for effective diagnosis and management in clinical practice.

10. Which sign of inflammation is characterized by warmth in the affected area?

A. Pain

B. Heat

C. Redness

D. Swelling

The sign of inflammation characterized by warmth in the affected area is heat. Inflammation is a biological response of tissues to harmful stimuli, and warmth occurs due to increased blood flow to the inflamed region. This is often a result of the body's immune response, where blood vessels dilate to allow more immune cells to reach the site of injury or infection. The increased metabolic activity and blood flow can lead to a noticeable increase in temperature, contributing to the feeling of warmth. The presence of heat is one of the classic signs of inflammation, alongside pain, redness, and swelling. Each of these signs reflects different aspects of the inflammatory process, but heat specifically denotes the increased blood circulation and metabolic activity in the affected tissues.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://radiationtherapyboard.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE