# REGISTERED NURSE CERTIFIED IN MATERNAL NEWBORN NURSING (RNC-MNN) PRACTICE TEST (SAMPLE)

## STUDY GUIDE



## SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

**VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE**

https://rnc-mnn.examzify.com

**Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.**

**ALL RIGHTS RESERVED.**

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

# Table of Contents


Copyright ..... 1

Table of Contents ..... 2

Introduction ..... 3

How to Use This Guide ..... 4

Questions ..... 5

Answers ..... 8

Explanations ..... 10

Next Steps ..... 16


SAMPLE

# Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

# How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

### 1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

### 2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

### 3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

### 4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

### 5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

### 6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

# Questions

SAMPLE

1. **If a patient with amniotic fluid embolism has dyspnea and pulmonary edema, which stage does this represent?**

   A. Stage I

   B. Stage II

   C. Stage III

   D. All of the above

2. **Which factor is least likely to increase risk for postpartum depression?**

   A. Prenatal anxiety

   B. Older maternal age

   C. Low socioeconomic status

   D. Unplanned pregnancy

3. **What condition is characterized by the failure to clear normal fetal lung fluid?**

   A. Transient tachypnea of the newborn

   B. Respiratory distress syndrome

   C. Cystitis

   D. Bacteriuria

4. **What assessment finding in a newborn might indicate an immediate need for intervention?**

   A. A heart rate below 100 beats per minute

   B. Skin that is pink and warm

   C. A weight above average for gestational age

   D. Normal respiratory rate for age

5. **What is the hallmark sign of a placental abruption?**

   A. Persistent back pain

   B. Sudden onset of vaginal bleeding accompanied by abdominal pain

   C. Light spotting without pain

   D. Excessive fetal movement

SAMPLE

**6. What is the primary concern if a postpartum woman presents with a boggy fundus?**

A. infection

B. postpartum hemorrhage

C. hormonal imbalance

D. retained placenta

**7. Which of the following is least likely to be a risk factor for postpartum hemorrhage?**

A. Prolonged labor

B. Induced or augmented labor

C. Prior vaginal birth

D. Difficult third stage of labor

**8. What is the minimum proficiency criteria defined in the clinical area?**

A. Ethical standards

B. Standard of care

C. Scope of practice

D. Moral standards

**9. Which pathology is least likely to cause persistent bradycardia in a newborn?**

A. Congenital heart block

B. Sepsis

C. Congestive heart failure

D. Increased intracranial pressure

**10. During the first week postpartum, the cervix generally narrows to what diameter?**

A. 1 cm

B. 2 cm

C. 0.5 cm

D. 3 cm

SAMPLE

# Answers

SAMPLE

1. A
2. B
3. A
4. A
5. B
6. B
7. C
8. B
9. C
10. A

SAMPLE

# Explanations

SAMPLE

## 1. If a patient with amniotic fluid embolism has dyspnea and pulmonary edema, which stage does this represent?

A. Stage I

B. Stage II

C. Stage III

D. All of the above

*In the context of amniotic fluid embolism (AFE), the condition is characterized by a progression of symptoms that reflect the severity of the situation. Stage I typically represents an early phase in which the patient may exhibit symptoms such as dyspnea or difficulty breathing, which is a common initial response to the infiltration of amniotic fluid into the maternal circulation. This phase may also include signs of hypoxemia and mild pulmonary edema, indicating the body is starting to react to the embolic event. As AFE progresses, additional clinical manifestations can develop, leading to worsening respiratory distress and even more severe pulmonary complications. Thus, identifying dyspnea and pulmonary edema as signs in the early stage of AFE aligns well with the characteristics described for Stage I. In more advanced stages, such as Stage II and Stage III, additional symptoms would likely be present, indicating a more severe systemic response. Understanding the progression of amniotic fluid embolism is crucial for recognizing the stage of the condition and providing timely intervention. The presence of dyspnea and pulmonary edema suggests that the situation is in its initial phase, making this correct answer significant in clinical practice for effective monitoring and management of the patient.*

## 2. Which factor is least likely to increase risk for postpartum depression?

A. Prenatal anxiety

B. Older maternal age

C. Low socioeconomic status

D. Unplanned pregnancy

*The factor of older maternal age being least likely to increase the risk for postpartum depression can be understood through various aspects of maternal health and psychosocial factors. Research indicates that older mothers, particularly those who are well-prepared and sought after their pregnancies, often display greater emotional stability and resilience. This demographic may have more life experience, support systems, and a sense of readiness for motherhood, which can contribute to better mental health outcomes postpartum. In contrast, prenatal anxiety, low socioeconomic status, and unplanned pregnancies are all associated with higher levels of stress and psychological distress. Prenatal anxiety can significantly impact mental health, leading to an increased likelihood of postpartum depression due to the ongoing emotional instability it can create. Low socioeconomic status often correlates with a lack of resources and support, resulting in heightened stress and challenges that can adversely affect a mother's mental health. Unplanned pregnancies may carry added emotional and logistical burdens that can also contribute to mental health challenges, as individuals may feel unprepared or overwhelmed by the sudden responsibility. Thus, the overall context indicates that older maternal age does not carry the same risk factors for postpartum depression as the other options, which are linked to heightened stress and anxiety levels surrounding pregnancy and motherhood.*

SAMPLE

## 3. What condition is characterized by the failure to clear normal fetal lung fluid?

A. Transient tachypnea of the newborn

B. Respiratory distress syndrome

C. Cystitis

D. Bacteriuria

*The condition characterized by the failure to clear normal fetal lung fluid is transient tachypnea of the newborn. This occurs when a newborn has difficulty expelling the fluid that normally fills the lungs during gestation. In most cases, this fluid is cleared during labor and immediately after birth. However, if the fluid is not adequately removed, the newborn may exhibit breathing difficulties, leading to a condition that resembles respiratory distress. Transient tachypnea typically presents within the first few hours after birth and is often seen in infants delivered via cesarean section or those who experienced a delayed clearance of lung fluid. While it usually resolves within 24 to 72 hours, it is essential for healthcare providers to monitor the infant's respiratory status during this time. The other conditions listed, like respiratory distress syndrome, cystitis, and bacteriuria, do not specifically relate to the failure to clear fetal lung fluid. Respiratory distress syndrome often involves complications due to surfactant deficiency, cystitis refers to bladder inflammation (not related to lung fluid), and bacteriuria pertains to the presence of bacteria in urine, which is unrelated to respiratory issues. Therefore, transient tachypnea of the newborn is the most accurate option regarding the described condition.*

## 4. What assessment finding in a newborn might indicate an immediate need for intervention?

A. A heart rate below 100 beats per minute

B. Skin that is pink and warm

C. A weight above average for gestational age

D. Normal respiratory rate for age

*A heart rate below 100 beats per minute in a newborn is a critical assessment finding that suggests the need for immediate intervention. In newborns, a normal heart rate typically ranges from 120 to 160 beats per minute. A heart rate less than 100 beats per minute can indicate bradycardia, which may compromise the newborn's oxygenation and lead to further complications if not addressed promptly. Immediate assessment and potentially resuscitative measures are warranted to ensure the newborn is stable and to prevent adverse outcomes such as hypoxia. In contrast, pink and warm skin indicates adequate perfusion and a stable thermal environment, while a weight above average for gestational age may require careful monitoring but does not inherently indicate an immediate need for intervention. A normal respiratory rate also suggests that the newborn is likely maintaining adequate ventilation and oxygenation. These findings, while important to note, do not present the same level of urgency as a significantly low heart rate does.*

SAMPLE

## 5. What is the hallmark sign of a placental abruption?

A. Persistent back pain

B. Sudden onset of vaginal bleeding accompanied by abdominal pain

C. Light spotting without pain

D. Excessive fetal movement

*The hallmark sign of a placental abruption is characterized by a sudden onset of vaginal bleeding that is often accompanied by abdominal pain. This condition occurs when the placenta detaches from the uterus before delivery, which can lead to significant maternal and fetal complications. The bleeding may be visible or concealed, and the abdominal pain results from the separation and subsequent irritation of the uterine wall. Vaginal bleeding in placental abruption is typically more severe than light spotting and usually indicates an emergency situation that may require immediate medical intervention. The abdominal pain can range from mild to severe and is often described as a constant cramping or sharp pain in the abdomen. This combination of symptoms highlights the seriousness of the condition and assists healthcare providers in making prompt decisions for the safety of both the mother and the fetus. Other options do not effectively capture the critical nature of placental abruption. For example, persistent back pain may not specifically indicate this condition and can be associated with various other pregnancy-related changes. Light spotting without pain is generally not concerning and is often not related to placental abruption, while excessive fetal movement is not indicative of abruption but rather can be a sign of fetal well-being in the absence of maternal distress. Thus, the distinct combination of sudden*

## 6. What is the primary concern if a postpartum woman presents with a boggy fundus?

A. infection

B. postpartum hemorrhage

C. hormonal imbalance

D. retained placenta

*A boggy fundus in a postpartum woman indicates that the uterus is not contracting effectively. This condition is often associated with postpartum hemorrhage, which is a critical and immediate concern following childbirth. The uterus should contract firmly following delivery to minimize bleeding. If the fundus is boggy and not firm, it suggests there may be excessive blood loss already occurring, making it imperative to address the situation quickly to prevent severe complications. While the other options may relate to postpartum conditions, they don't directly correlate with the immediate implications of a boggy fundus. For example, infection can lead to other signs and symptoms such as fever or foul-smelling lochia, and a hormonal imbalance could manifest later in the postpartum period without the immediate urgency that a boggy fundus presents. Retained placenta is a potential cause of a boggy fundus but is not the primary concern—monitoring for and managing postpartum hemorrhage takes precedence in this clinical scenario.*

SAMPLE

## 7. Which of the following is least likely to be a risk factor for postpartum hemorrhage?

A. Prolonged labor

B. Induced or augmented labor

C. Prior vaginal birth

D. Difficult third stage of labor

*Prior vaginal birth is considered the least likely risk factor for postpartum hemorrhage because women who have previously had a vaginal delivery generally have a better understanding of the birthing process, and their uterine tone may be stronger due to prior contractions. Additionally, they may have more effective healing processes in subsequent pregnancies and deliveries. In contrast, prolonged labor can increase the chances of uterine atony, which is a significant cause of postpartum hemorrhage. Induced or augmented labor may also lead to increased risk due to potential uterine overdistension or inadequate uterine tone if contractions are too strong or too frequent. Meanwhile, a difficult third stage of labor, characterized by challenges in delivering the placenta, can directly contribute to bleeding complications such as retained placental fragments or uterine atony, further elevating the risk of postpartum hemorrhage.*

## 8. What is the minimum proficiency criteria defined in the clinical area?

A. Ethical standards

B. Standard of care

C. Scope of practice

D. Moral standards

*The minimum proficiency criteria in the clinical area is best represented by the standard of care. This term refers to the level and type of care that healthcare professionals, including nurses, must provide to patients, ensuring it meets established benchmarks for quality and safety. The standard of care is derived from legal and professional guidelines and is fundamental in determining whether a healthcare provider's actions are appropriate in the context of patient treatment. It encompasses knowledge, skills, and practices essential for delivering effective care. The other options, while important in the context of healthcare, do not specifically define minimum proficiency criteria. Ethical standards focus on the moral principles guiding healthcare professionals, the scope of practice outlines the boundaries of what nurses are permitted to do, and moral standards pertain to personal values and principles of right and wrong. None of these directly establishes the baseline for clinical competency as the standard of care does.*

SAMPLE

## 9. Which pathology is least likely to cause persistent bradycardia in a newborn?

A. Congenital heart block

B. Sepsis

C. Congestive heart failure

D. Increased intracranial pressure

*Persistent bradycardia in a newborn can occur due to a variety of pathologies, and understanding the mechanisms behind these conditions is crucial for effective management. Congestive heart failure, while it can lead to a variety of cardiovascular issues, is more often associated with tachycardia rather than bradycardia. This is because when the heart is unable to effectively pump blood, compensatory mechanisms usually result in an increased heart rate to maintain adequate circulation. In contrast, congenital heart block is a direct cause of bradycardia since this condition affects the electrical conduction system of the heart. Sepsis can lead to bradycardia as well, particularly due to the systemic effects of infection and inflammatory responses. Increased intracranial pressure can cause bradycardia through mechanisms like the Cushing reflex, which results from pressure on the brain and subsequent changes in heart rate. Therefore, out of the listed conditions, congestive heart failure is least likely to be associated with persistent bradycardia in a newborn, making it the correct response in this context.*

## 10. During the first week postpartum, the cervix generally narrows to what diameter?

A. 1 cm

B. 2 cm

C. 0.5 cm

D. 3 cm

*During the first week postpartum, the cervix typically narrows to about 1 centimeter in diameter. This change is part of the body's natural process of recovery following childbirth. Immediately after delivery, the cervix remains significantly dilated, often ranging from 10 centimeters during active labor to wider. However, over the first few days and into the first week postpartum, it gradually begins to reduce in size due to the involution process, which is the body's way of returning to its pre-pregnancy state. The narrowing of the cervix is influenced by various factors, including hormonal changes, the physical healing that takes place after the trauma of delivering a baby, and uterine contractions that help reduce the size of the uterus and cervix. By the end of the first week, the cervix appears narrower and firmer, preparing for its roles in protection and fertility in future pregnancies. This physiological adaptation is critical for the postpartum recovery process and overall reproductive health.*

SAMPLE

# Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://rnc-mnn.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE