ROSH GASTROINTESTINAL PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://rosh-gastrointestinal.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. An elderly woman presents with difficulty swallowing after antibiotic treatment. What is the most likely causative agent for her symptoms?

A. Herpes simplex virus type I

B. Cytomegalovirus

C. Mycobacterium tuberculosis

D. Candida

2. Which characteristic is associated with Crohn disease?

A. Continuous uninterrupted inflammation of the colonic mucosa

B. Rectal involvement occurs in all cases

C. Smoking appears to confer a protective effect

D. Transmural involvement with cobblestone appearance

3. Which complication is associated with untreated pyloric stenosis?

A. Dehydration

B. Intestinal ischemia

C. Colonic perforation

D. Gastroesophageal reflux

4. In patients presenting with symptoms of gastroesophageal reflux disease (GERD), which weight-related factor should be considered?

A. Increased lean body mass

B. Obesity

C. Age

D. Physical activity

5. Which metabolic abnormality is most commonly associated with bulimia nervosa due to vomiting?

A. Hyperalbuminemia

B. Hypernatremia

C. Hypokalemia

D. Metabolic acidosis

SAMPLE

6. Following a cholecystectomy, what complication is often associated with gallbladder removal?

A. Biliary colic

B. Chronic diarrhea

C. Gallstones

D. Gallbladder cancer

7. In diagnosing rectal masses, which marker is most commonly elevated in patients with suspected rectal cancer?

A. Alpha fetoprotein

B. Carcinoembryonic antigen

C. Cancer antigen 125

D. Carcinoma antigen 15-3

8. Which demographic is most commonly affected by pyloric stenosis?

A. Infants

B. Adolescents

C. Adults

D. Elderly

9. Which combination of hepatitis B serologic markers indicates immunity due to vaccination?

A. HBsAg negative, anti-HBc negative, anti-HBs positive

B. HBsAg negative, anti-HBc positive, anti-HBs positive

C. HBsAg positive, anti-HBc positive, IgM anti-HBc negative, anti-HBs negative

D. HBsAg positive, anti-HBc positive, IgM anti-HBc positive, anti-HBs negative

10. What vitamin deficiency is most likely indicated in a woman with a diagnosis of osteomalacia?

A. Hypervitaminosis E

B. Hypervitaminosis K

C. Hypovitaminosis A

D. Hypovitaminosis D

SAMPLE

Answers

SAMPLE

1. D
2. D
3. A
4. B
5. C
6. B
7. B
8. A
9. A
10. D

SAMPLE

Explanations

SAMPLE

1. An elderly woman presents with difficulty swallowing after antibiotic treatment. What is the most likely causative agent for her symptoms?

A. Herpes simplex virus type I

B. Cytomegalovirus

C. Mycobacterium tuberculosis

D. Candida

In the scenario described, an elderly woman experiencing difficulty swallowing after antibiotic treatment suggests the development of esophagitis, commonly associated with Candida. Antibiotics can disrupt the normal flora of the mouth and gastrointestinal tract, leading to an overgrowth of Candida species, which are opportunistic fungi. Candida esophagitis typically presents with symptoms such as odynophagia (painful swallowing) and dysphagia (difficulty swallowing). This condition is especially prevalent in immunocompromised patients and those with altered microbial balance due to antibiotic use. The presentation aligns well with the effects of antibiotics leading to a proliferation of Candida, making it the most likely causative agent in this case. In contrast, the other pathogens are less common in this context. Herpes simplex virus type I usually causes oropharyngeal lesions and may lead to swallowing difficulties, but it is not as directly associated with antibiotic use. Cytomegalovirus can cause esophageal ulcers, particularly in immunocompromised individuals, but it is less frequently implicated immediately following antibiotic treatment. Mycobacterium tuberculosis is associated with extrapulmonary manifestations but is an unlikely cause of swallowing difficulty in this particular scenario.

2. Which characteristic is associated with Crohn disease?

A. Continuous uninterrupted inflammation of the colonic mucosa

B. Rectal involvement occurs in all cases

C. Smoking appears to confer a protective effect

D. Transmural involvement with cobblestone appearance

The characteristic associated with Crohn disease is transmural involvement with a cobblestone appearance. Crohn disease is known for its ability to affect any part of the gastrointestinal tract, from the mouth to the anus, and it typically involves all layers of the bowel wall, leading to transmural inflammation. This depth of inflammation can cause a variety of structural changes in the intestine, including the characteristic "cobblestone" appearance seen during endoscopy or imaging studies, where the mucosal surface shows irregularities and ulcerations. This feature is significant because it helps differentiate Crohn disease from other forms of inflammatory bowel disease, particularly ulcerative colitis, which primarily involves only the mucosal layer of the colon and typically presents with continuous inflammation. The cobblestone appearance is indicative of the focal nature of the disease, where segments of normal intestine may be interspersed with inflamed areas. In addition to the transmural involvement, Crohn disease can also lead to complications such as fistulas and strictures due to the extensive inflammation affecting the entire bowel wall. Understanding these characteristics is crucial for effective diagnosis and management of the disease.

SAMPLE

3. Which complication is associated with untreated pyloric stenosis?

A. Dehydration

B. Intestinal ischemia

C. Colonic perforation

D. Gastroesophageal reflux

In the context of untreated pyloric stenosis, dehydration is a well-recognized complication. Pyloric stenosis is a condition where the pylorus, the opening from the stomach into the small intestine, becomes narrowed, which obstructs the normal passage of food. As a result, infants with this condition frequently experience forceful vomiting after feeding, leading to significant fluid loss. Because of this ongoing vomiting and the inability to retain and absorb nutrients and fluids, dehydration can rapidly ensue. Dehydration in infants can manifest through various symptoms, including dry mouth, reduced urine output, lethargy, and decreased skin turgor. Left untreated, dehydration can lead to severe electrolyte imbalances and decrease the infant's overall health, making prompt recognition and treatment crucial. While intestinal ischemia, colonic perforation, and gastroesophageal reflux are serious medical issues, they are not directly associated with untreated pyloric stenosis in the same way that dehydration is. Intestinal ischemia is more commonly related to blood flow issues, colonic perforation is typically associated with conditions affecting the lower gastrointestinal tract, and gastroesophageal reflux while common, is not a direct result of pyloric stenosis. Therefore, the most direct complication arising from untreated pyl

4. In patients presenting with symptoms of gastroesophageal reflux disease (GERD), which weight-related factor should be considered?

A. Increased lean body mass

B. Obesity

C. Age

D. Physical activity

In patients presenting with symptoms of gastroesophageal reflux disease (GERD), obesity is a significant weight-related factor to consider. Obesity can lead to increased abdominal pressure, which in turn can promote the reflux of stomach contents into the esophagus. The accumulation of excess body fat, particularly around the abdomen, can exacerbate GERD symptoms by pushing the stomach upwards against the diaphragm, compromising the lower esophageal sphincter (LES) function, and increasing the likelihood of reflux episodes. Evaluating a patient's weight and overall body composition helps to strategize effective treatment plans and lifestyle modifications that can alleviate GERD symptoms, such as weight reduction, dietary changes, and increased physical activity tailored to the individual's health needs. Other factors mentioned, such as lean body mass, age, and physical activity, can influence health overall but do not have the same direct impact on GERD symptom severity as obesity does.

SAMPLE

5. Which metabolic abnormality is most commonly associated with bulimia nervosa due to vomiting?

A. Hyperalbuminemia

B. Hypernatremia

C. Hypokalemia

D. Metabolic acidosis

Hypokalemia is the metabolic abnormality most commonly associated with bulimia nervosa due to vomiting. This condition primarily arises from the loss of potassium that occurs when a person vomits repeatedly. During vomiting, significant amounts of gastric acid and electrolytes, including potassium, are expelled from the body. Potassium is an essential electrolyte that plays a critical role in various bodily functions, including muscle contraction and nerve transmission. When potassium levels fall significantly, it can lead to serious health issues, including cardiac complications, muscular weakness, and fatigue. Therefore, individuals with bulimia nervosa, particularly those engaging in self-induced vomiting, are at high risk of developing hypokalemia as a consequence of their eating disorder. In contrast, other metabolic abnormalities mentioned are less directly connected to the mechanism of vomiting seen in bulimia nervosa. Hyperalbuminemia and hypernatremia do not typically result from the act of vomiting in this context, while metabolic acidosis may occur in other scenarios, such as severe dehydration or renal impairment, but is not the most characteristic finding specifically linked to the vomiting seen in bulimia nervosa.

6. Following a cholecystectomy, what complication is often associated with gallbladder removal?

A. Biliary colic

B. Chronic diarrhea

C. Gallstones

D. Gallbladder cancer

After a cholecystectomy, which is the surgical removal of the gallbladder, chronic diarrhea can occur in some patients. This condition, often referred to as postcholecystectomy syndrome, can result from changes in bile flow and absorption after the gallbladder, which normally stores bile, has been removed. Without the gallbladder, bile is released continuously into the intestine, which can lead to fat malabsorption and potentially result in diarrhea. The other options are not typically associated complications directly resulting from gallbladder removal. Biliary colic usually occurs due to gallstones obstructing the bile ducts rather than after a cholecystectomy. Chronic diarrhea, while it can develop post-surgery, is less common. Gallstones can still form in the bile ducts after the gallbladder is removed, but this isn't a primary complication of the procedure itself. Gallbladder cancer is a distinct condition that is independent of the surgery performed and is not a complication associated with cholecystectomy.

SAMPLE

7. In diagnosing rectal masses, which marker is most commonly elevated in patients with suspected rectal cancer?

A. Alpha fetoprotein

B. Carcinoembryonic antigen

C. Cancer antigen 125

D. Carcinoma antigen 15-3

Carcinoembryonic antigen (CEA) is the most commonly elevated tumor marker in patients with suspected rectal cancer. CEA is a glycoprotein involved in cell adhesion, and it is produced during fetal development; however, its levels typically decrease after birth. In adults, elevated levels of CEA can indicate various types of cancers, particularly colorectal cancer, which includes rectal cancer. Testing for CEA is particularly useful in the context of colorectal cancer because it can assist in the diagnosis, provide prognostic information, and monitor treatment response or detect recurrence after primary treatment. In cases of rectal cancer, elevated CEA levels often correlate with tumor burden and can be used in conjunction with imaging studies and other diagnostic methods to assess the disease's extent. The other markers mentioned have different clinical applications. For instance, alpha-fetoprotein is primarily associated with liver cancer and germ cell tumors, while cancer antigen 125 is mainly utilized to track ovarian cancer. Carcinoma antigen 15-3 is primarily associated with breast cancer. Therefore, the elevation of CEA is specifically indicative of the pathological process associated with rectal and colorectal cancers, making it the most relevant choice in this context.

8. Which demographic is most commonly affected by pyloric stenosis?

A. Infants

B. Adolescents

C. Adults

D. Elderly

Pyloric stenosis is a condition that primarily affects infants, particularly those who are less than six months old. In this condition, the pylorus, which is the opening from the stomach into the small intestine, becomes narrowed, leading to severe vomiting and dehydration. The reasons for its higher incidence in infants may be related to developmental factors in utero or early in life, as it is not commonly found in older children or adults. The condition is typically diagnosed within the first few weeks after birth, and the majority of cases occur in male infants compared to females. Understanding this demographic is crucial for timely diagnosis and management, as untreated pyloric stenosis can lead to serious complications in this vulnerable age group. In contrast, pyloric stenosis is rarely seen in adolescents, adults, or the elderly, as these populations are outside the typical age range for this diagnosis. Conditions affecting the gastrointestinal tract in these older age groups usually present very differently and may involve other types of gastrointestinal pathology.

SAMPLE

9. Which combination of hepatitis B serologic markers indicates immunity due to vaccination?

A. HBsAg negative, anti-HBc negative, anti-HBs positive

B. HBsAg negative, anti-HBc positive, anti-HBs positive

C. HBsAg positive, anti-HBc positive, IgM anti-HBc negative, anti-HBs negative

D. HBsAg positive, anti-HBc positive, IgM anti-HBc positive, anti-HBs negative

The combination of serologic markers indicating immunity due to vaccination is characterized by the presence of anti-HBs (antibody to hepatitis B surface antigen) and the absence of HBsAg (hepatitis B surface antigen) and anti-HBc (antibody to hepatitis B core antigen). When a person has been vaccinated against hepatitis B, they develop specific antibodies to the surface antigen (anti-HBs), which signifies that the immune system has responded to the vaccine. In this correct scenario, the presence of anti-HBs suggests effective immunization, while the lack of HBsAg means there is no active infection. The absence of anti-HBc indicates that there has been no past or present infection, confirming that the immunity observed is entirely due to vaccination. This distinct pattern sets apart vaccinated individuals from those who have had a natural infection, which would show a positive anti-HBc. Understanding these markers is crucial for interpreting hepatitis B serology, especially in distinguishing between post-vaccination immunity and past or current infections.

10. What vitamin deficiency is most likely indicated in a woman with a diagnosis of osteomalacia?

A. Hypervitaminosis E

B. Hypervitaminosis K

C. Hypovitaminosis A

D. Hypovitaminosis D

The most likely vitamin deficiency indicated in a woman with a diagnosis of osteomalacia is a deficiency in vitamin D. Osteomalacia is characterized by the softening of bones due to a lack of mineralization, primarily caused by inadequate levels of vitamin D. This vitamin is essential for the absorption of calcium and phosphate, two critical minerals for maintaining bone health. A deficiency in vitamin D leads to decreased calcium absorption in the gut and subsequently can contribute to reduced bone mineralization, resulting in osteomalacia. Symptoms may include bone pain, muscle weakness, and an increased risk of fractures. While hypervitaminosis E, hypervitaminosis K, and hypovitaminosis A are related to other nutritional or metabolic disorders, they do not typically manifest as osteomalacia. Therefore, recognizing the importance of vitamin D in maintaining proper bone health is crucial in understanding the diagnosis of osteomalacia and the associated vitamin deficiencies.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://rosh-gastrointestinal.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE