SAEM EMERGENCY DEPARTMENT (ED) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://saemed.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What is a key indicator of potential ankle injury according to the Ottawa Ankle Rules?**

   A. Inability to walk 4 steps immediately and in the ED

   B. Tenderness over the lateral malleolus

   C. Swelling of the ankle

   D. Difficulty in wiggling toes

2. **What is the primary time-sensitive intervention in the 3-hour bundle for sepsis?**

   A. Administer blood culture

   B. Administer broad-spectrum antibiotics

   C. Measure blood pressure

   D. Check electrolyte levels

3. **What does the acronym "RACE" stand for in stroke recognition?**

   A. Rapid Assessment of Critical Emergencies

   B. Reassessment of Acute Care Environment

   C. Rapid Arterial oCclusion Evaluation

   D. Real-time Assessment of Cardiac Events

4. **What is the first step in the management of a patient with suspected sepsis?**

   A. Administering intravenous fluids

   B. Administering broad-spectrum antibiotics

   C. Performing a thorough history and physical examination

   D. Contacting a specialist

5. **What vital sign change is most indicative of shock?**

   A. Increased blood pressure

   B. Decreased respiratory rate

   C. Increased heart rate

   D. Elevation in temperature

SAMPLE

6. What percentage sensitivity can be achieved by having the patient sit upright for 10 minutes in cases of perforated viscus?

A. 10-30%

B. 30-80%

C. 80-100%

D. 0-10%

7. What does the term "decompensated shock" refer to?

A. A stage of shock with successful compensation

B. A stage of shock where organ function deteriorates

C. A reversible phase of hypovolemic shock

D. A stage of shock requiring surgical intervention

8. What is the first step in the preparation for intubation?

A. Preoxygenate the patient

B. Administer sedation

C. Gather intubation equipment

D. Check for contraindications

9. What is the most sensitive and specific test for diagnosing perforated viscus?

A. X-ray

B. CT scan

C. Ultrasound

D. MRI

10. What is the most important management step for inhalation injuries?

A. Immediate intubation and ventilation support

B. CXR and laryngoscopy

C. Administration of bronchodilators

D. Oral corticosteroids for inflammation

SAMPLE

Answers

SAMPLE

1. A
2. B
3. C
4. B
5. C
6. B
7. B
8. C
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is a key indicator of potential ankle injury according to the Ottawa Ankle Rules?

A. Inability to walk 4 steps immediately and in the ED

B. Tenderness over the lateral malleolus

C. Swelling of the ankle

D. Difficulty in wiggling toes

In the context of the Ottawa Ankle Rules, the key indicator of a potential ankle injury is the inability to bear weight. This rule helps to identify patients who are at risk for significant ankle and foot injuries and who may need further imaging to rule out fractures. The specific criterion states that a person is considered to have a potential fracture if they are unable to walk four steps both at the time of injury and during the evaluation in the emergency department. This aspect is crucial because it reflects a patient's functional capacity and pain level; if an individual is unable to walk even a short distance due to pain or instability, it suggests a higher likelihood of a serious injury, such as a fracture. The Ottawa Ankle Rules are designed to streamline the evaluation process, ensuring that only those who are at significant risk for fractures receive the necessary X-rays, thus minimizing unnecessary imaging for those who do not meet the criteria.

2. What is the primary time-sensitive intervention in the 3-hour bundle for sepsis?

A. Administer blood culture

B. Administer broad-spectrum antibiotics

C. Measure blood pressure

D. Check electrolyte levels

The primary time-sensitive intervention in the 3-hour bundle for sepsis is administering broad-spectrum antibiotics. Early initiation of antibiotics is crucial in managing sepsis because timely treatment directly correlates with improved patient outcomes. Sepsis can lead to rapid deterioration due to the body's overwhelming response to infection. The sooner broad-spectrum antibiotics are administered, the quicker the effective management of the underlying infection can occur, which significantly reduces morbidity and mortality associated with sepsis. While other actions like administering blood cultures, measuring blood pressure, and checking electrolyte levels are important components of sepsis management, they do not hold the same urgency as the initiation of antibiotic therapy. Blood cultures should be obtained before antibiotics are given to identify the causative organism, but this process can sometimes take longer than the critical window for antibiotic administration. Monitoring blood pressure and electrolyte levels are essential for ongoing assessment and management but do not serve as primary interventions that need to be executed within the first three hours. Therefore, administering broad-spectrum antibiotics stands out as the most immediate and critical step in the treatment protocol for sepsis.

SAMPLE

3. What does the acronym "RACE" stand for in stroke recognition?

A. Rapid Assessment of Critical Emergencies

B. Reassessment of Acute Care Environment

C. Rapid Arterial oCclusion Evaluation

D. Real-time Assessment of Cardiac Events

The acronym "RACE" stands for Rapid Arterial oCclusion Evaluation, which is a tool specifically designed to facilitate the quick identification of large vessel occlusions in patients with acute strokes. The importance of this acronym lies in its focus on improving the speed and accuracy of stroke assessment, which can significantly impact patient outcomes. Recognizing large vessel occlusions is critical because they are associated with more severe strokes and can often benefit from timely interventions, such as thrombectomy. In emergency settings, where time is of the essence, the RACE scale helps healthcare providers efficiently assess and triage patients, ultimately enhancing the overall quality of care and ensuring that patients receive the necessary treatment as quickly as possible. This is particularly vital in stroke management, where "time is brain," meaning that delays in treatment can lead to increased morbidity and mortality. The other options, while they might refer to relevant concepts in emergency medicine, do not pertain to the specific and established RACE framework used for stroke recognition.

4. What is the first step in the management of a patient with suspected sepsis?

A. Administering intravenous fluids

B. Administering broad-spectrum antibiotics

C. Performing a thorough history and physical examination

D. Contacting a specialist

The first step in the management of a patient with suspected sepsis involves administering broad-spectrum antibiotics. This is crucial because sepsis is a life-threatening condition that requires immediate intervention to combat high levels of infection and prevent progression to severe sepsis or septic shock. Starting antibiotics within the first hour of recognizing sepsis significantly improves patient outcomes, including reductions in mortality rates. In the context of sepsis, early identification and treatment are key. While intravenous fluids are also an important part of sepsis management to support blood pressure and organ perfusion, the priority is to control the underlying infection as quickly as possible. Furthermore, performing a thorough history and physical examination is important for diagnosis, but it should not delay the administration of antibiotics. Contacting a specialist can also play a role in sepsis management, especially in complex cases, but immediate antibiotic therapy is paramount. This timely intervention is supported by extensive clinical guidelines, emphasizing the urgency of addressing infection in sepsis cases.

SAMPLE

5. What vital sign change is most indicative of shock?

A. Increased blood pressure

B. Decreased respiratory rate

C. Increased heart rate

D. Elevation in temperature

In the context of shock, an increased heart rate is a critical vital sign change that indicates the body is attempting to compensate for diminished blood flow or oxygen delivery to tissues. When shock occurs, the body recognizes a deficit in perfusion and often responds by elevating the heart rate, known as tachycardia. This compensatory mechanism helps to maintain blood pressure and ensure that vital organs receive sufficient blood supply. Increased heart rate can be seen in various types of shock, including hypovolemic, cardiogenic, and septic shock, as the autonomic nervous system triggers the heart to pump faster in response to low blood volume or systemic vasodilation. Monitoring the heart rate provides essential information about a patient's hemodynamic status and can prompt further evaluation and intervention. In contrast, elevated blood pressure is not typically associated with shock, as it often decreases or remains normal in early stages of shock due to compensatory mechanisms failing over time. Decreased respiratory rate does not signify shock and may actually indicate deterioration. Lastly, an elevated temperature can indicate an infectious cause of shock but is not a direct indicator of shock itself. Thus, increased heart rate serves as a reliable indicator of the body's response to shock conditions.

6. What percentage sensitivity can be achieved by having the patient sit upright for 10 minutes in cases of perforated viscus?

A. 10-30%

B. 30-80%

C. 80-100%

D. 0-10%

Having the patient sit upright for 10 minutes can significantly enhance the sensitivity of detecting free air in cases of perforated viscus. The rationale for this practice involves the dynamics of air movement within the abdominal cavity. When a perforation occurs, air can escape from the gastrointestinal tract into the peritoneal cavity. By positioning the patient upright, the free air rises to the highest point in the peritoneal cavity, typically underneath the diaphragm, making it more visible on imaging studies like X-rays or CT scans. The sensitivity range of 30-80% indicates that while the technique improves detection, it is not foolproof. Factors such as the amount of free air present, the patient's body position before assessing, and the quality of imaging can all influence the final outcome. This percentage reflects that when patients are upright, there's a notable but not complete enhancement in the ability to identify free air associated with perforated organs.

SAMPLE

7. What does the term "decompensated shock" refer to?

A. A stage of shock with successful compensation

B. A stage of shock where organ function deteriorates

C. A reversible phase of hypovolemic shock

D. A stage of shock requiring surgical intervention

The term "decompensated shock" specifically refers to a stage where the body can no longer maintain adequate blood circulation and perfusion to vital organs, leading to a deterioration in organ function. At this point, compensatory mechanisms that initially helped to stabilize blood pressure and blood flow are overwhelmed. As a result, patients may experience progressive organ failure, drop in blood pressure, increased heart rate, and altered mental status. Identifying this stage is critical in emergency medicine, as it indicates the need for immediate intervention to restore perfusion and prevent irreversible damage to organs. The other options describe different stages or types of shock, such as successful compensation or a reversible phase, which do not accurately capture the essence of decompensated shock. In decompensated shock, the body's attempts to compensate have failed, necessitating urgent medical attention.

8. What is the first step in the preparation for intubation?

A. Preoxygenate the patient

B. Administer sedation

C. Gather intubation equipment

D. Check for contraindications

The initial step in preparing for intubation is to gather intubation equipment. This is crucial because having all necessary tools readily available ensures that the intubation process can proceed smoothly and efficiently. Key items include the endotracheal tubes, laryngoscope, suction device, bag-valve mask, and any adjuncts that might be needed to secure an airway. By organizing the equipment beforehand, the healthcare provider can minimize the risk of delays during the procedure, which can be critical in emergency situations where time is of the essence. Additionally, knowing that all the required items are at hand allows the clinician to focus on the patient during the intubation process, ensuring both safety and adherence to best practices. Following the gathering of equipment, subsequent steps like preoxygenation, administering sedation, and checking for contraindications can be performed. Each of those steps is important, but they are contingent on being prepared with the appropriate tools for the intubation itself.

SAMPLE

9. What is the most sensitive and specific test for diagnosing perforated viscus?

A. X-ray

B. CT scan

C. Ultrasound

D. MRI

A CT scan is indeed the most sensitive and specific test for diagnosing a perforated viscus. This imaging modality allows for detailed visualization of the abdominal and thoracic cavities, enabling the identification of free air, which indicates perforation in the gastrointestinal tract. The cross-sectional images produced by a CT scan provide clear information about the location and extent of any perforations, as well as associated complications such as abscess formation or peritonitis, which are crucial for guiding management. In contrast, while X-rays can show free air under the diaphragm, they are less sensitive and specific compared to CT scans, potentially missing smaller perforations. Ultrasound can detect fluid and can be useful in certain clinical scenarios, but it may not provide the same level of detail as a CT scan and can be operator-dependent. MRI is generally not used in the acute setting for diagnosing perforated viscus due to time constraints and its limited availability in emergency departments. Therefore, the superior diagnostic capability of CT scans makes them the preferred choice for this clinical scenario.

10. What is the most important management step for inhalation injuries?

A. Immediate intubation and ventilation support

B. CXR and laryngoscopy

C. Administration of bronchodilators

D. Oral corticosteroids for inflammation

For inhalation injuries, the pivotal management step is a thorough assessment of the airway, which is best accomplished through laryngoscopy. This procedure allows the clinician to directly visualize the airway, assessing for any edema, burns, or obstruction that could compromise breathing. Not all patients with inhalation injuries present with immediate respiratory distress; however, airway swelling can develop rapidly, and early intervention can be critical. While immediate intubation and ventilation support is crucial in cases where the airway is compromised or breathing is severely impaired, it may not always be necessary at the outset for every inhalation injury patient. Laryngoscopy provides a clearer understanding of the situation and allows for timely decisions regarding whether intubation is necessary. The administration of bronchodilators may help alleviate bronchospasm, but it does not address the potential upper airway obstruction that inhalation injuries may cause. Similarly, oral corticosteroids can reduce inflammation over time but should not be the primary focus in the immediate management of inhalation injuries. The initial step must focus on assessing the airway to ensure it is patent and safe for ventilation.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://saemed.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE