
ST. JOHN AMBULANCE FIRST AID PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://stjohnambulancefirstaid.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What should you do for a person that has fainted?

A. Keep them upright

B. Lay them down and elevate their legs

C. Give them sugary snacks

D. Sprinkle water on their face

2. What is a critical first step when dealing with a suspected fracture?

A. Move the injured limb

B. Immobilize the area

C. Apply ice immediately

D. Check for pulse

3. What could be a sign of a concussion during a primary survey?

A. Pain in the stomach

B. Nausea or vomiting

C. Rash on the skin

D. Difficulty breathing

4. What is a common sign of internal bleeding?

A. Rapid heartbeat

B. Skin rash

C. Headache

D. Fever

5. Which of the following does NOT relate to the assessment of an unconscious person?

A. Checking response to pain

B. Checking for smell of alcohol

C. Assessing if they are alert

D. Speaking to them to check for voice response

6. What is the first action to take when treating a serious burn?

A. Cover the burn with a cloth

B. Cool the burn under running water for at least 10 minutes

C. Apply ice directly to the burn

D. Wrap the burn with a bandage immediately

SAMPLE

7. Which of the following symptoms may suggest a serious medical condition?

A. Thirst

B. Thready pulse

C. Dry skin

D. Increased energy levels

8. What is the emergency action step for handling burns?

A. Cover the burn with ice immediately

B. Cool the burn with running water and cover it with a sterile dressing

C. Apply butter or ointment to the burn

D. Leave the burn uncovered for air exposure

9. Which piece of information is crucial to communicate to emergency responders?

A. The patient's medical history only

B. The condition of the patient and treatments administered

C. The weather conditions at the time of injury

D. The location of the incident solely

10. Which of the following is a sign of stroke?

A. Severe headache and fever

B. Sudden numbness or weakness in the face, arm, or leg

C. Loss of balance and coordination only

D. Slow, slurred speech and mild headache

SAMPLE

Answers

SAMPLE

1. B
2. B
3. B
4. A
5. B
6. B
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What should you do for a person that has fainted?

A. Keep them upright

B. Lay them down and elevate their legs

C. Give them sugary snacks

D. Sprinkle water on their face

When dealing with a person who has fainted, laying them down and elevating their legs is the appropriate response. This position assists in improving blood flow to the brain, as it helps return blood from the lower extremities to the upper body. Elevating the legs promotes proper circulation and can help the person regain consciousness more quickly. In situations of fainting, the goal is to reduce any potential injury from falling and stabilize the individual's condition until they recover. This method not only aids in physical recovery but also ensures that the person is in a more comfortable position where they can be monitored effectively. In contrast, keeping someone upright can exacerbate the issue, as gravitational forces may further decrease blood flow to the brain. Feeding sugary snacks is not advisable right after fainting because the person may have difficulty swallowing or could choke. Sprinkling water on their face does not address the underlying issue of reduced blood flow and may even be counterproductive by causing discomfort.

2. What is a critical first step when dealing with a suspected fracture?

A. Move the injured limb

B. Immobilize the area

C. Apply ice immediately

D. Check for pulse

The critical first step when dealing with a suspected fracture is to immobilize the area. This is essential because moving the injured limb can exacerbate the injury, potentially causing further damage to the bone, surrounding muscles, nerves, or blood vessels. Immobilization helps to minimize pain and reduces the risk of complications. By stabilizing the area, you reduce movement and prevent any shifting of the fractured bone ends, which can lead to more severe injuries and delays in healing. It also prepares the injured person for transportation to a medical facility if necessary. Applying ice, while beneficial for swelling and pain relief, should not take precedence over immobilization. Checking for a pulse is important in assessing circulation but is not the first step in managing a suspected fracture.

3. What could be a sign of a concussion during a primary survey?

A. Pain in the stomach

B. Nausea or vomiting

C. Rash on the skin

D. Difficulty breathing

Nausea or vomiting is indeed a significant sign of a concussion that can be observed during a primary survey. When an individual experiences a concussion, which is a type of traumatic brain injury, symptoms may manifest in various ways, including gastrointestinal upset. The brain's response to the trauma can affect areas that control nausea and vomiting. In a primary survey, where the focus is on identifying life-threatening conditions, the presence of nausea or vomiting can indicate that the individual may have sustained a head injury and is experiencing cerebral effects. This symptom, alongside others such as confusion, headache, or balance issues, helps in assessing the extent of the injury and the need for immediate medical attention. While other options like pain in the stomach, a rash on the skin, or difficulty breathing can indicate other medical concerns, they are not specifically associated with concussion. Understanding the relationship between nausea and concussions assists first aid providers in recognizing the need for further evaluation and potential transport to a medical facility for comprehensive care.

SAMPLE

4. What is a common sign of internal bleeding?

A. Rapid heartbeat

B. Skin rash

C. Headache

D. Fever

A rapid heartbeat is a common sign of internal bleeding because it indicates that the body is responding to a potential loss of blood volume. When internal bleeding occurs, the body may lose significant amounts of blood, leading to decreased blood pressure. In an attempt to maintain adequate blood flow to vital organs, the heart compensates by beating faster. This physiological response is known as compensatory tachycardia, and it helps sustain blood circulation even when the overall blood volume is reduced. The other options, while they may indicate other health issues, do not typically correlate with internal bleeding. A skin rash might suggest an allergic reaction or skin condition, a headache could be caused by numerous factors unrelated to internal bleeding, and a fever usually indicates an infection rather than a bleeding scenario. Hence, recognizing a rapid heartbeat as a symptom can be critical for prompt medical intervention when internal bleeding is suspected.

5. Which of the following does NOT relate to the assessment of an unconscious person?

A. Checking response to pain

B. Checking for smell of alcohol

C. Assessing if they are alert

D. Speaking to them to check for voice response

When assessing an unconscious person, the primary focus is on evaluating their responsiveness and vital signs to determine the level of consciousness and any immediate medical needs. The correct response highlights an approach that does not align directly with assessing conscious awareness. Checking response to pain, assessing alertness, and speaking to the person to check for voice response all involve interacting with the individual to gauge their level of consciousness or responsiveness. Each of these methods helps first responders understand whether the person is conscious, semi-conscious, or fully unconscious, which is crucial for further treatment decisions. In contrast, checking for the smell of alcohol does not directly inform you about the level of consciousness of the person. While it might be relevant in a broader context of understanding potential causes of unconsciousness (such as intoxication), it is not a direct assessment technique for evaluating the state of awareness or responsiveness of an unconscious individual. Therefore, this option is not relevant to the primary assessment criteria used in determining consciousness in first aid scenarios.

6. What is the first action to take when treating a serious burn?

A. Cover the burn with a cloth

B. Cool the burn under running water for at least 10 minutes

C. Apply ice directly to the burn

D. Wrap the burn with a bandage immediately

Cooling the burn under running water for at least 10 minutes is critical because it helps to reduce the temperature of the affected area, alleviating pain and minimizing further tissue damage. This action also helps to prevent the burn from progressing into deeper tissue layers. Cooling the burn effectively halts the burning process by washing away any residual heat and can also reduce swelling. In the context of burn treatment, it's important to avoid applying ice directly to the burn, as this can cause further damage to the skin and tissue. Covering the burn with a cloth too soon can trap heat, and immediately wrapping it with a bandage may hinder the cooling process and lead to complications. Therefore, the initial step should focus on cooling the burn properly to ensure optimal care.

SAMPLE

7. Which of the following symptoms may suggest a serious medical condition?

A. Thirst

B. Thready pulse

C. Dry skin

D. Increased energy levels

A thready pulse is a significant clinical sign that may indicate a serious medical condition. It is characterized by a weak, thin pulse that is difficult to feel or is barely perceptible. This may signal various underlying issues, such as shock, severe dehydration, or significant blood loss, all of which require immediate medical attention. The presence of a thready pulse might reflect the body's response to compromised circulation or inadequate blood oxygenation, emphasizing its potential severity. In contrast, the other symptoms listed do not inherently suggest a serious medical condition. Thirst might indicate dehydration but is not solely indicative of a serious issue without additional context. Dry skin can also arise from mild conditions or environmental factors rather than pointing directly to a medical emergency. Increased energy levels are typically a positive sign and would generally not suggest a serious medical concern. Therefore, the thready pulse stands out as a critical symptom that necessitates a higher level of vigilance and possible intervention.

8. What is the emergency action step for handling burns?

A. Cover the burn with ice immediately

B. Cool the burn with running water and cover it with a sterile dressing

C. Apply butter or ointment to the burn

D. Leave the burn uncovered for air exposure

The recommended emergency action step for handling burns involves cooling the burn with running water and then covering it with a sterile dressing. This method is crucial for several reasons. First, cooling the burn helps to reduce pain and prevents further tissue damage by lowering the skin temperature. Running water, ideally at room temperature, should be used for at least 10-20 minutes. This is more effective than applying ice, which can cause further harm and frostbite to the already damaged tissue. After cooling the burn, covering it with a sterile dressing helps protect the area from infection, reduces pain by keeping air off the wound, and facilitates healing. A non-stick dressing is preferred, as it minimizes trauma when changing the dressing. In contrast, using ice can lead to additional injury to the skin, while applying butter or ointments can trap heat and lead to infection. Leaving a burn uncovered exposes it to germs, increasing the risk of infection and complicating the healing process. Ensuring proper care in this way is essential for promoting recovery and minimizing complications from burns.

SAMPLE

9. Which piece of information is crucial to communicate to emergency responders?

A. The patient's medical history only

B. The condition of the patient and treatments administered

C. The weather conditions at the time of injury

D. The location of the incident solely

Communicating the condition of the patient and the treatments that have been administered is crucial for several reasons. Emergency responders rely on this information to assess the current status of the patient and to determine the most appropriate course of action upon their arrival. Understanding the patient's condition can help responders prioritize interventions based on the severity of the situation, whether the patient is stable, unstable, or requires immediate resuscitation efforts. Furthermore, knowing what treatments have already been given ensures that responders do not duplicate efforts or inadvertently administer something that could worsen the patient's condition. While the patient's medical history is important, it often takes a backseat to the immediate assessments of current symptoms and any interventions taken prior to arrival. Weather conditions might provide context but do not typically impact the medical treatment provided to the patient. Similarly, while knowing the location of the incident is essential for responders to reach the scene quickly, it is the patient's current condition and prior treatment that directly informs the life-saving measures they will implement.

10. Which of the following is a sign of stroke?

A. Severe headache and fever

B. Sudden numbness or weakness in the face, arm, or leg

C. Loss of balance and coordination only

D. Slow, slurred speech and mild headache

Sudden numbness or weakness in the face, arm, or leg is recognized as a classic sign of a stroke, particularly when it affects one side of the body. This symptom occurs due to the sudden loss of blood flow to parts of the brain, leading to damage. Recognizing this symptom quickly is crucial because immediate medical attention can significantly improve outcomes for people experiencing a stroke. While other choices may include symptoms that could appear with various medical conditions, they do not align as directly with the typical presentations associated with a stroke. For instance, severe headache and fever may indicate other medical issues, such as an infection or migraine, rather than a stroke. Loss of balance and coordination alone could also result from numerous conditions, not specifically indicating a stroke. Slow, slurred speech combined with a mild headache could suggest other neurological issues but does not capture the urgency or specificity of the numbness or weakness that is characteristic of a stroke. Therefore, identifying sudden numbness or weakness is essential for effective stroke recognition and timely intervention.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://stjohnambulancefirstaid.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE