STEPP GI DISORDERS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://steppgidisorders.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What is a recommended action if a patient feels choking or pressure during ERCP?**

   A. Ask to stop the procedure

   B. Remain calm and allow the procedure to continue

   C. Immediately leave the examination room

   D. Raise a hand and signal the nurse

2. **Which medication is typically used for sedation during a colonoscopy?**

   A. Atropine

   B. Midazolam

   C. Ibuprofen

   D. Lidocaine

3. **What condition can computed tomography (CT) scans primarily identify?**

   A. Bone fractures

   B. Abnormalities or tumors

   C. Blood clotting issues

   D. Respiratory infections

4. **What complication is most commonly associated with gastric surgeries?**

   A. Infection

   B. Dehydration

   C. Excessive weight gain

   D. Hemorrhage

5. **What is the required dietary preparation for a cholecystogram?**

   A. Fat-free supper the night before

   B. High-carb breakfast

   C. Fasting for 24 hours

   D. No specific dietary restrictions

SAMPLE

6. What vital sign change may indicate hypovolemic shock due to complications after a paracentesis?

A. Increased blood pressure

B. Increased heart rate

C. Increased respiratory rate

D. Cold extremities

7. What is a common complication of chronic pancreatitis?

A. Chronic kidney disease

B. Diabetes mellitus

C. Hyperlipidemia

D. Diverticulitis

8. When should an ostomy bag be emptied?

A. When full of stool

B. When 1/3-1/2 full of stool or gas

C. Once a day, regardless of amount

D. When it begins leaking

9. Which of the following is a nursing consideration for barium tests?

A. Encourage chewing gum before the procedure

B. No bowel preparation if contraindications exist

C. Low residue diet and increase fluids

D. Normal diet before the test

10. Before a Small Bowel Capsule Endoscopy, what is instructive about the patient's activity level?

A. Complete bed rest is required

B. Normal activities are allowed

C. Limited movement is advisable

D. Heavy exercise should be avoided

SAMPLE

Answers

SAMPLE

1. B
2. B
3. B
4. B
5. A
6. B
7. B
8. B
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. What is a recommended action if a patient feels choking or pressure during ERCP?

A. Ask to stop the procedure

B. Remain calm and allow the procedure to continue

C. Immediately leave the examination room

D. Raise a hand and signal the nurse

In the context of an Endoscopic Retrograde Cholangiopancreatography (ERCP) procedure, if a patient experiences sensations of choking or pressure, it is crucial for them to remain calm and allow the procedure to continue. This recommendation is based on the understanding that some degree of discomfort is common during ERCP, as the procedure involves the manipulation of tools within the esophagus and duodenum, which can trigger such sensations. While it is essential for the patient to communicate any feelings of distress to the medical team, maintaining calmness allows the healthcare providers to assess the situation and proceed safely. Stopping the procedure abruptly could lead to complications or incomplete diagnostic evaluations. The medical team is trained to manage discomfort and can adjust their approach or provide assistance while ensuring the procedure's integrity and the patient's safety. The other options do not effectively support the procedural flow and the safety of the patient. Asking to stop the procedure might not be the best immediate action, and leaving the examination room could lead to further complications. Raising a hand to signal the nurse can be helpful, but it might not be as effective as the patient focusing on staying calm to help the team manage the situation while they continue the necessary interventions.

2. Which medication is typically used for sedation during a colonoscopy?

A. Atropine

B. Midazolam

C. Ibuprofen

D. Lidocaine

Midazolam is typically used for sedation during a colonoscopy because it is a short-acting benzodiazepine. It provides both sedative and anxiolytic effects, making it an ideal choice for procedures that may cause anxiety or discomfort to the patient. Midazolam induces a state of relaxation and can also produce amnesia regarding the procedure, which can help alleviate patient anxiety and improve the overall experience. Other options, such as atropine, are used for different purposes, such as reducing secretions or counteracting bradycardia. Ibuprofen is a nonsteroidal anti-inflammatory drug useful for pain relief but does not serve a sedative purpose. Lidocaine, while used for local anesthesia, does not provide the systemic sedation that is often required during colonoscopy procedures.

SAMPLE

3. What condition can computed tomography (CT) scans primarily identify?

A. Bone fractures

B. Abnormalities or tumors

C. Blood clotting issues

D. Respiratory infections

Computed tomography (CT) scans are particularly effective for identifying abnormalities or tumors within the body due to their ability to provide detailed cross-sectional images of internal structures. This imaging technique enhances visualization of soft tissues, organs, and masses, enabling healthcare providers to detect and assess the presence of tumors, cysts, and other pathological changes in a way that traditional X-rays may not. CT scans are adept at demonstrating changes in tissue density, which is crucial for diagnosing malignancies and determining their size and extent. This level of detail is especially important in oncology, where early detection can significantly impact treatment options and outcomes. Bone fractures, although visible on CT scans, are not the primary focus of this imaging method; other modalities like X-rays are more commonly used for that purpose. Likewise, while CT scans can help identify some complications related to blood clotting, such as in cases of pulmonary embolism, they are not primarily designed for evaluating clotting issues. As for respiratory infections, while CT can be used to assess complications of infections in the lungs, such as abscesses or pneumonia, the primary diagnosis of respiratory infections is usually made through other methods like chest X-rays and clinical evaluation. This specificity and precision in detecting tumors and other abnormalities underscore the role

4. What complication is most commonly associated with gastric surgeries?

A. Infection

B. Dehydration

C. Excessive weight gain

D. Hemorrhage

The most commonly associated complication with gastric surgeries is dehydration. Following such procedures, patients often experience significant alterations in their digestive system, which can lead to difficulties in nutrient absorption and fluid intake. Additionally, surgical alterations, especially those involving the stomach or intestines, may result in reduced capacity to retain fluids, further exacerbating the risk of dehydration. Patients may also develop new eating habits, such as decreased food intake or changes in their dietary choices, which can contribute to insufficient fluid consumption. Since dehydration can lead to serious health consequences, it is critical for healthcare providers to monitor patients post-operatively and encourage adequate hydration. While infection, excessive weight gain, and hemorrhage can be complications of gastric surgeries, they are not as consistently observed as dehydration in the immediate aftermath of the procedure. Each of these issues can arise under specific circumstances; however, dehydration remains a more prevalent concern in the early recovery phases due to the physiological changes that occur after gastric interventions.

SAMPLE

5. What is the required dietary preparation for a cholecystogram?

A. Fat-free supper the night before

B. High-carb breakfast

C. Fasting for 24 hours

D. No specific dietary restrictions

The required dietary preparation for a cholecystogram involves consuming a fat-free supper the night before the procedure. This preparation is crucial because the cholecystogram relies on the gallbladder's ability to concentrate a contrast material used during imaging. A fat-free meal ensures that the gallbladder is emptied of any residual bile and is in a state ready to absorb the contrast material effectively. This enhances visualization of the gallbladder and surrounding structures during the imaging process, allowing for better diagnostic clarity. In contrast, the other options do not provide the specific preparation needed for optimal results. A high-carb breakfast might lead to gallbladder contraction, which is not desirable before the procedure. Fasting for 24 hours is excessive and unnecessary for a cholecystogram, as it can lead to patient discomfort without improving the outcome. Lastly, having no specific dietary restrictions would not adequately prepare the gallbladder for imaging, potentially leading to suboptimal results. Thus, adhering to a fat-free dinner allows for the best preparation for the test.

6. What vital sign change may indicate hypovolemic shock due to complications after a paracentesis?

A. Increased blood pressure

B. Increased heart rate

C. Increased respiratory rate

D. Cold extremities

In the context of hypovolemic shock, particularly after a procedure such as a paracentesis, increased heart rate is a crucial sign to monitor. Hypovolemic shock occurs when there is a significant loss of blood volume, which can lead to reduced cardiac output and inadequate perfusion of tissues. As the body attempts to compensate for the decreased blood volume and maintain perfusion to vital organs, the heart compensates by increasing its rate of contraction, resulting in tachycardia. This compensatory mechanism occurs because the body releases catecholamines, which stimulate the heart to beat faster in an effort to maintain adequate blood flow and oxygen delivery despite the drop in circulating blood volume. Therefore, when monitoring a patient who has undergone a paracentesis, an increased heart rate indicates that the body is responding to potential fluid loss and is a warning sign of possible hypovolemic shock. While other vital signs may also change in the context of hypovolemic shock, such as increased respiratory rate or changes in temperature, the heightened heart rate is particularly significant as an initial response to significant volume loss. It's important to assess heart rate alongside other clinical indicators to gauge the overall condition of the patient.

SAMPLE

7. What is a common complication of chronic pancreatitis?

A. Chronic kidney disease

B. Diabetes mellitus

C. Hyperlipidemia

D. Diverticulitis

Chronic pancreatitis often leads to the destruction of pancreatic tissue and a decline in the organ's ability to produce essential hormones and enzymes. One significant complication arising from this condition is diabetes mellitus. The pancreas plays a crucial role in regulating blood sugar levels by producing insulin; when it becomes damaged from chronic inflammation, insulin production is compromised. This leads to insulin deficiency and subsequently diabetes mellitus, which can be a direct result of prolonged pancreatic damage. While chronic pancreatitis can have other implications such as malabsorption due to lack of digestive enzymes or potential complications related to nutrient deficiencies, the development of diabetes mellitus is particularly notable as it reflects a direct failure of the pancreas to perform its endocrine function.

8. When should an ostomy bag be emptied?

A. When full of stool

B. When 1/3-1/2 full of stool or gas

C. Once a day, regardless of amount

D. When it begins leaking

An ostomy bag should ideally be emptied when it is 1/3 to 1/2 full of stool or gas. This recommendation is based on several important factors. First, an ostomy bag that is too full can become heavy and may put strain on the adhesive seal, increasing the risk of leakage or detachment from the skin. Emptying the bag at this level helps maintain a secure fit, promoting better overall hygiene and comfort for the individual. Moreover, frequent emptying at this level can help manage odor and make the process of cleaning and changing the bag easier. It allows patients to be proactive in their care and limits the potential for accidents or irritations that can occur if the bag is left too full for extended periods. Additionally, monitoring the contents aids in assessing the health of the individual's digestive system; any significant changes in consistency or volume can signal the need for medical consultation. By emptying the bag at this recommended level, individuals with an ostomy can maintain better quality of life and prevent complications associated with improper ostomy care.

9. Which of the following is a nursing consideration for barium tests?

A. Encourage chewing gum before the procedure

B. No bowel preparation if contraindications exist

C. Low residue diet and increase fluids

D. Normal diet before the test

For a barium test, nursing considerations play a crucial role in ensuring that the procedure is effective and safe for the patient. The correct answer emphasizes the importance of adopting a low residue diet and increasing fluid intake before the test. A low residue diet is recommended because it minimizes the amount of undigested food in the intestines, allowing for clearer imaging during the barium examination. This type of diet helps reduce the amount of stool present, which can obscure imaging results. Additionally, increasing fluid intake is crucial as it helps to flush out the barium from the system after the test is complete, preventing constipation and ensuring the patient's comfort following the procedure. The other options do not align with best practices for preparing a patient for a barium test. Chewing gum before the procedure could potentially stimulate digestive processes that might interfere with the test results. Skipping bowel preparation when there are contraindications may lead to inadequate visualization during the test. Allowing a normal diet before the test can lead to excess residue in the bowel, making it difficult to obtain accurate images. Thus, the focus on a low residue diet and increased hydration is essential for maximizing the diagnostic quality of the barium test.

SAMPLE

10. Before a Small Bowel Capsule Endoscopy, what is instructive about the patient's activity level?

A. Complete bed rest is required

B. Normal activities are allowed

C. Limited movement is advisable

D. Heavy exercise should be avoided

The correct choice emphasizes that normal activities are allowed prior to a Small Bowel Capsule Endoscopy. Patients can typically maintain their usual routines without the need for significant restrictions on movement. This is crucial because the procedure involves swallowing a capsule that contains a camera, which then travels through the gastrointestinal tract, capturing images. Regular activity does not interfere with the capsule's passage through the intestines, and it also helps patients remain comfortable and less anxious about the procedure. In general, while some guidelines may suggest avoiding heavy exercise or vigorous activities just to ensure patient comfort and minimize any potential disruptions, normal daily activities should not pose a risk to the success of the endoscopy. Thus, allowing patients to engage in their typical activities is both practical and supportive of their well-being leading up to the exam.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://steppgidisorders.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE