
SWIFT RIVER PEDIATRICS PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://swiftriverpediatrics.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. In which category is Ampicillin classified?

A. Antihistamine

B. Antibiotic

C. Pain reliever

D. Vitamin and Minerals

2. Which action should be performed if chest tube drainage is observed to change in color?

A. Notify the physician immediately

B. Continue to monitor the drainage for further changes

C. Clamp the chest tube

D. Document the change and wait for further instructions

3. What is the typical age for the onset of puberty in girls?

A. 6-10 years

B. 8-13 years

C. 10-15 years

D. 12-16 years

4. Which form of Diphenhydramine is listed for administration?

A. 25 mg tablets

B. 50 mg tablets

C. Liquid suspension

D. Capsules

5. When should school-age children have their first dental exam?

A. By age 1

B. By age 3

C. By age 5

D. By age 7

6. Why is Bridgett's weight monitored closely during her recovery?

A. To determine medication side effects

B. To assess nutritional needs

C. To ensure it aligns with her baseline

D. To prepare for future surgeries

SAMPLE

7. What is a classic sign of intussusception in a child?

A. Severe abdominal pain

B. "Currant jelly" stool

C. Yellow-green vomiting

D. Diarrhea

8. What should be requested for administering a 1000 mL bag of 0.9% Normal Saline?

A. Request an LPN

B. Request an RN

C. Handle it independently

D. Request a CNA

9. What should the nurse document after administering digoxin?

A. Only the dosage given

B. Patient's heart rate and any rhythm abnormalities

C. Patient's weight and vital signs

D. Time of medication administration only

10. At what age should a child receive their first dose of the MMR vaccine?

A. 6-12 months

B. 12-15 months

C. 18-24 months

D. 2-3 years

SAMPLE

Answers

SAMPLE

1. B
2. B
3. B
4. A
5. A
6. C
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. In which category is Ampicillin classified?

A. Antihistamine

B. Antibiotic

C. Pain reliever

D. Vitamin and Minerals

Ampicillin is classified as an antibiotic. This classification is based on its mechanism of action, which involves inhibiting the growth of bacteria. Antibiotics are substances that combat bacterial infections by targeting bacterial cell wall synthesis, protein synthesis, or other vital processes within bacterial cells. Ampicillin is part of the penicillin family and is effective against a variety of gram-positive and some gram-negative bacteria. It is particularly useful in treating infections such as respiratory tract infections, urinary tract infections, and meningitis, making it a critical medication in pediatric care for managing bacterial infections in children. The other categories listed, such as antihistamines, pain relievers, and vitamins/minerals, serve entirely different purposes. Antihistamines are primarily used to treat allergic reactions, pain relievers address discomfort, and vitamins and minerals contribute to nutritional support. Each of these categories does not relate to the antibacterial properties that define ampicillin's function, which solidifies its classification as an antibiotic.

2. Which action should be performed if chest tube drainage is observed to change in color?

A. Notify the physician immediately

B. Continue to monitor the drainage for further changes

C. Clamp the chest tube

D. Document the change and wait for further instructions

Monitoring the drainage for further changes is a prudent approach when there is a change in color observed in chest tube drainage. Changes in color can be indicative of various underlying issues, such as the presence of blood, infection, or other fluid changes, but it may not necessitate immediate intervention unless there are accompanying symptoms or significant variations in volume or character of the drainage. By choosing to continue monitoring, the healthcare provider can gather more information about the situation, such as whether the color change persists or if other signs (like increased drainage quantity, changes in vital signs, or signs of distress in the patient) occur alongside it. This allows for a more informed decision-making process, understanding that the context of the change will help determine the seriousness of the situation and if further actions are warranted. Monitoring can also help establish a trend in drainage characteristics over time, which can be useful for assessing the patient's condition and the efficacy of treatment. If significant or concerning changes occur, that's when communication with the physician would be necessary to decide on further interventions.

SAMPLE

3. What is the typical age for the onset of puberty in girls?

A. 6-10 years

B. 8-13 years

C. 10-15 years

D. 12-16 years

The typical age for the onset of puberty in girls is indeed between 8 to 13 years. This range reflects the biological changes that occur during this developmental stage. Puberty in girls begins with thelarche, which is the development of breast tissue, often followed by the onset of menstruation, known as menarche, which usually occurs a couple of years after the initial breast development. The onset age varies due to a combination of genetic, nutritional, and environmental factors. Starting at around 8 years old is considered normal for some girls, while others may not experience signs of puberty until around 13 years old. The choice of this range aligns well with standard medical guidelines and growth charts used by pediatricians, highlighting the variability in individual maturation processes. In contrast, younger ages such as 6-10 years may not encompass the full range of typical onset and can signify early puberty, which can lead to concerns if it occurs too soon. Later ages, such as 10-15 years or 12-16 years, may imply a delayed onset, which, while not inherently problematic, would be beyond the typical development window for most girls. Understanding this developmental stage is crucial for monitoring growth and addressing any potential health issues.

4. Which form of Diphenhydramine is listed for administration?

A. 25 mg tablets

B. 50 mg tablets

C. Liquid suspension

D. Capsules

In this context, Diphenhydramine is a common antihistamine used to alleviate symptoms of allergies, motion sickness, or insomnia. The correct answer highlights that 25 mg tablets of Diphenhydramine are a standard dosage form that is often prescribed or recommended for administration, particularly in pediatric patients. When considering the other forms mentioned, 50 mg tablets, liquid suspension, and capsules do exist, but they may not always be the preferred options in pediatric medicine. The 25 mg tablet form is particularly significant for dosing accuracy in children, as it allows for easier adjustment and adherence to recommended dosages based on weight and age. The choice of using a specific dosage form can have implications related to safety, efficacy, and ease of administration. In a pediatric context, lower dosages are critical to minimize the risk of side effects and ensure effective management of allergic reactions or other indications for which Diphenhydramine is used. Thus, the mention of the 25 mg tablet aligns with established guidelines for pediatric use.

5. When should school-age children have their first dental exam?

A. By age 1

B. By age 3

C. By age 5

D. By age 7

The guidelines recommend that children should have their first dental visit by age 1. This early visit is important for several reasons. It allows the dentist to assess the child's oral health, provide guidance on proper dental care practices, and assist parents with questions regarding teething, bottle use, and dietary habits that can affect dental health. Additionally, this early examination helps in identifying any potential issues at a stage when they are easier to treat, thereby establishing a good foundation for a child's ongoing dental health and encouraging the development of a positive attitude towards oral care from a young age. While some might suggest later ages for the first dental visit, starting at age 1 provides the earliest opportunity for preventive care and education, which can have long-lasting benefits.

SAMPLE

6. Why is Bridgett's weight monitored closely during her recovery?

A. To determine medication side effects

B. To assess nutritional needs

C. To ensure it aligns with her baseline

D. To prepare for future surgeries

Monitoring Bridgett's weight closely during her recovery is crucial to ensure it aligns with her baseline. This is significant because a child's weight can provide critical insights into their overall health and recovery status. If Bridgett's weight is significantly different from her baseline, it could indicate complications, nutritional deficits, or over-nutrition possibly leading to other health issues. Maintaining awareness of her baseline weight helps healthcare providers identify any changes that may require intervention, ensuring that her recovery progresses as expected and that her current health status is properly managed. This focus on adherence to her baseline also helps in establishing confidence in her recovery trajectory and any adjustments necessary in her care plan. While assessing nutritional needs and monitoring for medication side effects are important aspects of recovery, the emphasis in this context is specifically on ensuring that her weight remains consistent with what is typical for her, making it a primary focus in her post-recovery monitoring.

7. What is a classic sign of intussusception in a child?

A. Severe abdominal pain

B. "Currant jelly" stool

C. Yellow-green vomiting

D. Diarrhea

A classic sign of intussusception in a child is "currant jelly" stool. This distinctive type of stool results from the mixing of blood and mucus that occurs when part of the intestine telescopes into itself, leading to ischemia and subsequent bleeding. The appearance of the stool resembles jelly, often described as dark red or maroon in color, which serves as an important diagnostic clue for healthcare providers when evaluating a child presenting with abdominal pain and other gastrointestinal symptoms. Severe abdominal pain is frequently seen in cases of intussusception, but it is more of a symptom rather than a unique sign that specifically points to this condition. While yellow-green vomiting can occur in various gastrointestinal issues, it is not exclusive to intussusception and can be seen in other conditions as well. Diarrhea is also a common symptom in many pediatric gastrointestinal disorders and does not specifically indicate intussusception. Thus, the presence of "currant jelly" stool is a hallmark feature that assists in identifying this particular medical emergency.

SAMPLE

8. What should be requested for administering a 1000 mL bag of 0.9% Normal Saline?

A. Request an LPN

B. Request an RN

C. Handle it independently

D. Request a CNA

Administering a 1000 mL bag of 0.9% Normal Saline typically involves responsibilities that fall under the scope of practice for a Registered Nurse (RN). An RN is trained to perform the necessary assessments, ensure proper technique, monitor the patient's response, recognize any signs of complications, and document the procedure appropriately. The act of initiating and overseeing IV fluid therapy is a critical nursing responsibility, as it requires knowledge of the patient's medical history, an understanding of fluid management, and the ability to intervene if complications arise. Because of these factors, requesting an RN ensures that the patient receives safe and effective care. In contrast, while Licensed Practical Nurses (LPNs) and Certified Nursing Assistants (CNAs) play important roles in healthcare, their training and scope of practice differ. LPNs may assist with some aspects of patient care, but they may not be authorized to independently initiate IV fluids. CNAs typically handle basic patient care tasks and do not administer medications or IV fluids. Therefore, involving an RN in this situation is crucial for ensuring a high standard of care.

9. What should the nurse document after administering digoxin?

A. Only the dosage given

B. Patient's heart rate and any rhythm abnormalities

C. Patient's weight and vital signs

D. Time of medication administration only

After administering digoxin, it is essential for the nurse to document the patient's heart rate and any rhythm abnormalities. Digoxin is a cardiac glycoside used primarily to treat heart conditions such as atrial fibrillation and heart failure, and it has a narrow therapeutic index. Monitoring the heart rate is crucial because digoxin can cause bradycardia (a slower than normal heart rate), and understanding the patient's cardiac rhythm is vital in ensuring the safe administration of this medication. Documenting the heart rate helps determine if it is within the acceptable range before administration, and it provides a baseline for ongoing monitoring. Additionally, noting any rhythm abnormalities can indicate potential adverse effects or complications related to digoxin use, allowing for timely interventions if necessary. While other factors like dosage, patient weight, vital signs, and the time of administration are also important parts of medication documentation, they do not carry the same immediate significance in relation to the pharmacological effects and safety of digoxin as monitoring the heart rate and rhythm do.

SAMPLE

10. At what age should a child receive their first dose of the MMR vaccine?

A. 6-12 months

B. 12-15 months

C. 18-24 months

D. 2-3 years

The administration of the first dose of the MMR (measles, mumps, and rubella) vaccine is recommended between 12 and 15 months of age. This timing is crucial because it allows for the child to develop a robust immune response to the live attenuated viruses present in the vaccine. By vaccinating within this age range, the child is provided protection against these serious diseases at a time when they are particularly vulnerable, especially before they may be exposed to the viruses. The MMR vaccine is typically given as part of the routine immunization schedule, which is carefully designed to ensure optimal protection during early childhood when the risk of disease transmission is high. Receiving the vaccine too early, such as between 6 to 12 months of age, may not provide adequate immunity because it can be affected by the presence of maternal antibodies. Therefore, the guidelines specifically advise against administering it during that timeframe. Vaccinating at 18-24 months or 2-3 years delays the protection unnecessarily, leaving the child at risk for measles, mumps, and rubella during their formative years of social interaction. Adherence to the 12-15 month window balances both the safety and efficacy of vaccination, aligning with public health efforts to control

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://swiftriverpediatrics.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE