TACTICAL COMBAT CASUALTY CARE (TCCC) TIER 3 PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://tccctier3.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Why is it important to document prehospital combat casualty care?

A. It is a legal requirement only

B. To make assessments easier for the next provider

C. To avoid duplication of care

D. It helps with future medical training

2. What should be assessed before and after applying a splint?

A. Breathing pattern and heart rate

B. Pulses, skin color, and sensorimotor function

C. Level of consciousness and blood pressure

D. Body temperature and swelling

3. What is the difference between the TCCC Card (DD Form 1380) and the TCCC After Action Report (AAR)?

A. The TCCC Card is retrospective, while the AAR is real-time

B. The TCCC Card is for administrative use, while the AAR is for care documentation

C. The TCCC Card is filled in real-time at the point of injury, while the AAR is completed after the mission

D. The TCCC Card is only for medics, while the AAR is for everyone

4. In a TCCC scenario, what is the best initial step for managing a casualty with a suspected massive bleed?

A. Apply a pressure bandage

B. Administer fluids immediately

C. Assess and control the bleeding

D. Notify medical support

5. How is shock defined in a medical context?

A. Temporary loss of consciousness

B. Progressive cellular and tissue hypoxia

C. Sudden drop in blood pressure only

D. Acute pain in the chest

SAMPLE

6. What is the maximum duration a tourniquet can be left on a patient?

A. 1 hour
B. 2 hours
C. 4 hours
D. 6 hours

7. Which of the following is a contraindication for the use of opioids?

A. Severe hypertension
B. Hyperglycemia
C. Respiratory distress
D. Mild anxiety

8. Which airway device is used for patients who have suffered facial burns?

A. Endotracheal tube
B. Oropharyngeal airway
C. Nasopharyngeal airway
D. Cricothyrotomy

9. If a patient is experiencing an acute hemolytic reaction, which medication is the treatment of choice?

A. 50 mg of diphenhydramine
B. 25 mg of diphenhydramine
C. 10 mg of diphenhydramine
D. 100 mg of diphenhydramine

10. When should Priority Evac be initiated?

A. Within 2 hours of injury
B. Within 4 hours of injury
C. Within 24 hours of injury
D. As soon as resources are available

SAMPLE

Answers

SAMPLE

1. B
2. B
3. C
4. C
5. B
6. B
7. C
8. D
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. Why is it important to document prehospital combat casualty care?

A. It is a legal requirement only

B. To make assessments easier for the next provider

C. To avoid duplication of care

D. It helps with future medical training

Documenting prehospital combat casualty care serves several crucial purposes, one of which is to facilitate smoother transitions in patient care among providers. Accurate documentation ensures that the next provider has access to vital information about the patient's condition, treatments administered, and any interventions performed. This continuity allows for more effective assessments, enabling the subsequent care team to make informed clinical decisions quickly based on the latest information available, ultimately improving patient outcomes. Additionally, well-documented care can enhance communication among the various medical personnel involved and help prevent misinterpretations, thereby ensuring that care is coherent and comprehensive. This aspect supports the overall goal of TCCC to provide optimal care in combat situations, which is critical for the survival and recovery of the casualty. While other reasons for documentation are also significant, such as legal accountability, avoiding duplication of efforts, and contributing to future training, the immediate benefit of aiding the next provider is paramount in the context of prehospital care.

2. What should be assessed before and after applying a splint?

A. Breathing pattern and heart rate

B. Pulses, skin color, and sensorimotor function

C. Level of consciousness and blood pressure

D. Body temperature and swelling

The assessment of pulses, skin color, and sensorimotor function before and after applying a splint is crucial for ensuring that the extremity being splinted remains viable and does not suffer from further injury or compromise. Pulses provide vital information about blood flow to the area distal to the injury. If pulses are weak or absent after the application of the splint, it may indicate that circulation has been compromised, which is a critical concern that needs immediate attention. Skin color is another key indicator of circulation; for instance, a pale or cyanotic appearance could signal insufficient blood flow. Sensorimotor function includes assessing the patient's ability to move their fingers or toes and their sensation in the affected limb. A loss of motor function or sensation could suggest nerve injury or severe swelling that may arise due to improper splint application. These assessments ensure that the splint is applied correctly and that the injured limb has not developed any complications that would require adjustment to the splint or alternative interventions. The other options, while important in different contexts, do not directly address the specific concerns related to the implications of splinting an injured limb.

SAMPLE

3. What is the difference between the TCCC Card (DD Form 1380) and the TCCC After Action Report (AAR)?

A. The TCCC Card is retrospective, while the AAR is real-time

B. The TCCC Card is for administrative use, while the AAR is for care documentation

C. The TCCC Card is filled in real-time at the point of injury, while the AAR is completed after the mission

D. The TCCC Card is only for medics, while the AAR is for everyone

The TCCC Card (DD Form 1380) is designed to be filled out in real-time at the point of injury. It serves as a vital tool for documenting the immediate medical care provided to a casualty during tactical combat operations. This card captures crucial information about the injuries sustained, the interventions performed, and the patient's condition, all of which are critical for ongoing care as the casualty is evacuated to higher echelons of medical support. In contrast, the TCCC After Action Report (AAR) is a comprehensive document completed after the mission or operation has concluded. The AAR serves to analyze the overall effectiveness of the combat operation, including the medical response and outcomes, and provides recommendations for future operations. This distinction highlights the timing and purpose of each document: real-time documentation during an emergency versus an analytical review after the fact. Understanding this difference is essential for ensuring that proper medical and operational feedback processes are in place. The AAR can also serve to improve TCCC practices and protocols based on the insights derived from the experiences documented.

4. In a TCCC scenario, what is the best initial step for managing a casualty with a suspected massive bleed?

A. Apply a pressure bandage

B. Administer fluids immediately

C. Assess and control the bleeding

D. Notify medical support

In managing a casualty with a suspected massive bleed, the most critical initial step is to assess and control the bleeding. This step is foundational because uncontrolled hemorrhage is the leading cause of preventable death in traumatic injuries. Prioritizing the assessment allows the rescuer to determine the severity of the bleeding and the best method to manage it effectively, whether through direct pressure, the use of tourniquets, or other bleeding control measures. By focusing on controlling the bleed first, the rescuer can significantly increase the likelihood of saving the casualty's life. The determination of how and where to apply control measures is essential, as massive blood loss can lead to shock rapidly, thus requiring immediate intervention. Other steps, like applying a pressure bandage or administering fluids, may be necessary later but should follow the immediate action of controlling the hemorrhage. Notifying medical support is also important, but it does not address the immediate threat posed by the massive bleed. Therefore, assessing and controlling the bleeding is crucial in ensuring that the casualty receives the best chance of survival.

SAMPLE

5. How is shock defined in a medical context?

A. Temporary loss of consciousness

B. Progressive cellular and tissue hypoxia

C. Sudden drop in blood pressure only

D. Acute pain in the chest

In a medical context, shock is defined as a progressive cellular and tissue hypoxia, which indicates a critical condition where the body's tissues are not receiving enough oxygen to function properly. This condition results from inadequate blood flow, leading to impaired delivery of oxygen and nutrients and the accumulation of metabolic waste products. The deficiency in oxygen in the tissues triggers a cascade of physiological responses, ultimately leading to cellular dysfunction and, if not resolved, multi-organ failure. Recognizing shock is crucial for timely intervention, as it can occur due to various factors, such as blood loss, infection, or severe dehydration. The other options do not accurately encompass the definition of shock. For instance, a temporary loss of consciousness may occur in some forms of shock but does not encompass the broader systemic effects. A sudden drop in blood pressure is a symptom that can be associated with shock but does not define it holistically since shock can exist with normal or high blood pressure in its initial stages. Acute chest pain is not a defining characteristic of shock but rather a symptom of other possible conditions such as myocardial infarction or pulmonary embolism. Understanding shock in terms of cellular and tissue hypoxia allows for a more comprehensive approach to diagnosis and treatment.

6. What is the maximum duration a tourniquet can be left on a patient?

A. 1 hour

B. 2 hours

C. 4 hours

D. 6 hours

The maximum duration a tourniquet can be left on a patient depends on various factors, including the specific situation and the need to prevent limb loss while providing life-saving treatment. The accepted guideline generally indicates that a tourniquet can be safely left on for up to two hours in a tactical environment when assessing the risk versus the benefits of bleeding control. After two hours, tissue viability begins to be compromised due to lack of blood flow, which can lead to tissue necrosis and other complications. In prolonged scenarios, if a tourniquet must remain in place beyond this timeframe, it is crucial to monitor the patient closely for signs of tissue damage and to be prepared to perform additional medical interventions as soon as possible. While some training may touch on longer durations, such as four or six hours, these are typically associated with very specific circumstances and should be approached with caution. The two-hour guideline is the most widely recognized standard for practice in emergency situations, balancing effective life-saving measures with the risks associated with prolonged ischemia.

SAMPLE

7. Which of the following is a contraindication for the use of opioids?

A. Severe hypertension

B. Hyperglycemia

C. Respiratory distress

D. Mild anxiety

Opioids are powerful analgesics commonly used for pain management, but their use comes with specific contraindications due to their effects on the body. Respiratory distress is a critical condition where there is inadequate breathing or oxygenation. The administration of opioids can lead to respiratory depression, a serious side effect where the breathing rate decreases, further exacerbating respiratory distress. In such cases, the risks of administering opioids outweigh the benefits, making it a primary contraindication. The other conditions listed, while they may have their own considerations for treatment, do not directly contraindicate the use of opioids in the way that respiratory distress does. Severe hypertension may require careful monitoring or different treatment approaches, but it does not have the immediate life-threatening implications associated with respiratory depression. Hyperglycemia, which refers to high blood sugar levels, can be addressed through other medical means and is not an immediate concern with opioid use. Mild anxiety is generally not a contraindication for opioid usage, as they are often used to provide relief in a variety of painful situations. Thus, respiratory distress stands out as a compelling reason to withhold opioid treatment.

8. Which airway device is used for patients who have suffered facial burns?

A. Endotracheal tube

B. Oropharyngeal airway

C. Nasopharyngeal airway

D. Cricothyrotomy

In the context of treating patients who have sustained facial burns, the correct choice is the cricothyrotomy. This procedure is particularly relevant when there is the potential for airway compromise due to edema or obstruction caused by the burn injuries. Facial burns can lead to significant swelling of the soft tissues in the neck and face, which can complicate traditional methods of securing an airway, such as using an endotracheal tube or other airway adjuncts. A cricothyrotomy provides a direct route to the airway by making an incision through the skin over the cricothyroid membrane, allowing for ventilation when other methods may be ineffective or impossible due to swelling, trauma, or severe burns in the facial area. This method is preferable in emergency situations where rapid access to the airway is crucial for survival, especially when dealing with thermal injuries that could obstruct the airway. Endotracheal intubation may be considered in many cases, but it can be challenging in patients with facial burns due to the decreased visibility and potential for airway obstruction. Oropharyngeal and nasopharyngeal airways are generally not effective in patients with facial injuries since they could exacerbate injury or cause additional trauma. Therefore, the cricothyrotomy is

SAMPLE

9. If a patient is experiencing an acute hemolytic reaction, which medication is the treatment of choice?

A. 50 mg of diphenhydramine

B. 25 mg of diphenhydramine

C. 10 mg of diphenhydramine

D. 100 mg of diphenhydramine

In the case of an acute hemolytic reaction, the use of diphenhydramine is aimed at managing symptoms such as allergic reactions, which may present in this context. Diphenhydramine is an antihistamine that can help alleviate symptoms like itching, hives, and other mild allergic responses that can occur during transfusion reactions or similar situations. The specific dosage of 25 mg is considered appropriate for adults experiencing moderate symptoms related to an allergic reaction. It is a well-established starting dose that provides effective symptom relief without the risk of overmedication, which could lead to increased side effects. Higher doses, such as 50 mg or 100 mg, might lead to more pronounced sedation and other side effects without significantly enhancing the therapeutic effect for mild to moderate symptoms in most patients. Therefore, the choice of 25 mg strikes a balance between efficacy and safety, making it the treatment of choice in this context. Overall, administering the correct dose is crucial in managing acute hemolytic reactions effectively while minimizing potential adverse effects on the patient.

10. When should Priority Evac be initiated?

A. Within 2 hours of injury

B. Within 4 hours of injury

C. Within 24 hours of injury

D. As soon as resources are available

Priority Evacuation is a critical component of Tactical Combat Casualty Care, aimed at ensuring that casualties who are most likely to die without immediate medical intervention receive timely evacuation to definitive care. The appropriate timeframe for initiating Priority Evac is typically within 4 hours of injury. This timeframe is based on the understanding that certain traumatic injuries, especially those involving severe hemorrhaging or potential airway compromise, require urgent medical attention to prevent deterioration and increase the chances of survival. Initiating evacuation within this timeframe appropriately aligns with the need for rapid intervention to address life-threatening conditions that may not be immediately treatable in the field. The 4-hour window is recognized as a benchmark in the tactical environment, beyond which the risks of mortality significantly increase for certain types of injuries. In contrast, initiating Priority Evac within 2 hours, while beneficial, may not always be a realistic expectation in varied operational scenarios where resource availability and environmental conditions come into play. Waiting for 24 hours could lead to catastrophic outcomes for the casualty, as the likelihood of complications and death increases significantly after such a long delay. Therefore, evacuating as soon as resources are available is a philosophy that emphasizes readiness, but does not specifically address the critical window for intervention, which is essential in this

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://tccctier3.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE