TRANSFUSION THERAPY PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://transfusiontherapy.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1
Table of Contents 2
Introduction 3
How to Use This Guide 4
Questions 5
Answers 8
Explanations 10
Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What causes an acute hemolytic transfusion reaction?

A. ABO incompatibility

B. Allergic reaction

C. Bacterial contamination

D. Febrile non-hemolytic

2. Steps of the coagulation cascade starting from Factor X activation

A. Factor X Prothrombin Thrombin Fibrinogen Fibrin

B. Factor X Thrombin Prothrombin Fibrinogen Fibrin

C. Fibrin Fibrinogen Prothrombin Thrombin

D. Prothrombin Factor X Thrombin Fibrin

3. TXA dosing for patients weighing more than 40 kg: initial loading dose

A. 500 mg over 10 minutes

B. 1 g over 10 minutes

C. 2 g over 10 minutes

D. 1 g over 1 hour

4. Approximately what percentage reduction in factor levels is typically needed before aPTT becomes prolonged?

A. 10%

B. 60%

C. 30%

D. 90%

5. When should the warming cartridge be inserted in the Ranger blood warmer system?

A. Before priming with PRBCs or fluid

B. After priming with PRBCs

C. During transfusion

D. Never use a cartridge

SAMPLE

6. According to ASA guidelines, RBC transfusion is almost always indicated when Hb falls below which value?

A. Hb < 6-7 g/dL

B. Hb < 9-10 g/dL

C. Hb < 12-13 g/dL

D. Hb < 3-4 g/dL

7. What is the hematocrit of the blood product returned during blood salvage?

A. 30-40%

B. 70-80%

C. 50-60%

D. 40-50%

8. What precipitate remains when FFP is thawed slowly?

A. Leukocytes

B. Cryoprecipitate

C. RBCs

D. Platelets

9. Which component is assessed in a Type & Screen?

A. Blood type and antibody screening

B. Coagulation profile

C. Platelet count

D. Hemoglobin level

10. Before elective surgery for a Jehovah's Witness patient, which action should be completed?

A. Blood product use

B. Patient signs a blood refusal form

C. Do not discuss blood management

D. Postpone surgery

SAMPLE

Answers

SAMPLE

1. A
2. A
3. B
4. C
5. A
6. A
7. C
8. B
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. What causes an acute hemolytic transfusion reaction?

A. ABO incompatibility

B. Allergic reaction

C. Bacterial contamination

D. Febrile non-hemolytic

The key idea is that acute hemolytic transfusion reactions happen when the recipient has preexisting antibodies against donor red blood cell antigens, most classically against ABO antigens. If blood with incompatible ABO type is transfused, these antibodies bind to the donor red cells and activate complement, causing rapid intravascular destruction of those cells. This clash leads to symptoms such as fever, severe back or chest pain, hypotension, hemoglobinemia, and often hemoglobinuria, and can progress to shock and kidney injury. Laboratory testing typically shows a positive direct antiglobulin (DAT) test, reflecting in vivo red cell destruction. Other transfusion reactions do not involve this immediate and widespread red cell destruction. An allergic reaction is due to sensitivity to plasma proteins, usually presenting as hives or angioedema rather than intravascular hemolysis. A febrile non-hemolytic reaction is driven by cytokines or antibodies to donor leukocytes and presents with fever without hemolysis. Bacterial contamination causes a septic transfusion reaction with fever, chills, hypotension, and sepsis-like symptoms, but not the classic hemolysis pattern of acute intravascular destruction of donor red cells.

2. Steps of the coagulation cascade starting from Factor X activation

A. Factor X Prothrombin Thrombin Fibrinogen Fibrin

B. Factor X Thrombin Prothrombin Fibrinogen Fibrin

C. Fibrin Fibrinogen Prothrombin Thrombin

D. Prothrombin Factor X Thrombin Fibrin

In the coagulation cascade, the common pathway starts when Factor X is activated to Xa. On a phospholipid surface with calcium and Factor V, Xa forms the prothrombinase complex that converts prothrombin (Factor II) into thrombin (Factor IIa). Thrombin then cleaves soluble fibrinogen (Factor I) to form fibrin, which polymerizes to stabilize the clot. So the sequence beginning with Factor X activation proceeds as: Factor X activation Prothrombin Thrombin Fibrinogen Fibrin. This order is correct because each step depends on the previous one: prothrombin must be converted to thrombin, and thrombin must then convert fibrinogen to fibrin. Directly going from Factor X to thrombin would bypass the essential prothrombinase-driven conversion, and forming fibrin before fibrinogen would be impossible.

3. TXA dosing for patients weighing more than 40 kg: initial loading dose

A. 500 mg over 10 minutes

B. 1 g over 10 minutes

C. 2 g over 10 minutes

D. 1 g over 1 hour

The main idea is to rapidly achieve therapeutic plasma levels of tranexamic acid to inhibit fibrinolysis during bleeding. For patients weighing more than 40 kg, the initial loading dose is 1 g given IV over 10 minutes, so peak levels are reached quickly and treatment starts promptly. This aligns with common trauma and surgical protocols that emphasize a fast, short infusion to maximize early clot stability. After the loading dose, a maintenance dose of 1 g is typically given over several hours (often 8 hours). The other options either underdose, overdose, or deliver the drug too slowly to be fully effective at the outset, which is why they're not used as the initial loading dose.

SAMPLE

4. Approximately what percentage reduction in factor levels is typically needed before aPTT becomes prolonged?

A. 10%

B. 60%

C. 30%

D. 90%

The test is focusing on how sensitive the aPTT is to decreases in intrinsic pathway factors (such as VIII, IX, XI, XII). aPTT tends to become prolonged only after a notable drop in these factor activities. A commonly cited approximate threshold is about a 30% reduction from normal factor levels, meaning the aPTT starts to lengthen when factor activity falls to around 70% of normal. In other words, you don't need a huge deficiency to see prolongation; roughly one third less activity is enough to affect the aPTT. So, the best choice is the option representing about a 30% reduction. Smaller drops (like 10%) usually won't prolong the aPTT, while larger drops (60% or 90%) would certainly prolong, but the question asks for the approximate threshold, which is around 30%.

5. When should the warming cartridge be inserted in the Ranger blood warmer system?

A. Before priming with PRBCs or fluid

B. After priming with PRBCs

C. During transfusion

D. Never use a cartridge

The warming cartridge must be in place before any priming so that every drop that enters the patient is warmed. The cartridge is the heating element in the Ranger system, and if you prime the circuit with PRBCs or fluids first, the initial volume would flow through unheated pathways, producing a cold bolus and compromising temperature control. By inserting the cartridge before priming, you ensure the entire transfusion line is warmed from the start, improving safety and achieving the intended warmer output. Inserting during transfusion or never using a cartridge would defeat the warming purpose and risk temperature drop or instability.

6. According to ASA guidelines, RBC transfusion is almost always indicated when Hb falls below which value?

A. Hb < 6-7 g/dL

B. Hb < 9-10 g/dL

C. Hb < 12-13 g/dL

D. Hb < 3-4 g/dL

Transfusion decisions hinge on a balance between delivering enough oxygen to tissues and avoiding the risks of transfusion. The ASA supports a restrictive approach for stable patients, meaning transfuse only when hemoglobin is at a very low level where tissue oxygen delivery would be compromised. That is why this threshold is considered the best match for "almost always indicated." Transfusing at higher hemoglobin levels would expose patients to unnecessary risks when their oxygen delivery could be adequate without transfusion. In practice, there are exceptions for patients with active bleeding, ischemic heart disease, or symptomatic anemia, where clinicians might transfuse at higher levels, but the general guideline emphasizes waiting until hemoglobin is very low.

SAMPLE

7. What is the hematocrit of the blood product returned during blood salvage?

A. 30-40%

B. 70-80%

C. 50-60%

D. 40-50%

During blood salvage, collected shed blood is processed to concentrate the red cells while removing most plasma. The final washed red cell product is a concentrated suspension with a hematocrit in the range of about 50–60%, often around 60%. This reflects that the washing step removes most plasma and anticoagulant, yielding higher RBC concentration than whole blood, but not as high as undiluted red cells. If salvage is not washed, the hematocrit would be lower, roughly 30–40%, due to more plasma remaining.

8. What precipitate remains when FFP is thawed slowly?

A. Leukocytes

B. Cryoprecipitate

C. RBCs

D. Platelets

When plasma is slowly thawed in the cold, certain clotting proteins precipitate out of solution and form a visible solid. This solid, rich in fibrinogen, factor VIII, factor XIII, and von Willebrand factor, is called cryoprecipitate. So the precipitate that remains after slow thawing of fresh frozen plasma is cryoprecipitate, which is then collected for clinical use. The other options aren't formed as a precipitate from plasma in this process because red cells, platelets, and leukocytes are cellular components or are not the cold-insoluble protein group that cryoprecipitate represents.

9. Which component is assessed in a Type & Screen?

A. Blood type and antibody screening

B. Coagulation profile

C. Platelet count

D. Hemoglobin level

In a Type & Screen, the focus is compatibility for transfusion: first, determine the patient's blood type (ABO and Rh) and then screen for antibodies against red cell antigens. The blood type tells you which donor blood can safely be given, and the antibody screen looks for unexpected antibodies that could cause a reaction if donor cells with corresponding antigens are transfused. If no antibodies are detected, a crossmatch may be performed to confirm compatibility before transfusion. Coagulation profile evaluates clotting function, not compatibility of blood products. Platelet count assesses bleeding risk or platelet transfusion needs, but it's not part of the Type & Screen. Hemoglobin level indicates oxygen-carrying capacity and overall anemia status, not the compatibility testing performed in a Type & Screen.

SAMPLE

10. Before elective surgery for a Jehovah's Witness patient, which action should be completed?

A. Blood product use

B. Patient signs a blood refusal form

C. Do not discuss blood management

D. Postpone surgery

The essential idea is to honor the patient's autonomy by obtaining explicit, written consent that they refuse blood transfusions. Before elective surgery, it's crucial to have the patient sign a blood refusal form and have it documented in the medical record. This clearly communicates that the patient declines transfusion of whole blood and major blood components, guiding the entire perioperative team to pursue bloodless management strategies and to discuss alternatives and risks with the patient beforehand. Ensure the patient understands what is refused, any acceptable alternatives, and confirm capacity to consent; involve a surrogate only if the patient lacks capacity. With a signed refusal, the team can plan accordingly and avoid transfusions unless the patient revokes the decision. Postponing surgery isn't inherently required—the priority is clear, documented wishes to respect the patient's beliefs and legal/ethical obligations.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://transfusiontherapy.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE