
TRAUMA CARE AFTER RESUSCITATION (TCAR) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://tcar.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **Central venous pressure is most commonly used to estimate which physiologic parameter?**

   A. Preload

   B. Afterload

   C. Contractility

   D. Heart rate

2. **If dissection is suspected, which chest X-ray finding is most likely?**

   A. Hyperinflation

   B. Normal mediastinum

   C. Cardiac silhouette enlargement

   D. Widened mediastinum.

3. **In management of hemothorax, what is the typical sequence after chest tube placement?**

   A. CT scan to evaluate, then intrapleural tPA or VATS

   B. Immediate VATS without imaging

   C. Discharge

   D. MRI

4. **In an Rh- patient exposed to Rh+ blood, what prophylaxis within 72 hours is indicated?**

   A. Rhogam within 72 hours

   B. Vaccination with anti-D

   C. High-dose steroids

   D. No intervention

5. **In the setting of massive transfusion, what electrolyte management is indicated?**

   A. Calcium replacement

   B. Potassium replacement

   C. Sodium bicarbonate infusion

   D. Magnesium sulfate infusion

SAMPLE

6. On chest radiograph, air appears as which color?

A. White

B. Gray

C. Black

D. Purple

7. Which statement accurately reflects the 20 mm Hg threshold discussed in relation to capillary perfusion?

A. Pressures above 20 mm Hg are harmless

B. Pressures below 5 mm Hg are dangerous

C. 20 mm Hg is unrelated to capillaries

D. Pressures above 20 mm Hg can risk capillary collapse

8. What is the relationship between creatine kinase and creatinine as described?

A. Creatine Kinase Is a Muscle Enzyme; Creatinine Is a Different Substance

B. CK Is a Waste Product Cleared by Kidneys

C. CK Measures Liver Function

D. CK Is a Hormone

9. In C4-C5 cervical spinal cord injury, edema tends to blossom with time and may necessitate ventilatory support. Which option reflects this?

A. Edema resolves quickly without support

B. Edema blossoms with time and may require ventilatory support

C. Edema is minimal

D. Edema is limited to the injury level

10. Which agents are considered alpha-adrenergic SNS agonists?

A. Norepinephrine, Epinephrine, Phenylephrine

B. Dobutamine, Albuterol

C. Dopamine, Acetylcholine

D. Atropine, Ephedrine

SAMPLE

Answers

SAMPLE

1. A
2. D
3. C
4. A
5. A
6. C
7. D
8. A
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. Central venous pressure is most commonly used to estimate which physiologic parameter?

A. Preload

B. Afterload

C. Contractility

D. Heart rate

Central venous pressure reflects the pressure in the right atrium and, by extension, the filling pressure of the right heart. This makes it a rough proxy for preload—the amount of ventricular stretch at end-diastole due to venous return. In practice, CVP helps gauge how much volume is returning to the heart and how full the ventricles are before they contract. It's not a direct measure of afterload (the resistance the heart must push against), contractility (the heart's intrinsic pumping strength), or heart rate (which is governed by the rate of electrical impulses). So the parameter CVP most commonly estimates is preload.

2. If dissection is suspected, which chest X-ray finding is most likely?

A. Hyperinflation

B. Normal mediastinum

C. Cardiac silhouette enlargement

D. Widened mediastinum.

When aortic dissection is suspected, the most likely chest X-ray finding is widening of the mediastinum. The dissection creates a separation and expansion of the aortic wall, and the intramural hematoma plus the enlarged aortic segment push the mediastinal contours outward, making the mediastinal shadow appear broader on the film. This radiographic clue is the classic tip-off on plain X-ray, though it isn't perfect and a normal chest X-ray doesn't rule out dissection. In contrast, hyperinflation points to obstructive lung disease, a normal mediastinum would not suggest dissection, and an enlarged cardiac silhouette indicates cardiomegaly rather than an aortic catastrophe. If dissection is still suspected clinically, CT angiography provides a definitive diagnosis.

3. In management of hemothorax, what is the typical sequence after chest tube placement?

A. CT scan to evaluate, then intrapleural tPA or VATS

B. Immediate VATS without imaging

C. Discharge

D. MRI

After placing a chest tube for a hemothorax, the next step is to reassess with imaging to determine how much blood remains and whether clots are present. A CT scan is especially helpful because it can reveal residual, organized blood that a simple X-ray might miss, as well as how well the lung has re-expanded and whether there is ongoing bleeding. Based on what you find, you tailor the next move: if there's a significant residual collection or organized clot, you can use intrapleural tPA to liquefy and drain it, or proceed to video-assisted thoracoscopic surgery (VATS) to evacuate the clot and achieve definitive drainage and hemostasis. Discharging the patient right after chest tube placement would overlook ongoing issues; MRI isn't typically used in this acute setting, and jumping straight to VATS without imaging bypasses crucial assessment.

SAMPLE

4. In an Rh- patient exposed to Rh+ blood, what prophylaxis within 72 hours is indicated?

A. Rhogam within 72 hours

B. Vaccination with anti-D

C. High-dose steroids

D. No intervention

The concept being tested is preventing Rh sensitization after exposure to Rh-positive blood. In an Rh-negative person, encountering Rh-positive red cells can trigger the development of anti-D antibodies, which can cause hemolytic problems in future pregnancies. The best protection is giving Rh immune globulin (Rhogam) within 72 hours of exposure. Rhogam provides passive anti-D antibodies that bind and clear Rh-positive cells before the recipient's immune system can mount its own response, effectively preventing sensitization. It's not a vaccine and doesn't create lasting immunity in the recipient; it prevents an immune reaction to future exposures. Timing matters because the sooner the Rh-positive cells are cleared, the less likely an immune response will develop. The other options don't fit: there is no anti-D vaccine to administer, high-dose steroids aren't used for prophylaxis against Rh sensitization, and doing nothing risks sensitization and potential hemolytic disease in future pregnancies.

5. In the setting of massive transfusion, what electrolyte management is indicated?

A. Calcium replacement

B. Potassium replacement

C. Sodium bicarbonate infusion

D. Magnesium sulfate infusion

Massive transfusion uses large amounts of citrated blood products. The citrate acts as an anticoagulant, but as it circulates, it binds ionized calcium in the patient's blood. This chelation lowers ionized calcium levels, which are essential for normal coagulation and for cardiac contractility. When calcium becomes low, coagulation can worsen and the heart may lose contractile strength, making bleeding harder to control and hemodynamics less stable. Replacing calcium directly addresses this citrate-induced hypocalcemia, helping restore clot formation and stabilize the patient during rapid transfusion. Typically, calcium is given as calcium gluconate or calcium chloride, with monitoring of ionized calcium to guide dosing. Keep in mind that stored blood can contribute to other shifts, such as potential hyperkalemia, but the key immediate management during massive transfusion is maintaining calcium levels.

6. On chest radiograph, air appears as which color?

A. White

B. Gray

C. Black

D. Purple

In chest radiographs, the color you see reflects how much the substance attenuates the X-ray beam. Air has the lowest density in the chest, so it allows most X-ray photons to pass through and reaches the receptor with little attenuation. That makes air appear radiolucent, or black, on the image. Soft tissues and fluids attenuate more and show up as lighter gray, while bone attenuates the most and appears white. The option of purple isn't a radiographic color at all. So air in the chest appears black.

SAMPLE

7. Which statement accurately reflects the 20 mm Hg threshold discussed in relation to capillary perfusion?

A. Pressures above 20 mm Hg are harmless

B. Pressures below 5 mm Hg are dangerous

C. 20 mm Hg is unrelated to capillaries

D. Pressures above 20 mm Hg can risk capillary collapse

Capillary perfusion depends on a favorable pressure gradient across the capillary wall. The capillary pressure is typically around 20 mm Hg. When external or interstitial pressure on a tissue rises above this level, the transmural pressure across the capillary walls drops enough to compress the vessels. As a result, the capillaries can collapse and microcirculatory flow is impaired, increasing the risk of tissue ischemia. That's why pressures above 20 mm Hg are concerning for capillary patency. The other statements don't fit this physiology: pressures above 20 mm Hg aren't harmless, very low pressures (like below 5 mm Hg) aren't defined as inherently dangerous to capillaries, and 20 mm Hg is indeed relevant to capillary behavior.

8. What is the relationship between creatine kinase and creatinine as described?

A. Creatine Kinase Is a Muscle Enzyme; Creatinine Is a Different Substance

B. CK Is a Waste Product Cleared by Kidneys

C. CK Measures Liver Function

D. CK Is a Hormone

Creatine kinase and creatinine are two distinct substances serving different roles. Creatine kinase is an enzyme found mainly in muscle that helps regenerate ATP by transferring a phosphate to creatine, forming phosphocreatine. When muscle tissue is damaged, CK leaks into the bloodstream, so elevated CK levels indicate muscle injury (and in specific isoforms, heart muscle injury as well). Creatinine, by contrast, is a waste product produced from creatine metabolism and is cleared by the kidneys; it's used primarily to assess kidney function, not muscle injury. So, the correct description is that CK is a muscle enzyme and creatinine is a different substance.

9. In C4-C5 cervical spinal cord injury, edema tends to blossom with time and may necessitate ventilatory support. Which option reflects this?

A. Edema resolves quickly without support

B. Edema blossoms with time and may require ventilatory support

C. Edema is minimal

D. Edema is limited to the injury level

Edema after a cervical spinal cord injury often evolves and can worsen over time, especially at higher levels where respiratory muscles are affected. At C4–C5, the diaphragm and accessory respiratory muscles are at risk because their innervation lies around this region; swelling can progressively impair diaphragmatic function and breathing. Because of this progression, edema blossoming with time and potentially requiring ventilatory support best captures the clinical reality. Edema being minimal or resolving quickly would not reflect the typical secondary-injury cascade in this area, and limiting edema to the exact injury level wouldn't account for evolving respiratory compromise.

SAMPLE

10. Which agents are considered alpha-adrenergic SNS agonists?

A. Norepinephrine, Epinephrine, Phenylephrine

B. Dobutamine, Albuterol

C. Dopamine, Acetylcholine

D. Atropine, Ephedrine

Alpha receptors, when activated, produce vasoconstriction and increased vascular resistance—key actions for supporting blood pressure in resuscitation. Norepinephrine and epinephrine directly stimulate these alpha receptors, with norepinephrine acting strongly on alpha-1 (and some beta-1) receptors, and epinephrine stimulating both alpha and beta receptors (vasoconstriction via alpha while also affecting heart rate and contractility via beta). Phenylephrine is a classic alpha-1–selective agonist, producing vasoconstriction without direct beta effects. Other agents listed either don't primarily activate alpha receptors. Dobutamine mainly targets beta-1 receptors to boost cardiac output; albuterol is a beta-2 agonist for bronchodilation; dopamine's effects vary with dose and aren't limited to alpha stimulation; acetylcholine and atropine act on cholinergic receptors rather than adrenergic receptors; ephedrine has mixed indirect and direct sympathomimetic actions and is not a pure alpha-adrenergic agonist.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://tcar.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE