
UNITED STATES MEDICAL LICENSING EXAMINATION (USMLE) STEP 3 PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://usmlestep3.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the gold standard for confirming iron deficiency?

A. Serum ferritin measurement

B. Bone marrow biopsy

C. Soluble transferrin receptor assay

D. Transferrin saturation

2. What creatinine level indicates a need for caution when using contrast in CT scans?

A. 1.0

B. 1.5

C. 2.0

D. 2.5

3. What should be considered a potential cause of folliculitis in whirlpool bath users?

A. Group A Streptococcus

B. Staphylococcus aureus

C. Pseudomonas aeruginosa

D. Escherichia coli

4. For a patient with Lyme disease presenting with a rash, joint pain, or Bell's palsy, what is the appropriate first-line treatment?

A. IV ceftriaxone

B. Doxycycline or amoxicillin

C. Clindamycin

D. Quinine

5. Which type of rash is associated with coccidioidomycosis?

A. Erythema nodosum

B. Maculopapular rash

C. Diffuse urticarial rash

D. Vesicular rash

SAMPLE

6. What is the most common cardiac manifestation of Lyme disease?

A. Myocarditis
B. AV conduction block or defect
C. Pericardial effusion
D. Arrhythmias

7. What is the most common cause of urticaria?

A. Mast cell activation and IgG response
B. Activation of IgE and mast cells
C. Autoimmune mechanism
D. Drug hypersensitivity reaction

8. What demographic is most commonly affected by Takayasu's arteritis?

A. Young Asian females
B. Middle-aged Caucasian males
C. Older African American males
D. Young Hispanic females

9. What is the primary treatment option for diffuse esophageal spasm?

A. Surgical resection
B. Supportive care
C. Proton pump inhibitors
D. Chemotherapy

10. What is a distinguishing feature of pemphigus foliaceus?

A. Presence of oral lesions
B. Deeper blisters that are less likely to rupture
C. Association with other autoimmune disorders
D. High likelihood of malignancy

SAMPLE

Answers

SAMPLE

1. B
2. B
3. C
4. B
5. A
6. B
7. B
8. A
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. What is the gold standard for confirming iron deficiency?

A. Serum ferritin measurement

B. Bone marrow biopsy

C. Soluble transferrin receptor assay

D. Transferrin saturation

The gold standard for confirming iron deficiency is a bone marrow biopsy, particularly when looking for the presence of iron stores. In a healthy individual, bone marrow is expected to have adequate iron stores, which can be assessed histologically. If iron deficiency is present, there will be a marked decrease or absence of iron deposits in the marrow. While serum ferritin measurement is commonly used and reflects the body's iron stores, it can be influenced by inflammatory states and may not always accurately represent iron deficiency. Similarly, transferrin saturation and soluble transferrin receptor assays provide useful insights into iron metabolism but do not definitively confirm iron deficiency at the tissue level in the same way that a bone marrow biopsy does. Therefore, when direct evidence of iron stores is required to confirm a diagnosis of iron deficiency, a bone marrow biopsy serves as the definitive method.

2. What creatinine level indicates a need for caution when using contrast in CT scans?

A. 1.0

B. 1.5

C. 2.0

D. 2.5

When considering the use of contrast in CT scans, a creatinine level of 1.5 mg/dL is a significant threshold requiring caution. This level is associated with an increased risk of contrast-induced nephropathy, especially in patients with pre-existing kidney issues or other comorbidities. When renal function diminishes, the kidneys become less capable of filtering out contrast agents, which can lead to an accumulation of the agent in the body, potentially causing injury to renal tubules. At creatinine levels above 1.5 mg/dL, there is a notable heightened risk of complications, including acute kidney injury, following the administration of iodinated contrast material. Creatinine levels lower than 1.5 mg/dL, such as 1.0 mg/dL, may be considered safer for the use of contrast, as they typically indicate better renal function. Levels above 1.5 mg/dL, specifically those at or above 2.0 mg/dL and higher, significantly raise the concern regarding patient safety and necessitate more stringent protocols before proceeding with imaging that requires contrast. Thus, while a creatinine level of 1.5 mg/dL does warrant caution, it is generally still within a range where the benefits

SAMPLE

3. What should be considered a potential cause of folliculitis in whirlpool bath users?

A. Group A Streptococcus

B. Staphylococcus aureus

C. Pseudomonas aeruginosa

D. Escherichia coli

In whirlpool bath users, folliculitis is commonly caused by Pseudomonas aeruginosa, which is a type of bacteria well-known for thriving in warm, moist environments such as hot tubs and whirlpools. This organism can colonize the skin's hair follicles, leading to inflammation and infection, particularly in individuals who use these facilities often. Pseudomonas aeruginosa is capable of surviving in chlorinated water, and its presence in whirlpools, especially those that are poorly maintained, can lead to outbreaks of folliculitis. The clinical presentation often includes the development of itchy, painful red bumps around hair follicles, which typically occur in areas of skin that have been in contact with contaminated water. While other bacteria can cause skin infections, they are less specifically associated with whirlpool bath settings compared to Pseudomonas aeruginosa. For instance, Staphylococcus aureus is a common skin pathogen but is not unique to the aquatic environment. Group A Streptococcus is known for causing abscesses and cellulitis rather than folliculitis in this context, and Escherichia coli is primarily associated with gastrointestinal infections and not typically linked to skin infections from whirlpool baths. Hence, when considering the unique environmental

4. For a patient with Lyme disease presenting with a rash, joint pain, or Bell's palsy, what is the appropriate first-line treatment?

A. IV ceftriaxone

B. Doxycycline or amoxicillin

C. Clindamycin

D. Quinine

The appropriate first-line treatment for Lyme disease, particularly when the patient presents with conditions such as rash, joint pain, or Bell's palsy, is doxycycline or amoxicillin. These antibiotics are effective against the bacterium Borrelia burgdorferi, which causes Lyme disease. Doxycycline is often preferred due to its efficacy, ease of dosing, and the added benefit of treating co-existing infections like anaplasmosis, especially in endemic areas. However, amoxicillin is a suitable alternative for patients who may have contraindications to doxycycline, such as pregnant women or young children. Using either doxycycline or amoxicillin promptly addresses the bacterial infection and helps prevent potential complications associated with Lyme disease. Starting treatment early is crucial in reducing symptoms and preventing the progression to more serious manifestations. Other options like IV ceftriaxone would be more appropriate for patients with severe or refractory cases, such as those with neurological manifestations requiring a more aggressive approach. Clindamycin and quinine are not standard treatments for Lyme disease. Clindamycin is ineffective against Borrelia, while quinine is used for malaria but has no role in the treatment of Lyme disease. Thus, doxycycline or amoxicillin remains the

SAMPLE

5. Which type of rash is associated with coccidioidomycosis?

A. Erythema nodosum

B. Maculopapular rash

C. Diffuse urticarial rash

D. Vesicular rash

Coccidioidomycosis, also known as Valley Fever, is a fungal infection caused by Coccidioides species. One of the notable cutaneous manifestations associated with this infection is erythema nodosum. Erythema nodosum presents as painful, red nodules that typically appear on the lower extremities. It is an inflammatory reaction often triggered by various underlying conditions, including infections like coccidioidomycosis. The relationship between coccidioidomycosis and erythema nodosum highlights the body's immune response to the infection. This type of rash usually indicates a systemic reaction and is often seen in patients with a hypersensitivity response to the fungal antigens. The other types of rashes listed are not characteristically associated with coccidioidomycosis. For example, while a maculopapular rash could appear in various infections, it is not specific to coccidioides. Similarly, diffuse urticarial rashes and vesicular rashes are linked with different conditions or infections and do not correlate with the immune response observed in coccidioidomycosis. Hence, erythema nodosum is the most relevant type of rash associated with this fungal

6. What is the most common cardiac manifestation of Lyme disease?

A. Myocarditis

B. AV conduction block or defect

C. Pericardial effusion

D. Arrhythmias

The most common cardiac manifestation of Lyme disease is indeed an atrioventricular (AV) conduction block. Lyme disease, caused by the spirochete Borrelia burgdorferi, can lead to a variety of clinical manifestations, including those affecting the heart. While myocarditis, pericardial effusion, and arrhythmias can occur as manifestations of Lyme disease, they do not present as frequently or characteristically as AV conduction abnormalities. Specifically, the AV block seen in Lyme disease is often a first-degree or, more notably, a high-degree AV block that can result in bradycardia or syncope. This occurs as the spirochete affects the cardiac conduction system, leading to impaired electrical conduction. The presence of an AV block in a patient with Lyme disease usually indicates a more severe or systemic form of the disease, emphasizing the need for timely treatment with appropriate antibiotics to prevent further complications. Overall, the predominance of AV conduction defects in Lyme disease underscores the importance of recognizing the signs and symptoms of this illness in patients presenting with relevant clinical features, such as the characteristic erythema migrans rash, joint pain, or neurological symptoms, in order to provide effective management and prevent serious cardiac complications.

SAMPLE

7. What is the most common cause of urticaria?

A. Mast cell activation and IgG response

B. Activation of IgE and mast cells

C. Autoimmune mechanism

D. Drug hypersensitivity reaction

The most common cause of urticaria, also known as hives, is the activation of IgE and mast cells. In this process, specific allergens, which could be food, medications, insect stings, or environmental factors, lead to the cross-linking of IgE antibodies bound to mast cells. This results in the release of histamine and other mediators, causing the characteristic wheals, redness, and itching associated with urticaria. When mast cells degranulate, they release histamine, which increases vascular permeability, leading to fluid leakage from the blood vessels into the surrounding tissues, causing the swelling and redness. This immediate hypersensitivity reaction typically manifests within minutes after exposure to an allergen, highlighting IgE's central role in the pathogenesis of urticaria. While autoimmune mechanisms and drug hypersensitivity reactions can also cause urticaria, they are not the most common causes. Autoimmune urticaria can occur when the immune system mistakenly targets the body's own tissues, but this is less prevalent compared to allergic responses involving IgE. Drug hypersensitivity can also lead to urticaria, especially with certain medications, but it is also not as frequent as the IgE-mast cell activation route. Therefore, the role

8. What demographic is most commonly affected by Takayasu's arteritis?

A. Young Asian females

B. Middle-aged Caucasian males

C. Older African American males

D. Young Hispanic females

Takayasu's arteritis is a large vessel vasculitis that primarily affects the aorta and its major branches. The demographic most commonly affected by this condition is young Asian females, typically between the ages of 10 and 40. This association with young females in Asian populations is significant, as epidemiological studies have shown a higher prevalence of the disease in these demographics compared to others. The reason for the gender and racial predisposition is not entirely understood but may involve genetic and environmental factors. The other demographics mentioned do not align with the established patterns of Takayasu's arteritis. Middle-aged Caucasian males, older African American males, and young Hispanic females are less commonly affected, making them less relevant in the context of this specific vasculitis. The distinct population characteristics associated with Takayasu's arteritis emphasize the importance of recognizing specific demographic trends in clinical assessments and diagnoses of this condition.

SAMPLE

9. What is the primary treatment option for diffuse esophageal spasm?

A. Surgical resection

B. Supportive care

C. Proton pump inhibitors

D. Chemotherapy

Diffuse esophageal spasm (DES) is characterized by intermittent contractions of the esophagus that can lead to chest pain and dysphagia. The primary treatment option focuses on providing symptomatic relief and supportive care rather than a specific curative approach. Supportive care may include advice on dietary modifications, such as avoiding large meals and consuming softer foods, as well as the avoidance of triggers that may exacerbate symptoms, such as hot or cold foods. Additionally, medications that help relax the esophagus or decrease esophageal spasm, such as nitrates or calcium channel blockers, may be utilized. These approaches aim to reduce discomfort and improve the quality of life for patients rather than targeting a reversible lesion or specific pathology. Other options like surgical resection, proton pump inhibitors, and chemotherapy are not suitable as primary treatments for DES. Surgical options usually address structural problems, and DES does not respond to PPIs since it is not related to acid reflux. Chemotherapy is irrelevant in this context as it pertains to oncology and would not address the spasm experienced in the esophagus. Thus, supportive care remains the cornerstone of management for diffuse esophageal spasm.

10. What is a distinguishing feature of pemphigus foliaceus?

A. Presence of oral lesions

B. Deeper blisters that are less likely to rupture

C. Association with other autoimmune disorders

D. High likelihood of malignancy

Pemphigus foliaceus is primarily characterized by the presence of superficial blisters that are easily ruptured, resulting in crusted lesions, typically on the scalp, face, and trunk. Unlike its more severe counterpart, pemphigus vulgaris, pemphigus foliaceus does not present with oral lesions. This absence of oral involvement is a significant distinguishing feature between the two conditions. The condition is indeed associated with other autoimmune disorders, particularly in patients who may have underlying conditions such as systemic lupus erythematosus or rheumatoid arthritis. This association highlights the autoimmune nature of pemphigus foliaceus, where antibody production against desmogleins leads to the disruption of cell adhesion in the epidermis. While there may be concerns regarding the risk of malignancy in some autoimmune conditions, pemphigus foliaceus does not have a high likelihood of being associated with malignancy compared to other autoimmune blistering disorders. The condition typically does not present with deeper blisters, as seen in other forms of pemphigus, and oral lesions are also not a feature of this condition. The focus on its autoimmune associations makes the correct answer particularly relevant in distinguishing pemphigus foliaceus from other disease processes.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://usmlestep3.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE