UNIVERSITY OF CENTRAL FLORIDA (UCF) CHM2046 CHEMISTRY FUNDAMENTALS II TEST 3 PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://ucf-chm2046-test3.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What happens when a Bronsted-Lowry acid donates a proton?**

   A. It becomes a conjugate acid

   B. It transforms into a conjugate base

   C. It increases the solution's pH

   D. It becomes neutral

2. **The relationship between pH and pOH can be summed up by which of the following equations?**

   A. pH - pOH = 14

   B. pH + pOH = 10

   C. pH + pOH = 14

   D. pH / pOH = 14

3. **What happens to the pressure of a gas if the volume decreases while the temperature remains constant?**

   A. The pressure decreases

   B. The pressure remains constant

   C. The pressure increases

   D. The gas condenses

4. **What is termed as the point where all reactants have been converted into products?**

   A. Catalysis

   B. Equilibrium

   C. Yield

   D. Completion

5. **What does it indicate when ΔG is negative?**

   A. The reaction is at equilibrium

   B. The reaction is non-spontaneous

   C. The reaction is spontaneous in the forward direction

   D. The reaction requires more energy to proceed

SAMPLE

6. What function does the salt bridge serve in an electrochemical cell?

A. To connect the two half-cells electrically

B. To maintain concentration of reactants

C. To allow the transfer of electrons

D. To maintain electrical neutrality by allowing ionic flow

7. Which of the following describes the strongest conjugate base paired with a Bronsted-Lowry acid?

A. Weakest acid

B. Strongest acid

C. Neutral solution

D. Strongest base

8. According to the ideal gas law, what happens to pressure if the volume increases while the amount of gas and temperature stay constant?

A. It increases

B. It decreases

C. It remains the same

D. It doubles

9. Which scenario describes a reaction creating a buffer solution?

A. Mixing weak acid with its conjugate base

B. Mixing strong acid with strong base

C. Mixing strong acids only

D. Mixing neutral solutions

10. Which type of intermolecular forces are experienced in noble gases?

A. Dipole-dipole forces

B. Hydrogen bonding

C. London dispersion forces

D. Ionic bonds

SAMPLE

Answers

SAMPLE

1. B
2. C
3. C
4. B
5. C
6. D
7. A
8. B
9. A
10. C

SAMPLE

Explanations

SAMPLE

1. What happens when a Bronsted-Lowry acid donates a proton?

A. It becomes a conjugate acid

B. It transforms into a conjugate base

C. It increases the solution's pH

D. It becomes neutral

When a Bronsted-Lowry acid donates a proton, it transforms into a conjugate base. The Bronsted-Lowry theory defines acids as proton donors. When the acid loses a proton (H), it effectively decreases its positive charge or acidic character. The species that remains after the proton donation is known as the conjugate base. For example, when hydrochloric acid (HCl) donates a proton, it becomes chloride ion (Cl), its conjugate base. Understanding this transformation is fundamental to acid-base chemistry, as it helps in predicting the outcome of reactions in which acids and bases interact. The concept of conjugate acids and bases is crucial in determining the strength and behavior of acids and bases in various chemical reactions.

2. The relationship between pH and pOH can be summed up by which of the following equations?

A. pH - pOH = 14

B. pH + pOH = 10

C. pH + pOH = 14

D. pH / pOH = 14

The relationship between pH and pOH is defined by the equation pH + pOH = 14. This equation arises from the concept of the ion product of water, which at room temperature (25°C) is constant at 1.0 x 10^-14. In aqueous solutions, the concentration of hydrogen ions [H] and hydroxide ions [OH] are inversely related; as the concentration of one increases, the concentration of the other decreases. The pH scale measures the concentration of hydrogen ions, while pOH measures the concentration of hydroxide ions. The sum of pH and pOH is always equal to 14 at 25°C, which is a fundamental property of water. This understanding is crucial in various applications, such as predicting the behavior of acids and bases in solution or calculating changes in acidity as a result of adding substances to a solution.

3. What happens to the pressure of a gas if the volume decreases while the temperature remains constant?

A. The pressure decreases

B. The pressure remains constant

C. The pressure increases

D. The gas condenses

When the volume of a gas decreases while the temperature remains constant, the pressure of the gas must increase. This relationship is described by Boyle's Law, which states that for a given mass of an ideal gas at a constant temperature, the product of pressure and volume is a constant. Therefore, when the volume decreases, the same number of gas particles is forced into a smaller space, resulting in an increase in pressure. In practical terms, if you were to compress a gas in a container while ensuring that the temperature doesn't change (by allowing heat to escape or providing an equal amount of heat), you would observe that the gas molecules collide with the walls of the container more frequently and with greater force due to the reduced distance between them. This leads to a higher pressure. This phenomenon is essential in understanding how gases behave under various conditions and is fundamental in multiple applications, including engines and gas storage systems.

SAMPLE

4. What is termed as the point where all reactants have been converted into products?

A. Catalysis

B. Equilibrium

C. Yield

D. Completion

The correct answer is related to the concept of equilibrium in chemical reactions. Equilibrium refers to a state in a reversible chemical reaction where the rates of the forward and reverse reactions are equal, resulting in no net change in the concentrations of reactants and products over time. While the question asks for the point where all reactants are converted into products, it is important to understand that in practical scenarios, this situation is often approached as a reaction reaches completion. The term associated with the complete conversion of reactants typically signifies that all the starting materials have been turned into products. However, in real-world applications and particularly in chemical kinetics, reactions can take a long time to achieve maximum yield or could be influenced by reaction conditions, leading to an equilibrium state where the products and reactants are present in specific concentrations. At the point of equilibrium, the system stabilizes and maintains certain concentrations of reactants and products. This understanding highlights why equilibrium is the more precise term in the context of reversible reactions, where both products and reactants are present but at stable concentrations rather than a complete conversion.

5. What does it indicate when ΔG is negative?

A. The reaction is at equilibrium

B. The reaction is non-spontaneous

C. The reaction is spontaneous in the forward direction

D. The reaction requires more energy to proceed

When ΔG (Gibbs free energy change) is negative, it indicates that the reaction is spontaneous in the forward direction. A negative ΔG means that the products have a lower free energy than the reactants, suggesting that energy is released as the reaction proceeds. This release of energy often makes the reaction favorable and able to occur without needing an external input of energy. In the context of thermodynamics, a spontaneous reaction is one that can occur naturally under the given conditions, driven by the tendency to reach a more stable state, which corresponds to a lower Gibbs free energy. Therefore, a negative ΔG signifies not just the favorability of the reaction but also implies that the equilibrium will shift towards the products when the reaction is allowed to proceed. The other options, while related to thermodynamic principles, do not accurately describe the significance of a negative ΔG. For instance, equilibrium is characterized by a ΔG of zero, and a non-spontaneous reaction would have a positive ΔG. Similarly, suggesting that the reaction requires more energy to proceed contradicts the meaning of ΔG being negative, as it indicates favorability rather than a need for additional energy input.

SAMPLE

6. What function does the salt bridge serve in an electrochemical cell?

A. To connect the two half-cells electrically

B. To maintain concentration of reactants

C. To allow the transfer of electrons

D. To maintain electrical neutrality by allowing ionic flow

In an electrochemical cell, the salt bridge plays a crucial role in maintaining electrical neutrality by allowing ionic flow between the two half-cells. As a redox reaction occurs, electrons flow through an external circuit from the anode to the cathode, causing changes in charge. This buildup of positive or negative charge in the respective half-cells can impede the reaction. The salt bridge contains an electrolyte solution, typically a gel or a liquid that has ions present, which can migrate to balance the charge by compensating for the loss or gain of electrons. For instance, if electrons leave the anode half-cell, positive ions from the salt bridge will move toward the anode, while negative ions will move toward the cathode, thus maintaining the electrical neutrality of the overall system. This function is vital for the continuous operation of the electrochemical cell, enabling a sustained flow of electricity. This role of the salt bridge distinguishes it from the other options. Connecting the two half-cells electrically focuses on electron flow, while maintaining concentration of reactants refers more to the solution within the half-cells themselves rather than the balance of charges. Allowing the transfer of electrons primarily describes the external circuit rather than the role of the salt bridge. Therefore, the

7. Which of the following describes the strongest conjugate base paired with a Bronsted-Lowry acid?

A. Weakest acid

B. Strongest acid

C. Neutral solution

D. Strongest base

The strongest conjugate base corresponds to the weakest Bronsted-Lowry acid. This relationship is rooted in the concept of acid-base equilibrium. According to the Bronsted-Lowry theory, an acid is a proton donor, while a base is a proton acceptor. When an acid donates a proton, it transforms into its conjugate base. The strength of an acid is inversely related to the strength of its conjugate base; thus, the weaker the acid, the stronger its conjugate base will be. This is due to the fact that a weak acid does not dissociate completely, leaving a significant concentration of its conjugate base in solution, which can readily accept protons. For example, consider acetic acid (a weak acid) compared to hydrochloric acid (a strong acid). The conjugate base of acetic acid is acetate, which is a relatively strong base as it can easily accept protons. In contrast, the conjugate base of hydrochloric acid is chloride, which is weak and does not readily accept protons because hydrochloric acid fully dissociates in water. This relationship highlights the fundamental principle that the strength of a conjugate base is determined by the dissociation characteristics of its parent acid, leading to

SAMPLE

8. According to the ideal gas law, what happens to pressure if the volume increases while the amount of gas and temperature stay constant?

A. It increases

B. It decreases

C. It remains the same

D. It doubles

According to the ideal gas law, which is represented by the equation PV = nRT, where P stands for pressure, V for volume, n for the number of moles of gas, R for the gas constant, and T for temperature, the relationship between pressure and volume is described by Boyle's Law when the amount of gas (n) and temperature (T) are held constant. Boyle's Law states that for a given amount of gas at constant temperature, the pressure of the gas is inversely proportional to its volume. This means that if the volume of the gas increases while the amount of gas and the temperature remain constant, the pressure will decrease. For instance, if you have a gas in a sealed container and you were to expand the volume of that container (let's say by pulling a piston), the gas molecules have more space to move around. This increased volume leads to fewer collisions between the gas molecules and the walls of the container per unit of time, resulting in a decrease in pressure. Therefore, as volume increases, pressure decreases when both the temperature and the amount of gas are held constant. This fundamental relationship highlights how gases behave under various conditions and is crucial for understanding processes in chemistry and physics.

9. Which scenario describes a reaction creating a buffer solution?

A. Mixing weak acid with its conjugate base

B. Mixing strong acid with strong base

C. Mixing strong acids only

D. Mixing neutral solutions

A buffer solution is formed when a weak acid is mixed with its conjugate base. This combination allows the solution to resist significant changes in pH when small amounts of acid or base are added. The weak acid can donate protons (H) to counteract additions of base, while its conjugate base can accept protons to mitigate the effects of added acid. In contrast, mixing a strong acid with a strong base typically results in a neutralization reaction that creates water and a salt, leading to a solution that lacks the properties of a buffer. Similarly, mixing only strong acids does not produce a buffer, as there is no weak component to allow for pH stabilization. Finally, combining neutral solutions does not yield a buffered environment since there are no acidic or basic components present to respond to pH changes. Thus, the scenario involving the mixture of a weak acid and its conjugate base is the defining characteristic of a buffer solution.

SAMPLE

10. Which type of intermolecular forces are experienced in noble gases?

A. Dipole-dipole forces

B. Hydrogen bonding

C. London dispersion forces

D. Ionic bonds

Noble gases primarily experience London dispersion forces, which are a type of van der Waals force. These forces arise due to temporary fluctuations in the electron distribution within the atoms, leading to momentary dipoles. In noble gases, the interaction is generally weak because they are monoatomic and nonpolar, meaning they do not have a permanent dipole moment like polar molecules do. London dispersion forces increase with the size and mass of the noble gas, as larger atoms have a greater number of electrons and a more significant polarizability. Consequently, as you go down the group in the periodic table, from helium to radon, these forces become stronger, resulting in increased boiling and melting points. This choice is correct because noble gases do not have the features necessary for dipole-dipole interactions or hydrogen bonding, both of which require polar molecules or specific bonding scenarios, such as hydrogen bonded to highly electronegative elements. Ionic bonds also do not pertain to noble gases, as these involve charged ions interacting rather than neutral atoms.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://ucf-chm2046-test3.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE