U.S. PREVENTIVE SERVICES TASK FORCE (USPSTF) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://uspstf.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What does a grade of 'I' signify?

A. Insufficient evidence to assess the balance of benefits and harms of the service

B. The service is recommended, but with limitations

C. There is significant evidence against the service

D. The service is not recommended

2. If a woman has been screened for HIV during one pregnancy, what is the follow-up recommendation for subsequent pregnancies?

A. She should not be screened again

B. She should be rescreened

C. Rescreening is optional

D. She should wait 2 years between screenings

3. How are healthcare providers recommended to use USPSTF recommendations?

A. To decide on all treatments based on personal preference

B. To guide clinical decision-making regarding prevention and screening

C. To develop new medications

D. To create their own screening protocols without guidelines

4. What does the USPSTF recommend for unsensitized Rh (D)-negative women during 24-28 weeks' gestation?

A. Do not retest if the father is Rh-negative

B. Repeat Rh (D) antibody testing

C. Screening is not necessary

D. Immediate transfusion therapy

5. What diseases does the USPSTF promote vaccination against for adults?

A. Only measles and mumps

B. Influenza and pneumococcal disease

C. HIV and Hepatitis C

D. Only tetanus

SAMPLE

6. What criteria does the USPSTF use to assess new preventive services?

A. Effectiveness and popularity

B. Balance of benefits vs. harms

C. Cost alone

D. Time required for implementation

7. What is the recommended frequency for rescreening groups at very high risk for HIV infection?

A. Every 6 months

B. Annually

C. Every 2 years

D. Every 3 years

8. What impact does racial and ethnic background have on STI prevalence according to USPSTF findings?

A. There is no impact

B. Black and Hispanic persons have higher rates than white persons

C. White persons have the highest rates

D. Prevalence is equal among all groups

9. What BMI threshold does the USPSTF specify for obesity screening in adults?

A. 25 kg/m^2

B. 30 kg/m^2

C. 35 kg/m^2

D. 40 kg/m^2

10. What should clinicians consider when providing STI screening recommendations?

A. Only the age of patients

B. The communities they serve and local public health guidance

C. The gender of patients exclusively

D. Only previous patient history

SAMPLE

Answers

SAMPLE

1. A
2. B
3. B
4. B
5. B
6. B
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What does a grade of 'I' signify?

A. Insufficient evidence to assess the balance of benefits and harms of the service

B. The service is recommended, but with limitations

C. There is significant evidence against the service

D. The service is not recommended

A grade of 'I' from the U.S. Preventive Services Task Force signifies that there is insufficient evidence to assess the balance of benefits and harms of a preventive service. This grading indicates that the available data does not allow for a clear conclusion regarding whether the service is beneficial or harmful based on the current evidence. This classification suggests that further research is necessary to better evaluate the service's effectiveness and overall impact on patient outcomes. It does not imply that the service is harmful, nor does it recommend its use; rather, it highlights the need for more data before a definitive recommendation can be made. This is critical as it guides healthcare providers in their clinical decision-making, ensuring that they understand the uncertainties surrounding certain services.

2. If a woman has been screened for HIV during one pregnancy, what is the follow-up recommendation for subsequent pregnancies?

A. She should not be screened again

B. She should be rescreened

C. Rescreening is optional

D. She should wait 2 years between screenings

For women who have been screened for HIV during one pregnancy, the recommendation is to undergo rescreening in subsequent pregnancies. This guideline is in place because a woman's risk for HIV infection can change over time, depending on various factors such as new sexual partners or changes in risk behavior. The USPSTF emphasizes the importance of screening during each pregnancy, as early detection and treatment of HIV can significantly improve health outcomes for both the mother and the infant. Identifying HIV infection early allows for interventions that can reduce the transmission of the virus to the baby during childbirth, and enable effective management of the mother's health. By advocating for routine rescreening during each pregnancy, public health recommendations aim to ensure the best possible outcomes and significantly lower the risk of mother-to-child transmission of HIV.

SAMPLE

3. How are healthcare providers recommended to use USPSTF recommendations?

A. To decide on all treatments based on personal preference

B. To guide clinical decision-making regarding prevention and screening

C. To develop new medications

D. To create their own screening protocols without guidelines

Healthcare providers are recommended to use USPSTF recommendations to guide clinical decision-making regarding prevention and screening. The USPSTF offers evidence-based guidelines that are rigorously researched and evaluated, helping clinicians make informed decisions that are consistent with the best available evidence. These recommendations are intended to improve patient outcomes by assisting providers in determining which preventive services and screenings are appropriate for specific populations based on age, risk factors, and health history. Utilizing these guidelines helps standardize care, ensuring that patients receive appropriate preventive measures and screening tests in a timely manner. It also fosters a shared understanding among healthcare professionals regarding best practices, ultimately contributing to better health outcomes at the population level. By relying on the USPSTF recommendations, providers can avoid arbitrary decision-making and instead focus on evidence-based approaches that have been shown to be effective in improving health.

4. What does the USPSTF recommend for unsensitized Rh (D)-negative women during 24-28 weeks' gestation?

A. Do not retest if the father is Rh-negative

B. Repeat Rh (D) antibody testing

C. Screening is not necessary

D. Immediate transfusion therapy

The recommendation from the USPSTF for unsensitized Rh (D)-negative women during 24-28 weeks' gestation is to repeat Rh (D) antibody testing. This is essential as unsensitized Rh-negative women are at risk of developing antibodies against Rh-positive blood if their fetus is Rh-positive. By conducting this test at 24-28 weeks, healthcare providers can identify any potential sensitization before delivery. If sensitization occurs, it can lead to complications in future pregnancies, such as hemolytic disease of the newborn. Therefore, monitoring and testing during this critical period helps ensure appropriate intervention and management, should the mother show any signs of sensitization. The other options do not align with the established guidelines for prenatal care in this context. They either imply a lack of necessary follow-up or suggest irrelevant interventions that do not address the risk of Rh incompatibility in the pregnancy. Thus, regular screening and testing become crucial to safeguard both maternal and fetal health.

SAMPLE

5. What diseases does the USPSTF promote vaccination against for adults?

A. Only measles and mumps

B. Influenza and pneumococcal disease

C. HIV and Hepatitis C

D. Only tetanus

The correct answer focuses on the promotion of vaccination against influenza and pneumococcal disease for adults, reflecting the recommendations made by the U.S. Preventive Services Task Force (USPSTF). Vaccination is a critical preventive healthcare measure, especially for diseases that can lead to severe complications in adults. Influenza, or the flu, can cause significant morbidity and mortality, particularly among older adults and those with underlying health conditions. The annual flu vaccine is recommended to reduce the incidence of flu illness and its complications. Pneumococcal disease refers to infections caused by the Streptococcus pneumoniae bacteria, which can lead to pneumonia, meningitis, and other serious infections. Vaccination against pneumococcus is recommended for adults to prevent these potentially life-threatening diseases, especially in groups at higher risk, such as those over 65 or with certain chronic conditions. The other options highlight diseases that are either not primarily emphasized for adult vaccination or are involved in different preventive measures rather than vaccination. For example, while there are vaccines related to measles and mumps, the focus on adult preventive care typically centers around the most impactful vaccines, like those for influenza and pneumococcus. Therefore, the targeted vaccination for adults by the USPSTF reflects the need to

6. What criteria does the USPSTF use to assess new preventive services?

A. Effectiveness and popularity

B. Balance of benefits vs. harms

C. Cost alone

D. Time required for implementation

The USPSTF evaluates new preventive services primarily by assessing the balance of benefits versus harms. This criterion is essential because it ensures that any recommended preventive service demonstrates a clear advantage in improving health outcomes when weighed against potential negative effects or risks associated with that service. For example, a screening test may identify a disease early, which is a benefit, but if the test carries significant risks or leads to unnecessary invasive procedures without clear advantages, the overall assessment could be negative. The focus on benefits and harms is fundamental to the mission of the USPSTF, as it aims to promote evidence-based recommendations that genuinely enhance patient health and wellbeing. This systematic evaluation process is integral to making informed decisions about what preventive services to endorse, ensuring that the health interventions recommended are both safe and effective for the population. Other options, while potentially relevant in certain contexts, do not encapsulate the core principle the USPSTF utilizes for its assessments.

SAMPLE

7. What is the recommended frequency for rescreening groups at very high risk for HIV infection?

A. Every 6 months

B. Annually

C. Every 2 years

D. Every 3 years

The recommended frequency for rescreening groups at very high risk for HIV infection is annually. This recommendation is based on the understanding that individuals at high risk, such as those with multiple sexual partners, those involved in injection drug use, or individuals with a partner who is HIV positive, may have higher rates of new infections. Annual rescreening targets the need for timely identification and treatment of new infections, which is essential to reduce transmission within the community. The goal is to ensure that high-risk individuals are consistently aware of their HIV status, can receive appropriate preventive measures such as PrEP (pre-exposure prophylaxis), and can access care and treatment if they are found to be HIV positive. In contrast, less frequent rescreening intervals such as every 2, 3 years, or 6 months do not align with the urgency that comes with managing high-risk populations effectively, where the potential for new infections is significant. Hence, the annual screening recommendation is designed to closely monitor and manage the risk factors prevalent in these groups.

8. What impact does racial and ethnic background have on STI prevalence according to USPSTF findings?

A. There is no impact

B. Black and Hispanic persons have higher rates than white persons

C. White persons have the highest rates

D. Prevalence is equal among all groups

The statement that Black and Hispanic persons have higher rates of sexually transmitted infections (STIs) than white persons aligns with findings from various public health studies, including those evaluated by the U.S. Preventive Services Task Force (USPSTF). This conclusion is supported by data that show disparities in health outcomes among different racial and ethnic groups, highlighting the importance of addressing social determinants of health, access to care, and education on STI prevention and treatment. Understanding these disparities is crucial for developing targeted public health interventions and policy decisions aimed at reducing STI prevalence in higher-risk populations. Recognizing that certain groups experience a higher burden of STIs allows healthcare providers to tailor screening and preventive measures effectively, ensuring that resources are allocated based on need. The other options do not reflect the evidence appropriately. The suggestion that there is no impact oversimplifies the complex interactions between race, ethnicity, social factors, and health outcomes. Claiming that white persons have the highest rates or asserting equal prevalence among all groups contradicts the established epidemiological data, which shows significant differences in STI rates that are often correlated with socioeconomic factors, healthcare access, and cultural differences. Thus, understanding these differences among racial and ethnic backgrounds is vital for addressing the public health challenges posed by STIs.

SAMPLE

9. What BMI threshold does the USPSTF specify for obesity screening in adults?

A. 25 kg/m²

B. 30 kg/m²

C. 35 kg/m²

D. 40 kg/m²

The U.S. Preventive Services Task Force (USPSTF) specifies a body mass index (BMI) threshold of 30 kg/m² for obesity screening in adults. This threshold is significant because it aligns with established definitions of obesity, which categorize individuals with a BMI of 30 or higher as obese. Screening for obesity is essential as it allows for the identification of individuals who may be at increased risk for various health conditions, including diabetes, heart disease, and certain cancers. Using a threshold of 30 kg/m² helps healthcare providers focus their efforts on individuals who are likely to benefit from interventions, such as lifestyle modifications, counseling, or pharmacotherapy when necessary. By emphasizing this threshold, the USPSTF aims to support early detection and management of obesity, thereby contributing to improved overall health outcomes. The other thresholds mentioned would include individuals who do not meet the clinical definition of obesity. For instance, a BMI of 25 kg/m² is classified as overweight but not obese, which would not meet the screening criteria set forth by the USPSTF. Similarly, 35 kg/m² and 40 kg/m², while indicative of higher classes of obesity, are above the primary threshold used for screening all adults. By focusing on a BMI

10. What should clinicians consider when providing STI screening recommendations?

A. Only the age of patients

B. The communities they serve and local public health guidance

C. The gender of patients exclusively

D. Only previous patient history

When providing STI screening recommendations, it is essential for clinicians to consider the communities they serve and local public health guidance. This approach allows clinicians to tailor their screening recommendations based on the epidemiological data specific to the community, such as the prevalence of STIs in certain populations. Public health guidance is often derived from research and statistics that indicate which groups are at increased risk for STIs, based on various factors like demographic characteristics and regional health trends. Considering the broader community context helps ensure that screening efforts are not only relevant but also effective in reducing the incidence of STIs. For instance, in areas with high rates of certain STIs, clinicians might recommend more frequent screening or target specific populations that are more vulnerable. While age, gender, and previous patient history are factors that play a role in individual assessments, they should not be the sole focus in isolation. Effective STI screening is best achieved when it reflects the dynamics of the community and aligns with local health initiatives that aim to address public health challenges holistically. By integrating community-specific data and public health guidelines into their recommendations, clinicians can optimize screening practices and contribute to better health outcomes for their patients.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://uspstf.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE