
USMLE STEP 2 ANTIBIOTICS AND TREATMENTS PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://usmlestep2antibioticstreatments.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Identify a side effect unique to Vancomycin.

A. Fever and rash

B. Red man syndrome

C. Photosensitivity

D. Tendon rupture

2. What is the first-line treatment for community-acquired pneumonia (CAP)?

A. Azithromycin and ceftriaxone

B. Vancomycin and zosyn

C. Doxycycline or moxifloxacin

D. Amoxicillin and nitrofurantoin

3. When can seizure prophylaxis be stopped in a patient?

A. No seizures for 1 year

B. No seizures for 2 years

C. No seizures for 3 years

D. No seizures for 5 years

4. Which antibiotics are typically used for Community-Acquired Pneumonia (CAP)?

A. Azithromycin and Doxycycline

B. Azithromycin and Ceftriaxone

C. Bactrim and Tetracycline

D. Amoxicillin and Clindamycin

5. What is the first-line treatment for bacterial conjunctivitis in patients who do not wear contacts?

A. Topical erythromycin

B. Ciprofloxacin drops

C. Oral antibiotics

D. Warm compresses

SAMPLE

6. What is the typical mechanism of action for beta-lactam antibiotics?

A. Inhibition of folate synthesis

B. Disruption of protein synthesis

C. Inhibition of cell wall synthesis

D. Disruption of DNA replication

7. What condition presents with a painful red eye, hard on palpation, and does not react to light?

A. Allergic conjunctivitis

B. Acute angle-closure glaucoma

C. Uveitis

D. Corneal abrasion

8. What is the common side effect of aminoglycosides?

A. Gastrointestinal upset

B. Nephrotoxicity

C. Rash

D. Photosensitivity

9. If a patient has neutropenic fever without resolution after 72 hours, what is the next step?

A. Switch to oral antibiotics

B. Add antifungal therapy

C. Increase dosage of current antibiotics

D. Consult infectious disease specialist

10. What antibiotic is used to treat urinary tract infections due to E. coli?

A. Trimethoprim-sulfamethoxazole

B. Ciprofloxacin

C. Ampicillin

D. Nitrofurantoin

SAMPLE

Answers

SAMPLE

1. B
2. A
3. B
4. B
5. A
6. C
7. B
8. B
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. Identify a side effect unique to Vancomycin.

A. Fever and rash

B. Red man syndrome

C. Photosensitivity

D. Tendon rupture

Vancomycin is known for a specific side effect called Red Man Syndrome, which occurs due to rapid infusion of the drug. This syndrome is characterized by flushing, rash, and sometimes hypotension, predominantly affecting the upper body and neck. It results from the release of histamine triggered by the drug, leading to these vascular effects. While the side effects like fever and rash can occur with other medications as well, Red Man Syndrome is particularly associated with Vancomycin and is a distinct reaction that healthcare providers need to be aware of when administering the drug. The management involves slowing the infusion rate and prescriptive measures, such as administering antihistamines prior to administration, if necessary.

2. What is the first-line treatment for community-acquired pneumonia (CAP)?

A. Azithromycin and ceftriaxone

B. Vancomycin and zosyn

C. Doxycycline or moxifloxacin

D. Amoxicillin and nitrofurantoin

The first-line treatment for community-acquired pneumonia (CAP) in otherwise healthy adults typically involves using a macrolide, such as azithromycin, or a respiratory fluoroquinolone, along with a beta-lactam antibiotic like ceftriaxone. This combination effectively targets the most common bacterial pathogens responsible for CAP, including Streptococcus pneumoniae and atypical organisms such as Mycoplasma pneumoniae. In contrast, other options proposed are not standard first-line therapies for CAP. While doxycycline or moxifloxacin could serve as alternatives if the patient has specific factors, they are not as broadly recommended as the combination of azithromycin and ceftriaxone in treatment guidelines. The pairing of vancomycin and zosyn is more suitable for hospital-acquired pneumonia or severe cases involving multi-drug-resistant organisms, not the typical outpatient settings. Lastly, amoxicillin is commonly used in mild cases, but nitrofurantoin is primarily indicated for urinary tract infections rather than pneumonia. Therefore, the answer affirmed reflects current practices for effective treatment of CAP in community settings.

3. When can seizure prophylaxis be stopped in a patient?

A. No seizures for 1 year

B. No seizures for 2 years

C. No seizures for 3 years

D. No seizures for 5 years

Seizure prophylaxis can typically be considered for discontinuation after a patient has been seizure-free for a period of at least 2 years. This recommendation is based on clinical guidelines that suggest that the risk of seizures decreases significantly after this time frame. Patients who remain seizure-free for 2 years have a better prognosis for continued seizure freedom once medication is discontinued. Stopping seizure prophylaxis sooner than 2 years may increase the risk of recurrence since seizures can still occur, particularly in certain populations or types of seizure disorders, such as those with specific etiologies or a history of status epilepticus. Meanwhile, waiting longer than 2 years, such as 3 or 5 years, could be unnecessary in many cases, as the risk of seizure recurrence diminishes significantly after the 2-year mark. Thus, the 2-year seizure-free interval provides a balanced approach between minimizing unnecessary medication exposure and the risk of recurrent seizures.

SAMPLE

4. Which antibiotics are typically used for Community-Acquired Pneumonia (CAP)?

A. Azithromycin and Doxycycline

B. Azithromycin and Ceftriaxone

C. Bactrim and Tetracycline

D. Amoxicillin and Clindamycin

Community-acquired pneumonia (CAP) is primarily managed with antibiotics that cover the most common pathogens, such as Streptococcus pneumoniae, Mycoplasma pneumoniae, and Chlamydophila pneumoniae. Azithromycin is a macrolide antibiotic that provides excellent coverage against atypical bacteria often implicated in CAP, while ceftriaxone is a broad-spectrum cephalosporin that is particularly effective against certain strains of Streptococcus pneumoniae, especially in more severe cases or in patients needing hospitalization. Using azithromycin in combination with ceftriaxone allows for a dual approach: the macrolide addresses atypical organisms while ceftriaxone covers typical organisms effectively. This combination is particularly beneficial for patients with moderate to severe CAP or those with comorbid conditions. Alternative options, like the ones involving Bactrim, doxycycline, or clindamycin, may sometimes be appropriate, but they are not the first-line choices recommended in the latest guidelines for the treatment of CAP. Bactrim, for instance, is not generally used for CAP, while clindamycin is more reserved for specific scenarios, such as aspiration pneumonia. Hence, the combination of azithromycin and ceftriaxone stands out as

5. What is the first-line treatment for bacterial conjunctivitis in patients who do not wear contacts?

A. Topical erythromycin

B. Ciprofloxacin drops

C. Oral antibiotics

D. Warm compresses

The first-line treatment for bacterial conjunctivitis in patients who do not wear contact lenses is topical erythromycin. This antibiotic works effectively against the common pathogens responsible for bacterial conjunctivitis, such as Staphylococcus aureus and Streptococcus pneumoniae. Erythromycin is preferred due to its safety profile, effectiveness, and ease of use. In cases of bacterial conjunctivitis not associated with contact lens use, monitoring often shows that patients do not require more aggressive treatments, such as fluoroquinolone drops like ciprofloxacin, which are typically reserved for more severe infections or in cases where there is a risk of Pseudomonas aeruginosa, particularly in contact lens wearers. Oral antibiotics are not typically indicated for uncomplicated cases of conjunctivitis, as local topical treatment is sufficient. Warm compresses may provide symptomatic relief but do not address the underlying infection. Thus, the selection of topical erythromycin as the first-line treatment aligns well with both clinical guidelines and the nature of common bacterial conjunctivitis in this patient population.

SAMPLE

6. What is the typical mechanism of action for beta-lactam antibiotics?

A. Inhibition of folate synthesis

B. Disruption of protein synthesis

C. Inhibition of cell wall synthesis

D. Disruption of DNA replication

Beta-lactam antibiotics primarily function by inhibiting cell wall synthesis in bacteria. This class of antibiotics, which includes penicillins, cephalosporins, and carbapenems, targets the bacterial cell wall's key structural component known as peptidoglycan. Specifically, they bind to and inhibit penicillin-binding proteins (PBPs), which are essential for the cross-linking of peptidoglycan layers. This action weakens the cell wall, making it susceptible to osmotic pressure and ultimately leading to bacterial lysis and death. The effectiveness of beta-lactam antibiotics is particularly notable against actively dividing bacteria, as they rely heavily on their cell wall for structural integrity. Because of this mechanism, beta-lactams are generally considered bactericidal. Their action is crucial in treating various bacterial infections and highlights the importance of the bacterial cell wall as a target for antibiotic therapy.

7. What condition presents with a painful red eye, hard on palpation, and does not react to light?

A. Allergic conjunctivitis

B. Acute angle-closure glaucoma

C. Uveitis

D. Corneal abrasion

Acute angle-closure glaucoma is characterized by a painful red eye that typically has a hardness on palpation due to elevated intraocular pressure. This condition is marked by a rapid onset of symptoms, which can include severe eye pain, headache, nausea, and vomiting. The eye may appear red and is often non-reactive to light due to the increased pressure and potential damage to the optic nerve. In contrast, allergic conjunctivitis usually presents with redness and itching, along with tearing, but does not usually cause significant pain or a hard feeling in the eye. Uveitis, while it can present with redness and pain, often does not produce the palpably hard eye unless associated with significant intraocular inflammation or secondary complications. A corneal abrasion usually results in intense pain and photophobia, and the eye is not typically hard to touch. Thus, the characteristics of acute angle-closure glaucoma—such as the painful red eye, hard feeling upon palpation, and lack of light response—clearly delineate it as the correct diagnosis in this scenario.

8. What is the common side effect of aminoglycosides?

A. Gastrointestinal upset

B. Nephrotoxicity

C. Rash

D. Photosensitivity

Aminoglycosides, which include medications such as gentamicin, tobramycin, and amikacin, are primarily used to treat serious infections caused by Gram-negative bacteria. One of the well-recognized common side effects associated with aminoglycosides is nephrotoxicity. This refers to their potential to cause damage to the kidneys, leading to acute kidney injury. The nephrotoxic effect occurs due to the accumulation of aminoglycosides in renal tubular cells, where they can cause cellular damage and impair kidney function. Monitoring renal function during treatment is critical, especially since the risk of nephrotoxicity increases in patients who are also receiving other nephrotoxic agents or who have pre-existing kidney issues. Understanding this side effect is crucial for clinicians, as it highlights the importance of dosing adjustments and the need for regular kidney function assessments when administering aminoglycosides to patients.

SAMPLE

9. If a patient has neutropenic fever without resolution after 72 hours, what is the next step?

A. Switch to oral antibiotics

B. Add antifungal therapy

C. Increase dosage of current antibiotics

D. Consult infectious disease specialist

In cases of neutropenic fever, particularly when a patient has not shown improvement after 72 hours of broad-spectrum antibiotic therapy, the addition of antifungal therapy is warranted. Patients who are neutropenic are at significantly increased risk for invasive fungal infections, especially if their neutropenia is prolonged or if they have other risk factors such as recent chemotherapy or bone marrow disorders. The reasoning for adding antifungal therapy stems from the fact that common bacterial pathogens may not fully explain persistent febrile reactions in these patients. Invasive aspergillosis and candidiasis can occur in the context of neutropenia and often require a different therapeutic approach. The initiation of antifungal treatment after 72 hours of inadequate response to antibiotics is a recognized strategy aimed at covering possible fungal pathogens that may be complicating the clinical picture. Switching to oral antibiotics, increasing the dosage of current antibiotics, or merely consulting an infectious disease specialist may not address the undiagnosed possibility of a fungal infection that could be responsible for the patient's fever and lack of response. Therefore, the proper and timely introduction of antifungal agents is critical for optimizing patient outcomes in this context.

10. What antibiotic is used to treat urinary tract infections due to E. coli?

A. Trimethoprim-sulfamethoxazole

B. Ciprofloxacin

C. Ampicillin

D. Nitrofurantoin

Trimethoprim-sulfamethoxazole is a commonly used antibiotic for treating urinary tract infections (UTIs) caused by Escherichia coli, as it effectively targets and inhibits bacterial folic acid synthesis. This combination of trimethoprim and sulfamethoxazole works synergistically to enhance bacterial suppression, making it an effective choice for uncomplicated UTIs, which are often caused by E. coli. While ciprofloxacin is also effective against E. coli and is sometimes used for treating UTIs, it is generally reserved for cases where other treatments may fail or for complicated infections due to concerns about antibiotic resistance and potential side effects. Ampicillin can be used for UTI treatment, particularly when the causative organism is known to be susceptible, but it is less preferred for E. coli, which has shown increasing resistance to aminopenicillins. Nitrofurantoin is another option for treating UTIs, particularly in uncomplicated cases due to its specific action in the urinary tract; however, it is more effective against some strains of E. coli than others and is not the first choice in areas with high resistance rates. Given the context of standard treatment protocols for uncomplicated UTIs caused by E. coli, trimethoprim

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://usmlestep2antibioticstreatments.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE