USMLE STEP 2 CLINICAL KNOWLEDGE (CK) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://usmlestep2ck.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the primary treatment for whooping cough in patients?

A. Corticosteroids

B. Macrolides

C. Aspirin

D. Inhaled bronchodilators

2. Which of the following treatments is often recommended for managing conditions associated with high potassium levels during anesthesia?

A. Succinylcholine

B. Atracurium

C. Pancuronium

D. Dantrolene

3. What is a common complication in patients with sickle cell anemia affecting the hip?

A. Osteomyelitis

B. Fracture

C. Avascular necrosis

D. Arthritis

4. How is the stool pH affected in lactose intolerance?

A. High pH

B. Normal pH

C. Acidic pH

D. Alkaline pH

5. What is the common imaging method to diagnose developmental dysplasia of the hip in infants less than 4 months old?

A. Magnetic resonance imaging (MRI)

B. X-ray

C. Ultrasound (US)

D. Computed tomography (CT)

SAMPLE

6. Which electrolyte imbalance is commonly seen in patients with Conn's syndrome?

A. Hyperkalemia

B. Hypocalcemia

C. Hyponatremia

D. Hypokalemia

7. What is the first-line treatment for Legionella pneumonia?

A. Penicillin

B. Ciprofloxacin

C. Azithromycin

D. Tetracycline

8. Which of the following indicates a diagnosis of Fanconi anemia?

A. Deficiency in clotting factors

B. Autosomal dominant inheritance pattern

C. Development of skin changes and congenital anomalies

D. Increased white blood cell count

9. What characteristic rash is associated with mononucleosis when treated with ampicillin?

A. Maculopapular rash

B. Polymorphous rash

C. Urticarial rash

D. Scaly rash

10. What visible finding is associated with Mycoplasma pneumonia?

A. Erythema multiforme

B. Herpes labialis

C. Psoriasis

D. Urticaria

SAMPLE

Answers

SAMPLE

1. B
2. B
3. C
4. C
5. C
6. D
7. C
8. C
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. What is the primary treatment for whooping cough in patients?

A. Corticosteroids

B. Macrolides

C. Aspirin

D. Inhaled bronchodilators

The primary treatment for whooping cough, also known as pertussis, is macrolides. Pertussis is a bacterial infection caused by Bordetella pertussis, and macrolides, particularly azithromycin or erythromycin, are effective in eradicating the bacteria, especially when administered in the early stages of the illness. Using macrolides not only helps alleviate the symptoms more rapidly but also reduces transmission to others, making it a critical public health measure. While the infection can resolve on its own, timely antibiotic treatment is essential to minimize complications and protect vulnerable populations, such as infants who are more likely to experience severe illness. Other treatments, such as corticosteroids, aspirin, and inhaled bronchodilators, are not indicated for the management of whooping cough and do not address the underlying bacterial cause. Corticosteroids may be utilized in certain inflammatory conditions but are not beneficial for treating bacterial infections like pertussis. Aspirin is not indicated due to the risk of Reye's syndrome in children, and inhaled bronchodilators are more typically associated with the management of asthma or other obstructive lung diseases rather than infections like whooping cough.

2. Which of the following treatments is often recommended for managing conditions associated with high potassium levels during anesthesia?

A. Succinylcholine

B. Atracurium

C. Pancuronium

D. Dantrolene

The management of high potassium levels, especially in the context of anesthesia, emphasizes the careful selection of muscle relaxants due to their effects on the neuromuscular system. Atracurium is often the preferred choice because it is metabolized through a process called ester hydrolysis and non-specific plasma esterases. This metabolism does not rely on renal function, making it a safer choice for patients who may have renal impairment or elevated potassium levels. In cases of hyperkalemia, there's a concern for the use of certain neuromuscular blockers, particularly succinylcholine, which can lead to further potassium release from muscle cells, worsening hyperkalemia. Non-depolarizing muscle relaxants like atracurium are thus preferable as they have a more stable profile in managing neuromuscular blockade in high-potassium scenarios. Additionally, other agents, such as pancuronium, can prolong neuromuscular block and may have cardiac implications in patients with electrolyte imbalances. Dantrolene is primarily used to treat malignant hyperthermia and is not a standard treatment for hyperkalemia. Therefore, atracurium emerges as the most suitable option for patients experiencing high potassium levels during anesthesia.

SAMPLE

3. What is a common complication in patients with sickle cell anemia affecting the hip?

A. Osteomyelitis

B. Fracture

C. Avascular necrosis

D. Arthritis

In the context of sickle cell anemia, avascular necrosis is a significant complication, particularly affecting the hip joint. This condition occurs due to impaired blood flow to the bone, which can be a direct result of sickled red blood cells occluding small blood vessels. The femoral head of the hip is one of the most commonly affected areas due to its high vascular demand and limited collateral circulation. In patients with sickle cell anemia, the repeated episodes of vaso-occlusive crises can result in ischemia and subsequent necrosis of the bone tissue. The hip is notably vulnerable because it is a weight-bearing joint, and such damage can lead to pain, limited mobility, and eventually, joint deterioration. The onset of avascular necrosis often manifests as hip pain, which may be exacerbated by activity, and is diagnosed through imaging studies, such as MRI or X-rays. Other complications can arise in patients with sickle cell anemia, but avascular necrosis is particularly common and directly tied to the pathophysiological processes of the disease, distinguishing it from conditions such as osteomyelitis or arthritis, which are not inherently linked to the sickling phenomenon in the same manner. Fractures may occur due to bone weakness from avascular

4. How is the stool pH affected in lactose intolerance?

A. High pH

B. Normal pH

C. Acidic pH

D. Alkaline pH

In cases of lactose intolerance, the ability to properly digest lactose (a sugar found in milk and dairy products) is impaired due to a deficiency of lactase, the enzyme responsible for breaking down lactose into glucose and galactose. When lactose is not adequately digested, it passes into the large intestine where it is fermented by bacteria. This fermentation process produces short-chain fatty acids and gases, which can result in gastrointestinal symptoms such as bloating, diarrhea, and abdominal cramping. The presence of undigested lactose in the bowel leads to an increase in the production of acidic metabolites during fermentation, specifically lactic acid and other organic acids. This accumulation of acids subsequently lowers the stool pH, making it more acidic. An acidic stool pH is a classic finding in lactose intolerance. Therefore, the stool pH in lactose intolerance is typically acidic, reflecting the fermentation of undigested lactose by colonic bacteria. This mechanism underscores why patients with lactose intolerance may present with diarrhea that has a lower stool pH compared to those with normal lactose digestion.

SAMPLE

5. What is the common imaging method to diagnose developmental dysplasia of the hip in infants less than 4 months old?

A. Magnetic resonance imaging (MRI)

B. X-ray

C. Ultrasound (US)

D. Computed tomography (CT)

Ultrasound is the preferred imaging method for diagnosing developmental dysplasia of the hip in infants less than 4 months old because it is safe, does not involve radiation, and is particularly effective in evaluating the hip joint and surrounding soft tissues in this age group. The cartilaginous structures of the hip are not well visualized on X-ray in very young infants, making ultrasound a superior choice for accurate assessment. Ultrasound allows for dynamic assessment of the hip joint and can help identify hip dislocation or instability, which are critical in the early detection and management of developmental dysplasia of the hip. In older infants, X-ray may become more useful due to the increased ossification of bone structures, but for those under 4 months, ultrasound remains the gold standard for evaluation. Other imaging modalities, such as MRI and CT, are generally reserved for situations where more detailed anatomical evaluation is necessary or when other diagnoses need to be ruled out, as they are often more complicated, less accessible, or involve more risk compared to ultrasound for this particular application.

6. Which electrolyte imbalance is commonly seen in patients with Conn's syndrome?

A. Hyperkalemia

B. Hypocalcemia

C. Hyponatremia

D. Hypokalemia

Conn's syndrome, also known as primary hyperaldosteronism, is characterized by the excessive production of the hormone aldosterone by the adrenal glands. Aldosterone plays a crucial role in regulating electrolytes by promoting sodium reabsorption and potassium excretion in the renal tubules. In patients with Conn's syndrome, the increased levels of aldosterone lead to retention of sodium, which often results in elevated sodium levels in the bloodstream, but more importantly, it causes increased excretion of potassium. This process results in hypokalemia, or low levels of potassium in the body. Symptoms of hypokalemia can include muscle weakness, fatigue, and arrhythmias. Therefore, the common electrolyte imbalance associated with Conn's syndrome is hypokalemia, making it the correct answer in this context.

SAMPLE

7. What is the first-line treatment for Legionella pneumonia?

A. Penicillin

B. Ciprofloxacin

C. Azithromycin

D. Tetracycline

Legionella pneumonia is a type of atypical pneumonia caused by the bacterium Legionella pneumophila. The first-line treatment for this infection is typically a macrolide antibiotic, with azithromycin being a common choice. Macrolides are effective against Legionella because they can penetrate human cells, where Legionella often resides, and they inhibit protein synthesis in the bacteria. Given that Legionella does not respond well to beta-lactam antibiotics (like penicillin) due to its unique cell wall structure, and since tetracyclines (while potentially effective) are not the preferred first-line agents, azithromycin stands out as the most reliable and effective treatment option. Ciprofloxacin, a fluoroquinolone, could also be effective against Legionella and might be used in certain cases, especially when a patient has other contraindications to macrolides or has a complicated presentation; however, it is considered secondary to azithromycin as initial treatment. In summary, azithromycin is recommended due to its effectiveness, safety profile, and ability to target the intracellular nature of the organism, securing its status as the first-line therapy for Legionella pneumonia.

8. Which of the following indicates a diagnosis of Fanconi anemia?

A. Deficiency in clotting factors

B. Autosomal dominant inheritance pattern

C. Development of skin changes and congenital anomalies

D. Increased white blood cell count

Fanconi anemia is a rare genetic disorder characterized by a variety of clinical features, primarily due to defects in the DNA repair process. A hallmark of this condition is the presence of congenital anomalies and skin changes, which can include café-au-lait spots, short stature, skeletal abnormalities, and genital abnormalities. These manifestations arise because Fanconi anemia affects multiple organ systems during development. Patients with Fanconi anemia often present with aplastic anemia due to bone marrow failure, which can lead to increased susceptibility to infections and hematologic malignancies. The recognition of congenital anomalies, such as upper limb malformations or bone defects, combined with specific skin changes, supports the diagnosis of this condition. While deficiencies in clotting factors, autosomal dominant inheritance patterns, and increased white blood cell counts may be observed in various other conditions or contexts, they do not specifically characterize Fanconi anemia. The distinctive combination of congenital features and skin changes is crucial for making an accurate diagnosis of this disorder.

SAMPLE

9. What characteristic rash is associated with mononucleosis when treated with ampicillin?

A. Maculopapular rash

B. Polymorphous rash

C. Urticarial rash

D. Scaly rash

In the context of mononucleosis, which is typically caused by the Epstein-Barr virus (EBV), the administration of ampicillin can lead to a specific type of rash. The characteristic polymorphous rash that appears is often maculopapular in nature but can exhibit varied forms and features across different areas of the skin. This rash can be seen as part of an atypical reaction to the medication in patients with mononucleosis, and although some may confuse it with other forms of rashes due to different conditions or medications, it is notably associated with this presentation. The polymorphous nature indicates variability in the presentation, making it distinct in its appearance compared to other rashes. Other types of rashes listed in the options, while potentially related to different etiologies or hypersensitivity reactions, do not specifically correlate with the treatment of mononucleosis with ampicillin in the way that a polymorphous rash does. This distinguishes the polymorphous rash as the correct response in this scenario.

10. What visible finding is associated with Mycoplasma pneumonia?

A. Erythema multiforme

B. Herpes labialis

C. Psoriasis

D. Urticaria

Mycoplasma pneumoniae is recognized for its association with a variety of extrapulmonary manifestations, with erythema multiforme being one of the notable skin findings. Erythema multiforme is an immune-mediated condition characterized by the development of distinctive target lesions on the skin. This condition can occur following infections, and Mycoplasma pneumoniae is a well-known trigger. The pathophysiological mechanism involves cross-reactivity between Mycoplasma antigens and host tissues, leading to an inflammatory skin response. While the other conditions listed—herpes labialis, psoriasis, and urticaria—can all be associated with various triggers and conditions, they are not specifically linked to Mycoplasma pneumonia as erythema multiforme is. Herpes labialis is primarily a result of herpes simplex virus infection, psoriasis is a chronic autoimmune condition influenced by genetic and environmental factors, and urticaria is a reaction often related to allergens or irritants, but none of these are commonly caused by Mycoplasma pneumoniae infection.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://usmlestep2ck.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE