UWORLD SURGERY PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://uworldsurgery.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which condition best describes the need for total parenteral nutrition?

A. Ability to absorb food orally

B. Chronic renal failure

C. Intestinal obstruction

D. Non-functioning intestines

2. Following a gastrectomy, what syndrome may patients be at risk for developing?

A. Dumping syndrome

B. Parkinsonism

C. Hepatic encephalopathy

D. Gastroesophageal reflux disease

3. What is the most common cause of acute pancreatitis?

A. Gallstones

B. Alcohol abuse

C. Hyperlipidemia

D. Medications

4. What does the term "malrotation" refer to in pediatric surgery?

A. Improper alignment of intestine after surgery

B. Normal bowel rotation

C. Abnormal rotation of the bowel during fetal development

D. Dislocation of the bowel

5. What is a significant complication after a total hip arthroplasty indicated by sudden hemodynamic changes?

A. Deep vein thrombosis

B. Pneumonia

C. Acute renal failure

D. Pulmonary embolism

SAMPLE

6. What type of anticoagulant is commonly used in the perioperative setting?

A. Heparin

B. Aspirin

C. Warfarin

D. Clopidogrel

7. What is the most likely diagnosis for a man presenting with penile curvature and pain during erections?

A. Peyronie's disease

B. Hypogonadism

C. Erectile dysfunction

D. Priapism

8. What is the typical response in laboratory findings post-acute pancreatitis?

A. Increased white blood cell count

B. Elevated liver enzymes

C. Decreased hemoglobin

D. Elevated lipids

9. What surgical approach is preferred for a thickened or ulcerated gastric lesion?

A. Gastrectomy

B. Colostomy

C. Appendectomy

D. Duodenectomy

10. What is the first-line treatment for superficial venous thrombosis?

A. Anticoagulation therapy

B. NSAIDs and compression stockings

C. Elevation of the affected limb

D. Surgical removal of the vein

SAMPLE

Answers

SAMPLE

1. D
2. A
3. A
4. C
5. D
6. A
7. A
8. A
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. Which condition best describes the need for total parenteral nutrition?

A. Ability to absorb food orally

B. Chronic renal failure

C. Intestinal obstruction

D. Non-functioning intestines

Total parenteral nutrition (TPN) is a medical treatment that provides nutrients directly into the bloodstream, bypassing the digestive system entirely. This approach is necessary when the gastrointestinal (GI) tract cannot be used for nutrition, which is the case in conditions where the intestines are non-functional. In situations where the intestines cannot absorb nutrients due to severe diseases or dysfunction—such as short bowel syndrome, bowel obstruction, or during periods of bowel rest after surgery—TPN becomes crucial. It ensures that patients receive essential nutrients, including proteins, carbohydrates, fats, vitamins, and minerals, to maintain their nutritional status and support metabolic needs despite an inability to use the digestive tract effectively. The other conditions listed do not warrant the use of TPN to the same extent. For instance, the ability to absorb food orally contradicts the need for TPN, as patients who can eat usually do not require this intervention. Chronic renal failure might necessitate dietary modifications, but it doesn't inherently require TPN unless accompanied by other complications. Intestinal obstruction could sometimes allow for partial digestive function, and managing that condition may not always necessitate bypassing the intestines entirely. Therefore, it is the presence of non-functioning intestines that most accurately defines

2. Following a gastrectomy, what syndrome may patients be at risk for developing?

A. Dumping syndrome

B. Parkinsonism

C. Hepatic encephalopathy

D. Gastroesophageal reflux disease

After a gastrectomy, patients may develop dumping syndrome, which is a condition that occurs when food moves too quickly from the stomach into the small intestine. This rapid transit can lead to symptoms such as nausea, vomiting, diarrhea, cramping, and even palpitations or dizziness. The syndrome often results from the loss of the stomach's ability to regulate the pace at which food is released into the intestine, particularly after partial or total gastric resection. Patients may experience this syndrome especially with the consumption of high-sugar meals, as these can exacerbate the rapid emptying process. The physiological changes following gastrectomy, including the loss of gastric control and altered hormonal responses, significantly increase the risk for dumping syndrome, making it a relevant concern for patients who have undergone this surgical procedure. In contrast, other conditions listed do not have a direct connection to the effects of gastrectomy. Parkinsonism and hepatic encephalopathy arise from different etiologies unrelated to surgical alteration of the gastrointestinal tract. Gastroesophageal reflux disease can occur after various types of surgery but is not as commonly or directly associated with gastrectomy as dumping syndrome.

SAMPLE

3. What is the most common cause of acute pancreatitis?

A. Gallstones

B. Alcohol abuse

C. Hyperlipidemia

D. Medications

The most common cause of acute pancreatitis is gallstones. This is primarily because gallstones can obstruct the common bile duct or the pancreatic duct, leading to an inflammatory response within the pancreas. When a gallstone obstructs the pancreatic duct, it causes the accumulation of digestive enzymes within the pancreas, which can lead to autodigestion and inflammation, resulting in acute pancreatitis. While alcohol abuse is also a significant cause of acute pancreatitis, especially in certain populations, gallstones are statistically the leading cause overall. Hyperlipidemia and medications can also contribute to the development of pancreatitis but are less frequently responsible than gallstones. Understanding that gallstones can directly impact the anatomy and function of the biliary and pancreatic systems is crucial for grasping why they are the primary cause of this condition.

4. What does the term "malrotation" refer to in pediatric surgery?

A. Improper alignment of intestine after surgery

B. Normal bowel rotation

C. Abnormal rotation of the bowel during fetal development

D. Dislocation of the bowel

Malrotation refers to an abnormal rotation of the bowel during fetal development, which can lead to the displacement of the small intestine and potentially result in intestinal obstruction or volvulus after birth. During the normal development of the gastrointestinal tract, the intestines undergo a specific process of rotation and fixation. When this process is disrupted, the bowel can be improperly positioned, making it susceptible to complications such as twisting or strangulation. Understanding the concept of malrotation is crucial in pediatric surgery, as it is a condition that may require surgical intervention to correct and prevent severe gastrointestinal complications.

5. What is a significant complication after a total hip arthroplasty indicated by sudden hemodynamic changes?

A. Deep vein thrombosis

B. Pneumonia

C. Acute renal failure

D. Pulmonary embolism

A significant complication after total hip arthroplasty that is indicated by sudden hemodynamic changes is pulmonary embolism. Following surgery, particularly in orthopedic procedures like total hip arthroplasty, patients are at an increased risk of developing deep vein thrombosis (DVT) due to prolonged immobility and venous stasis. If a thrombus from the deep veins dislodges, it can travel to the pulmonary circulation, resulting in a pulmonary embolism. The symptoms of pulmonary embolism can manifest as a sudden onset of dyspnea, chest pain, tachycardia, and sometimes altered mental status, which may contribute to sudden hemodynamic changes such as hypotension or decreased oxygen saturation. This acute presentation differentiates pulmonary embolism from other complications that may not present so abruptly or are related to gradual physiological changes. In contrast, while deep vein thrombosis can lead to significant morbidity, it typically does not cause immediate hemodynamic instability until a pulmonary embolism occurs. Pneumonia and acute renal failure, while they are possible postoperative complications, are generally associated with gradual changes rather than sudden, drastic shifts in hemodynamics. Thus, understanding the pathophysiology and risk factors associated with these surgical procedures highlights pulmonary embol

SAMPLE

6. What type of anticoagulant is commonly used in the perioperative setting?

A. Heparin

B. Aspirin

C. Warfarin

D. Clopidogrel

Heparin is commonly used in the perioperative setting due to its rapid onset of action, controllable dosing, and the ability to rapidly reverse its effects. It is, therefore, particularly useful in managing patients who require anticoagulation close to or during surgical procedures. In addition, heparin can be given subcutaneously or intravenously, allowing for flexible administration depending on the clinical situation. In the perioperative context, heparin is often used for prevention of venous thromboembolism (VTE) in patients who are at increased risk, especially in situations where there might be prolonged immobility or surgical manipulation of venous structures. Its effectiveness is due in part to its ability to inhibit thrombin and factor Xa, thus preventing clot formation. Other anticoagulants, while they have their own indications, are less favored in the immediate perioperative setting. Warfarin, for instance, has a delayed onset and requires monitoring of INR levels, making it less suitable for immediate adjustment around the time of surgery. Aspirin and clopidogrel are antiplatelet agents rather than anticoagulants; although they can also have roles in certain surgical settings, they do not provide the same breadth of anticoagulation as

7. What is the most likely diagnosis for a man presenting with penile curvature and pain during erections?

A. Peyronie's disease

B. Hypogonadism

C. Erectile dysfunction

D. Priapism

The presentation of penile curvature and pain during erections is highly indicative of Peyronie's disease. This condition is characterized by the development of fibrous scar tissue in the penis, which results in curved and sometimes painful erections. The curvature can lead to difficulties during sexual intercourse and is often accompanied by discomfort or pain. Individuals with Peyronie's disease typically notice changes in the shape of their penis, and the condition can have significant psychological and relational impacts due to the associated symptoms. The etiology is often linked to trauma or injury to the penis, which leads to the formation of plaque that causes curvature. In this scenario, symptoms such as curvature and pain specifically during erections align with the typical clinical presentation of Peyronie's disease, making it the most likely diagnosis. Evaluation of the patient's history and the timing of symptoms can further support this diagnosis. Other conditions, like hypogonadism, erectile dysfunction, and priapism, have different presentations and would not typically cause the curvature and associated pain described in this situation.

SAMPLE

8. What is the typical response in laboratory findings post-acute pancreatitis?

A. Increased white blood cell count

B. Elevated liver enzymes

C. Decreased hemoglobin

D. Elevated lipids

In the context of acute pancreatitis, the typical laboratory finding is an increased white blood cell count. This response is part of the body's inflammatory reaction to the condition. Acute pancreatitis often triggers a systemic inflammatory response, leading to the activation of the immune system. As inflammation occurs, the body mobilizes white blood cells, particularly neutrophils, to the site of injury in the pancreas, causing a characteristic rise in their count in the bloodstream. The other options do not represent typical findings directly associated with acute pancreatitis. Elevated liver enzymes, for instance, could suggest liver involvement or biliary obstruction rather than pancreatitis alone. Decreased hemoglobin is not a standard finding in the acute phase of pancreatitis; while bleeding can occur, it is not a direct result of the pancreatic inflammation itself. Elevated lipids may be related to other conditions but are not a diagnostic or typical laboratory finding in acute pancreatitis. Therefore, the increased white blood cell count most accurately reflects the inflammatory response seen in acute pancreatitis.

9. What surgical approach is preferred for a thickened or ulcerated gastric lesion?

A. Gastrectomy

B. Colostomy

C. Appendectomy

D. Duodenectomy

The preferred surgical approach for a thickened or ulcerated gastric lesion is gastrectomy. This procedure involves the partial or total removal of the stomach and is indicated in cases of gastric cancer, severe peptic ulcers, or other significant gastric pathology that presents as thickening or ulceration of the gastric wall. Gastrectomy allows for both diagnostic evaluation and therapeutic intervention, enabling the surgeon to remove the lesion and any surrounding tissue that may be involved while minimizing the risk of leaving residual cancerous tissue. By excising the affected area, gastrectomy also addresses complications that arise from gastric lesions, such as obstruction or bleeding. Alternative procedures, such as colostomy, appendectomy, or duodenectomy, do not appropriately address gastric lesions. Colostomy pertains to the colon and is used primarily in the management of colonic diseases. Appendectomy is the surgical removal of the appendix and is irrelevant in gastric conditions. Duodenectomy, which involves removal of the duodenum, does not directly target the gastric lesion and is reserved for specific duodenal pathologies. Hence, none of these alternatives provide the necessary treatment for a thickened or ulcerated gastric lesion compared to gastrectomy.

SAMPLE

10. What is the first-line treatment for superficial venous thrombosis?

A. Anticoagulation therapy

B. NSAIDs and compression stockings

C. Elevation of the affected limb

D. Surgical removal of the vein

For superficial venous thrombosis, the first-line treatment involves the use of nonsteroidal anti-inflammatory drugs (NSAIDs) and compression stockings. This approach is effective in managing the pain and inflammation associated with the condition. NSAIDs help alleviate discomfort and reduce inflammation, making them an appropriate choice for symptomatic relief. Additionally, the application of compression stockings aids in improving venous return and can help minimize swelling in the affected limb. This dual approach addresses both the symptoms and the underlying vascular issue without the need for more invasive procedures or pharmacologic interventions that are typically reserved for more severe cases. In contrast, while anticoagulation therapy may be considered in certain scenarios, it is not routinely used for superficial thrombosis, particularly when there is no extension into deeper venous systems or significant risk of complications. Elevation of the affected limb may provide some benefit, but it is more of an adjunctive measure rather than a primary treatment. Surgical removal of the vein is rarely indicated for superficial venous thrombosis, as the condition often resolves with conservative management.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://uworldsurgery.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE