
# VASCULAR ACCESS BOARD CERTIFICATION PRACTICE EXAM (SAMPLE)

## STUDY GUIDE




## SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

**VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE**

https://vascularaccessboard.examzify.com

**Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.**

**ALL RIGHTS RESERVED.**

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

# Table of Contents


Copyright ..... 1

Table of Contents ..... 2

Introduction ..... 3

How to Use This Guide ..... 4

Questions ..... 5

Answers ..... 8

Explanations ..... 10

Next Steps ..... 16


SAMPLE

# Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

# How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

### 1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

### 2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

### 3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

### 4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

### 5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

### 6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

# Questions

SAMPLE

1. **How long can a transparent dressing with a semi-permeable membrane remain in place?**

   A. 24 to 48 hours

   B. 5 days

   C. 7 days

   D. 10 days

2. **What is an indication for employing an aquapheresis catheter?**

   A. Need for frequent blood draws

   B. Fluid management in diuretic-resistant CHF

   C. Blood transfusion purposes

   D. Administration of blood thinners

3. **How often should needleless connectors be changed during continuous IV infusion?**

   A. Every 24 hours

   B. Every 48 hours

   C. Every 96 hours

   D. Every 72 hours

4. **What does "checking for blood return" during flushing help determine?**

   A. The catheter is obstructed

   B. Infusion is successful

   C. The catheter is correctly positioned

   D. The saline solution is contaminated

5. **What should be regularly assessed as part of monitoring a central line?**

   A. Length of the catheter

   B. Frequency of dressing changes

   C. Signs of infection and catheter patency

   D. Quality of the intravenous fluid

SAMPLE

**6. What type of fluid can be administered through an umbilical venous catheter?**

A. Only blood products

B. Clear fluids only

C. TPN and medications

D. Only isotonic solutions

**7. What is the maximum period recommended for a non-tunneled central line placement?**

A. No more than 7 days

B. No more than 10-14 days

C. No more than 21 days

D. No more than 30 days

**8. What type of solution is typically used to flush vascular access devices?**

A. Heparin solution

B. Normal saline

C. Ringer's lactate

D. Dextrose solution

**9. What segment of the spine gives rise to the musculocutaneous nerve?**

A. C5-C7

B. C7-T1

C. C4-C6

D. C3-C5

**10. What assessment finding most commonly indicates a catheter-associated thrombosis?**

A. Redness around the insertion site

B. Swelling or pain in the affected arm or limb

C. Fever and chills

D. Increased heart rate

SAMPLE

# Answers

SAMPLE

1. C
2. B
3. C
4. C
5. C
6. C
7. B
8. B
9. A
10. B

SAMPLE

# Explanations

SAMPLE

## 1. How long can a transparent dressing with a semi-permeable membrane remain in place?

A. 24 to 48 hours

B. 5 days

C. 7 days

D. 10 days

*The correct response is based on the guidelines for the use of transparent dressings with semi-permeable membranes, which are designed to maintain a moist environment while allowing for permeability to gases and moisture. These dressings are effective for covering vascular access devices and can typically remain in place for a duration of up to 7 days. The 7-day lifespan is supported by evidence indicating that such dressings effectively reduce the risk of infection while providing adequate protection and visibility of the insertion site. This duration also aligns with best practice recommendations aimed at balancing the need for access site surveillance with the potential for skin irritation or complications that may arise from longer application. While other timeframes mentioned may be part of recommended practices for different types of dressings or specific scenarios not related to transparent semi-permeable membranes, the 7-day limit is widely accepted for these types of dressings in order to ensure ongoing effectiveness and safety.*

## 2. What is an indication for employing an aquapheresis catheter?

A. Need for frequent blood draws

B. Fluid management in diuretic-resistant CHF

C. Blood transfusion purposes

D. Administration of blood thinners

*An aquapheresis catheter is specifically designed for managing fluid overload in conditions such as congestive heart failure (CHF), particularly when traditional diuretics are ineffective or have limited efficacy. In diuretic-resistant CHF, the body retains excess fluid, leading to symptoms like edema and shortness of breath. Employing an aquapheresis catheter allows for the continuous removal of excess fluid, thereby alleviating these symptoms and improving the patient's overall circulatory status. This method is particularly beneficial in chronic cases where patients struggle with volume overload, as it directly targets the underlying issue of fluid retention rather than just masking the symptoms. The use of an aquapheresis catheter integrates a form of ultrafiltration, which selectively removes excess fluid without significantly disrupting electrolyte balance or causing the complications often associated with more invasive procedures. In contrast, the other options may pertain to procedures that do not require the specialized use of an aquapheresis catheter. Frequent blood draws, for instance, typically utilize standard venous access methods. Blood transfusions and the administration of anticoagulants commonly rely on dedicated lines or peripheral access rather than an aquapheresis catheter, which is focused solely on fluid removal in specific clinical situations.*

SAMPLE

## 3. How often should needleless connectors be changed during continuous IV infusion?

A. Every 24 hours

B. Every 48 hours

C. Every 96 hours

D. Every 72 hours

*Needleless connectors are critical components in intravenous (IV) therapy, providing a way to administer fluids and medications while minimizing the risk of needle-stick injuries and contamination. The appropriate frequency for changing needleless connectors during continuous IV infusion is based on guidelines that prioritize patient safety and infection control. Changing needleless connectors every 96 hours aligns with recommendations provided by various health organizations, which recognize that while these connectors may become contaminated due to frequent use, they can remain functional and safe for longer periods when used appropriately. This timeframe takes into account the importance of maintaining a balance between ensuring cleanliness and reducing the risk of unnecessary disconnections that could lead to complications. It is essential to note that certain factors may necessitate more frequent changes of needleless connectors, such as any signs of visible contamination, patient's clinical condition, or institutional protocols influenced by specific patient populations. However, in a standard continuous IV infusion scenario without complications, the 96-hour guideline is a widely accepted practice. In summary, the recommendation to change needleless connectors every 96 hours is grounded in a goal of optimizing care while adhering to infection control practices, ensuring the connectors remain functions safe for the duration of IV therapy.*

## 4. What does "checking for blood return" during flushing help determine?

A. The catheter is obstructed

B. Infusion is successful

C. The catheter is correctly positioned

D. The saline solution is contaminated

*"Checking for blood return" during the flushing of a vascular catheter is a critical procedure that helps determine whether the catheter is correctly positioned within the vascular system. When a healthcare professional observes blood return during this process, it indicates that the catheter tip is situated within a blood vessel, allowing for proper circulation. This vital assessment serves to confirm that the pathway for delivering medications or fluids is functional and that the catheter has not migrated into inappropriate locations, such as tissue or a non-vascular space. This confirmation is crucial for ensuring effective treatment and avoiding complications, as a catheter that is not in the correct position might lead to insufficient delivery of fluids or medications, potentially causing harm to the patient. Therefore, the presence of blood return is a reliable indicator of correct catheter placement and functionality, which is essential for safe and effective patient care.*

SAMPLE

## 5. What should be regularly assessed as part of monitoring a central line?

A. Length of the catheter

B. Frequency of dressing changes

C. Signs of infection and catheter patency

D. Quality of the intravenous fluid

*Regular assessment of signs of infection and catheter patency is crucial in monitoring a central line. This practice is essential because central venous catheters (CVCs) are associated with risks such as bloodstream infections and thrombosis. Monitoring for signs of infection involves looking for symptoms like redness, swelling, warmth, or discharge at the insertion site, which could indicate infection. Additionally, assessing catheter patency ensures that the catheter is functioning properly, allowing for effective delivery of medications and fluids, and reducing the risk of complications such as occlusion. While other factors, such as the frequency of dressing changes and the quality of intravenous fluids, are important in overall patient care, they do not directly relate to the immediate risks associated with the catheter itself. The length of the catheter might indicate placement issues but is not a routine factor for ongoing monitoring. In contrast, regularly checking for infection and ensuring that the catheter is patent directly influences patient safety and treatment efficacy.*

## 6. What type of fluid can be administered through an umbilical venous catheter?

A. Only blood products

B. Clear fluids only

C. TPN and medications

D. Only isotonic solutions

*An umbilical venous catheter (UVC) is designed to provide access to the circulatory system of neonates for the administration of various types of fluids, and option C highlights two primary uses for this type of catheter: Total Parenteral Nutrition (TPN) and medications. One of the key roles of a UVC is to facilitate nutritional support, particularly in premature infants or those who cannot take feeds orally. TPN—which includes essential nutrients such as carbohydrates, proteins, fats, vitamins, and minerals—can be directly infused into the bloodstream through a UVC, ensuring that the neonate receives appropriate nutrition while bypassing the gastrointestinal tract. Additionally, a UVC can be used to administer various medications, allowing for rapid delivery into the systemic circulation, which is crucial for the management of different medical conditions commonly faced by neonates. This dual capability makes the UVC a vital tool in neonatal care. While other options may contain aspects of potential fluid administration, they do not capture the full range of capabilities afforded by the UVC. Therefore, recognizing that TPN and medications can be effectively delivered through an umbilical venous catheter underscores the versatility and importance of this vascular access method in neonatal medicine.*

SAMPLE

## 7. What is the maximum period recommended for a non-tunneled central line placement?

A. No more than 7 days

B. No more than 10-14 days

C. No more than 21 days

D. No more than 30 days

*The recommendation for the duration of non-tunneled central venous catheters is typically no more than 10-14 days. This timeframe is based on evidence that longer dwelling times can increase the risk of complications, including infections and thrombosis. The shorter duration helps to minimize these risks while still allowing for appropriate use of the catheter for necessary therapies, such as intravenous medications or fluid administration. Unsurprisingly, extending the use beyond 14 days can lead to a significantly higher incidence of complications without substantial benefit, which is why guidelines firmly advocate for the 10-14 day maximum for non-tunneled lines. This ensures that patients receive the necessary care while maintaining safety as a priority.*

## 8. What type of solution is typically used to flush vascular access devices?

A. Heparin solution

B. Normal saline

C. Ringer's lactate

D. Dextrose solution

*Normal saline is the most commonly used solution to flush vascular access devices due to its isotonic properties, which minimize the risk of fluid shifts and cellular damage. This solution helps to maintain the patency of the vascular access device by preventing the formation of blood clots and ensuring that the lumen remains clear for medication administration or fluid infusions. Because it closely matches the body's electrolyte balance and osmolarity, normal saline is safe and effective for maintaining vascular access devices. It is easy to prepare and store, making it practical for clinical use. This solution also helps to dilute any potentially harmful substances that may be left in the catheter or to remove any debris that could compromise the functionality of the device. While heparin solution is sometimes used in certain circumstances to prevent clot formation in long-term catheters, it is not typically used for routine flushing due to its anticoagulant properties and the need for careful monitoring. Ringer's lactate and dextrose solutions have specific clinical applications but are not suitable for flushing vascular access devices in general practice, as they alter electrolyte balance and do not effectively prevent clotting in the same way normal saline does.*

## 9. What segment of the spine gives rise to the musculocutaneous nerve?

A. C5-C7

B. C7-T1

C. C4-C6

D. C3-C5

*The musculocutaneous nerve is primarily formed from the spinal segments C5, C6, and C7. These segments are involved in the emergence of the nerve from the brachial plexus, which innervates the muscles of the anterior compartment of the arm, notably the biceps brachii and brachialis, as well as providing sensory innervation to part of the forearm. When considering the spinal segments that give rise to the musculocutaneous nerve, C5, C6, and C7 provide the necessary nerve roots for its formation. The roots originate from the upper trunk of the brachial plexus and contribute to the peripheral nerve that is essential for arm function. The other spinal segment ranges provided do not correlate with the musculocutaneous nerve. C7-T1 pertains to other nerves in the plexus, while C4-C6 and C3-C5 include segments that do not contribute to the musculocutaneous nerve pathway. Thus, it establishes that C5-C7 is indeed the correct choice for the segments that are responsible for the musculocutaneous nerve.*

SAMPLE

**10. What assessment finding most commonly indicates a catheter-associated thrombosis?**

A. Redness around the insertion site

B. Swelling or pain in the affected arm or limb

C. Fever and chills

D. Increased heart rate

*Swelling or pain in the affected arm or limb is the assessment finding that most commonly indicates a catheter-associated thrombosis. This is because a thrombosis leads to impaired venous return and can cause venous obstruction, resulting in noticeable swelling in the affected area. Additionally, the presence of pain occurs due to inflammation and pressure on surrounding tissues as the thrombus develops. The other signs mentioned may suggest different complications. Redness around the insertion site is often more related to infection or phlebitis rather than thrombosis. Fever and chills typically indicate an infectious process rather than a thrombotic event. Increased heart rate can be a generalized response to many factors, including pain or anxiety, and is not specific to catheter-associated thrombosis. Thus, the combination of localized swelling and pain is the most reliable indicator for identifying a potential catheter-associated thrombosis.*

SAMPLE

# Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://vascularaccessboard.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE