VESTIBULAR SYSTEM PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://vestibularsystem.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 9

Explanations 11

Next Steps 17

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What does vestibular evoked myogenic potential testing assess, and what do cVEMP and oVEMP reflect?

A. VEMP assesses semicircular canal function; cVEMP reflects saccular and inferior vestibular nerve function; oVEMP reflects utricular and superior vestibular nerve function

B. VEMP assesses auditory pathways; cVEMP reflects cochlear; oVEMP reflects vestibular

C. VEMP assesses otolithic pathway function; cVEMP reflects saccular and inferior vestibular nerve function; oVEMP reflects utricular and superior vestibular nerve function

D. VEMP assesses visual pathways; cVEMP reflects optic nerve function; oVEMP reflects oculomotor nerve

2. In unilateral VOR damage, compensation is achieved through which mechanism?

A. Head movements

B. Complete recovery

C. Increased pursuit

D. Nerve regrowth

3. Vestibular rehabilitation addresses which domains to improve function?

A. Hearing and language development

B. Gaze stabilization, habituation, and balance/gait training

C. Visual acuity and color perception

D. Cardiovascular endurance only

4. Distinguish the typical presentations of Menière disease versus vestibular schwannoma.

A. Both conditions typically present with progressive bilateral SNHL.

B. Menière disease has constant vertigo and stable hearing, while schwannoma has fluctuating hearing.

C. Menière: episodic vertigo with fluctuating low-frequency SNHL and tinnitus; vestibular schwannoma: progressive unilateral SNHL with vestibular symptoms.

D. Schwannoma presents with hearing loss but no vestibular symptoms.

SAMPLE

5. Where are the maculae located and what is their role in vestibular function?

A. Otoliths are located in the utricle and saccule; maculae detect linear accelerations and head tilt via otoliths overlying maculae.

B. Maculae are located in the cochlea; they detect sound.

C. Maculae are located in the semicircular canals; they detect angular velocity.

D. Maculae are located in the vestibulocochlear nerve; they transmit auditory signals.

6. Which nerve is associated with cVEMP responses?

A. Superior vestibular nerve

B. Cochlear nerve

C. Inferior vestibular nerve

D. Facial nerve

7. The cupula serves as a receptor-level structure because it does what?

A. Extends into the endolymph and bends hair cells when endolymph moves

B. Houses hair cells directly

C. Detects gravity

D. Produces endolymph

8. Differentiated pattern of nystagmus between canalolithiasis and cupulolithiasis in BPPV.

A. Canalolithiasis: delayed-onset geotropic nystagmus; cupulolithiasis: immediate-onset persistent nystagmus with sustained vertigo due to debris adheres to cupula.

B. Canalolithiasis: immediate-onset vertical nystagmus; cupulolithiasis: delayed horizontal nystagmus.

C. Canalolithiasis: immediate-onset persistent nystagmus; cupulolithiasis: delayed-onset geotropic nystagmus.

D. Canalolithiasis and cupulolithiasis have identical nystagmus patterns.

9. Which cerebellar regions are involved in modulating the vestibulo-ocular reflex and facilitating vestibular compensation?

A. Cerebellar cortex

B. Cerebral cortex

C. Basal ganglia

D. Flocculus and nodulus

SAMPLE

10. Which examination maneuver is commonly used to diagnose BPPV?

A. Romberg test

B. Head impulse test

C. Dix-Hallpike maneuver

D. Caloric test

SAMPLE

Answers

SAMPLE

1. C
2. A
3. B
4. C
5. A
6. C
7. A
8. A
9. D
10. C

SAMPLE

Explanations

SAMPLE

1. What does vestibular evoked myogenic potential testing assess, and what do cVEMP and oVEMP reflect?

A. VEMP assesses semicircular canal function; cVEMP reflects saccular and inferior vestibular nerve function; oVEMP reflects utricular and superior vestibular nerve function

B. VEMP assesses auditory pathways; cVEMP reflects cochlear; oVEMP reflects vestibular

C. VEMP assesses otolithic pathway function; cVEMP reflects saccular and inferior vestibular nerve function; oVEMP reflects utricular and superior vestibular nerve function

D. VEMP assesses visual pathways; cVEMP reflects optic nerve function; oVEMP reflects oculomotor nerve

VEMP testing taps the otolithic pathways of the vestibular system, focusing on how the otolith organs respond to stimulation. The cervical VEMP is recorded from the sternocleidomastoid muscle and primarily reflects saccular input carried by the inferior vestibular nerve, meaning it tells us about saccular function and its brainstem connections that influence neck muscles. The ocular VEMP is recorded from eye muscles and mainly reflects utricular input via the superior vestibular nerve, giving information about utricular function and its vestibulo-ocular pathways. So, VEMP assesses otolithic function rather than semicircular canal function, and cVEMP points to the saccule and inferior nerve pathway, while oVEMP points to the utricle and superior nerve pathway.

2. In unilateral VOR damage, compensation is achieved through which mechanism?

A. Head movements

B. Complete recovery

C. Increased pursuit

D. Nerve regrowth

When one vestibular system is damaged, the brain adapts by recalibrating how gaze is stabilized and by using other sensory cues to substitute for the lost input. A key driver of this compensation is head movement. Repeated head motion engages the intact vestibular pathways and drives plastic changes in the cerebellum and vestibular nuclei, which adjust the VOR gain and help the eyes generate appropriate compensatory eye movements (catch-up saccades) during head turns. Over time, this learning, combined with greater reliance on visual input and neck proprioception, improves gaze stability despite the unilateral loss. The other options don't explain the mechanism as well: complete recovery is not the mechanism itself (it can occur variably, but compensation is about the process, not a guaranteed full return). Increased pursuit is not the main compensatory strategy for VOR loss, since smooth pursuit is a different system used for tracking moving targets. Nerve regrowth is not how central compensation occurs in adults—the compensation relies on neural plasticity and sensory substitution, not regeneration of the damaged nerve.

SAMPLE

3. Vestibular rehabilitation addresses which domains to improve function?

A. Hearing and language development

B. Gaze stabilization, habituation, and balance/gait training

C. Visual acuity and color perception

D. Cardiovascular endurance only

The main idea is that vestibular rehabilitation focuses on three areas that directly support how the vestibular system helps you function: keeping vision stable during head movement, reducing dizziness through controlled exposure, and improving balance and walking. Gaze stabilization is about training the eyes to stay focused on a target when the head moves. This relies on the vestibulo-ocular reflex, so exercises that pair head and eye movements help you see clearly during daily tasks like walking down the street or reading while riding in a car. Habituation involves repeated exposure to movements or positions that provoke dizziness. Over time, the nervous system adapts, and symptoms lessen, making those movements more tolerable. Balance and gait training build safer, steadier posture and walking. This encompasses static balance tasks, dynamic challenges, and walking with varying surfaces or dual tasks, all aimed at reducing falls and increasing functional mobility. The other options don't fit as well: hearing and language aren't the primary aims of vestibular rehab, though some patients may have related issues; visual acuity and color perception aren't improved by vestibular therapy (vision is used as a cue, but acuity isn't increased); and while overall fitness matters, cardiovascular endurance alone doesn't address the specific domains targeted by vestibular rehabilitation.

4. Distinguish the typical presentations of Menière disease versus vestibular schwannoma.

A. Both conditions typically present with progressive bilateral SNHL.

B. Menière disease has constant vertigo and stable hearing, while schwannoma has fluctuating hearing.

C. Menière: episodic vertigo with fluctuating low-frequency SNHL and tinnitus; vestibular schwannoma: progressive unilateral SNHL with vestibular symptoms.

D. Schwannoma presents with hearing loss but no vestibular symptoms.

A clear way to distinguish these conditions is by how they affect hearing and balance over time. Menière disease typically presents with episodes of vertigo that come and go, paired with fluctuating low-frequency sensorineural hearing loss and tinnitus. The hearing loss isn't steady; it waxes and wanes with each attack, especially affecting the low frequencies early on. Vestibular schwannoma, on the other hand, usually shows a progressive, unilateral sensorineural hearing loss that worsens over time, along with vestibular symptoms such as imbalance or vertigo related to the tumor's impact on the vestibular nerve. The hearing loss tends to be gradual and persistent rather than fluctuating. So the best summary is: Menière disease features episodic vertigo with fluctuating low-frequency hearing loss and tinnitus, while vestibular schwannoma features progressive unilateral hearing loss with accompanying vestibular symptoms.

SAMPLE

5. Where are the maculae located and what is their role in vestibular function?

A. Otoliths are located in the utricle and saccule; maculae detect linear accelerations and head tilt via otoliths overlying maculae.

B. Maculae are located in the cochlea; they detect sound.

C. Maculae are located in the semicircular canals; they detect angular velocity.

D. Maculae are located in the vestibulocochlear nerve; they transmit auditory signals.

Maculae are the sensory patches that line the otolith organs of the inner ear—the utricle and saccule. Their purpose is to sense linear motion and head position relative to gravity. They sit beneath a gelatinous layer that contains otoliths, tiny calcium crystals. When you move in a straight line or tilt your head, gravity causes these crystals to tug on the gelatin and bend the hair cells of the macula. This bending converts mechanical movement into neural signals that travel via the vestibular nerve to the brain, providing information about linear acceleration and tilt. This data is essential for balance and stabilizing gaze, working alongside the semicircular canals, which detect angular (rotational) motion. The cochlea is involved in hearing, and the vestibulocochlear nerve transmits these sensory signals rather than housing the sensory cells themselves.

6. Which nerve is associated with cVEMP responses?

A. Superior vestibular nerve

B. Cochlear nerve

C. Inferior vestibular nerve

D. Facial nerve

cVEMP responses reflect a vestibulocollic reflex that starts with the saccule, an otolith organ, sending input through the inferior vestibular nerve. When a loud sound or tone burst stimulates the saccule, the signal travels via the inferior vestibular nerve to the brainstem vestibular nuclei and then down to the sternocleidomastoid muscle through the spinal accessory nerve. The muscle response is what we record in cVEMP testing. This is why the inferior vestibular nerve is the nerve associated with cVEMP. The superior vestibular nerve carries input from the utricle and semicircular canals, not primarily the saccule. The cochlear (auditory) nerve handles hearing, and the facial nerve is motor to facial muscles, not part of this reflex pathway.

7. The cupula serves as a receptor-level structure because it does what?

A. Extends into the endolymph and bends hair cells when endolymph moves

B. Houses hair cells directly

C. Detects gravity

D. Produces endolymph

The cupula is the movable, gelatinous lid of the crista ampullaris that extends into the endolymph and deflects when the endolymph moves. When the head rotates, inertia causes the endolymph to lag, pushing against the cupula and bending the hair cells' stereocilia. That deflection changes the receptors' ion flow, altering neurotransmitter release and sending signals to the vestibular nerve. This is how mechanical motion is translated into neural information, which is the essence of a receptor-level structure. The hair cells themselves sit in the sensory epithelium, not inside the cupula; gravity detection is done by the otolith organs, and endolymph is produced by other vascular channels, not by the cupula.

SAMPLE

8. Differentiated pattern of nystagmus between canalolithiasis and cupulolithiasis in BPPV.

A. Canalolithiasis: delayed-onset geotropic nystagmus; cupulolithiasis: immediate-onset persistent nystagmus with sustained vertigo due to debris adheres to cupula.

B. Canalolithiasis: immediate-onset vertical nystagmus; cupulolithiasis: delayed horizontal nystagmus.

C. Canalolithiasis: immediate-onset persistent nystagmus; cupulolithiasis: delayed-onset geotropic nystagmus.

D. Canalolithiasis and cupulolithiasis have identical nystagmus patterns.

A key distinction here is where the debris causing BPPV is located and how that affects the velocity and duration of the eye movements after you change head position. In canalolithiasis, the otoconia are free-floating in the canal. When a provocative position is assumed, the endolymph flow takes a moment to start moving the particles, so there is a brief delay before symptoms appear. The nystagmus is geotropic (toward the ground ear) and short-lived because the particles move with the flow and then settle, causing only a transient response. In cupulolithiasis, debris sticks to the cupula itself. Gravity continuously deflects the cupula as the head is held in a provocative position, so there is little to no delay and the vertigo with nystagmus can be persistent, lasting as long as the position is maintained. Thus, the description that matches this pattern is canalolithiasis with delayed-onset geotropic nystagmus, and cupulolithiasis with immediate-onset persistent nystagmus.

9. Which cerebellar regions are involved in modulating the vestibulo-ocular reflex and facilitating vestibular compensation?

A. Cerebellar cortex

B. Cerebral cortex

C. Basal ganglia

D. Flocculus and nodulus

The question tests understanding that the vestibulocerebellum, specifically the flocculus and nodulus, is the brain region responsible for fine-tuning gaze stability and adapting to vestibular changes. The flocculus and nodulus receive vestibular and eye-movement signals and use Purkinje cell outputs to modulate the vestibular nuclei, adjusting the vestibulo-ocular reflex gain and timing so eye movements compensate accurately for head movements. This same circuitry also supports vestibular compensation after unilateral vestibular loss by recalibrating how vestibular input translates into eye movements, helping to restore stable gaze. Other brain areas listed aren't primary players in this automatic reflex modulation or its compensation. The cerebral cortex and basal ganglia contribute to higher-level planning and voluntary movement control, not the immediate reflex refinement and adaptation of the VOR. The broader cerebellar cortex isn't specific enough; it's the vestibulocerebellum—the flocculus and nodulus—that handle these vestibular-specific adjustments.

SAMPLE

10. Which examination maneuver is commonly used to diagnose BPPV?

A. Romberg test

B. Head impulse test

C. Dix-Hallpike maneuver

D. Caloric test

The test is designed to provoke the specific vertigo caused by displaced otoconia in a semicircular canal, most commonly the posterior canal. By moving the head into a position that aligns gravity with that canal, the endolymph moves and triggers the abnormal eye movements and dizziness characteristic of BPPV. A positive result shows a brief onset of vertigo with a distinctive nystagmus pattern—usually upgoing with a torsional component toward the affected ear—often with a latency and fatigability on repeated testing. This maneuver directly tests the pathway most commonly involved in BPPV, making it the best choice for diagnosis. Other tests don't target the same mechanism. The Romberg test assesses static balance and proprioception rather than canalith-induced vertigo. The head impulse test evaluates high-frequency vestibulo-ocular reflex gain to detect acute unilateral vestibular weakness, not positional vertigo. The caloric test stretches the vestibular response to low-frequency stimulation and is used for broader vestibular function assessment, not specifically to diagnose BPPV.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://vestibularsystem.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE