VISCERAL CLET IRAT 3 PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://visceralcletirat3.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. How many functionally independent segments are defined by the Couinaud liver segmentation system?

A. 4

B. 6

C. 10

D. 8

2. Which sexually transmitted disease is implicated in anogenital cancer?

A. Syphilis

B. Chlamydia

C. Human Papilloma Virus

D. Gonorrhea

3. Which finding is most strongly associated with the described case of suspected multiple myeloma?

A. Increased serum albumin

B. Increased serum acid phosphatase

C. Decreased serum albumin

D. Bence Jones proteinuria

4. In pre-hepatic jaundice, which bilirubin form is elevated?

A. Direct bilirubin

B. Indirect bilirubin

C. Both direct and indirect

D. None of the above

5. Which artery runs along the top of the pancreas toward the spleen?

A. Gastroduodenal artery

B. Common hepatic artery

C. Splenic artery

D. Superior mesenteric artery

SAMPLE

6. Which organism is responsible for syphilis?

A. Staphylococcus aureus

B. Treponema pallidum

C. Streptococcus pyogenes

D. Mycobacterium tuberculosis

7. Outline the hepatic lymph drainage pathway.

A. To the stomach

B. Lymphatics accompany the portal triad to hepatic lymph nodes, then to celiac nodes, and finally to the thoracic duct via cisterna chyli.

C. Directly to the inferior vena cava

D. To the spleen

8. Which laboratory finding is characteristic of multiple myeloma?

A. Increased serum albumin

B. Bence Jones proteinuria

C. Decreased serum globulin

D. Normal sedimentation rate

9. Which vessels run along the pancreas's superior border?

A. Hepatic artery

B. Splenic vessels

C. Superior mesenteric artery

D. Gastroduodenal artery

10. Which migraine type is most common in the general population?

A. Basilar

B. Common

C. Classic

D. Status migrainosus

SAMPLE

Answers

SAMPLE

1. D
2. C
3. D
4. B
5. C
6. B
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. How many functionally independent segments are defined by the Couinaud liver segmentation system?

A. 4

B. 6

C. 10

D. 8

The idea being tested is how the liver is divided into functionally independent units. In the Couinaud system, eight segments are defined, each receiving its own portal inflow and biliary drainage, and each draining venously into the hepatic veins. This functional independence comes from how the portal triads branch and how the hepatic veins carve out boundaries, producing eight discrete segments numbered I through VIII (with I being the caudate lobe and the others arranged around it). Because each segment operates with its own vascular and biliary supply, surgeons can target specific segments for resection without disturbing the others. So, eight is the correct number.

2. Which sexually transmitted disease is implicated in anogenital cancer?

A. Syphilis

B. Chlamydia

C. Human Papilloma Virus

D. Gonorrhea

HPV is the sexually transmitted infection most strongly linked to anogenital cancer. High risk HPV types, especially 16 and 18, can persist in the cells lining the genital and anal areas. Over time, the viral proteins E6 and E7 disrupt the cell's tumor-suppressor pathways (p53 and Rb), which normally keep cell growth in check. When these pathways are inactivated, cells can accumulate genetic changes and progress from dysplasia to precancerous lesions and eventually to cancer. This is why HPV infection is a major driver of not only cervical cancer but other anogenital cancers as well as some oropharyngeal cancers. Vaccines targeting these high risk types reduce the risk by preventing infection. The other infections listed are serious STDs with different health impacts, but they do not have the same well-established causal link to cancer as HPV.

3. Which finding is most strongly associated with the described case of suspected multiple myeloma?

A. Increased serum albumin

B. Increased serum acid phosphatase

C. Decreased serum albumin

D. Bence Jones proteinuria

In multiple myeloma, malignant plasma cells overproduce a monoclonal immunoglobulin and, more characteristically, its free light chains. These light chains are small enough to pass through the glomerulus and are not fully reabsorbed by the proximal tubules, so they appear in the urine as Bence Jones proteins. Their presence is a classic and strong clue toward this diagnosis and they can also contribute to kidney damage (myeloma kidney) through tubular precipitation and cast formation. Other options don't point as clearly to this disease: serum albumin levels can fluctuate with many conditions and aren't specific to myeloma, and serum acid phosphatase isn't a hallmark feature of multiple myeloma.

SAMPLE

4. In pre-hepatic jaundice, which bilirubin form is elevated?

A. Direct bilirubin

B. Indirect bilirubin

C. Both direct and indirect

D. None of the above

Pre-hepatic jaundice is driven by increased production of bilirubin, usually from hemolysis or ineffective erythropoiesis. The liver's capacity to conjugate bilirubin isn't the limiting factor here; there's simply more bilirubin arriving than can be processed. This leads to accumulation of unconjugated (indirect) bilirubin in the blood, since it is not yet water-soluble and cannot be excreted in urine. Direct (conjugated) bilirubin stays normal or only mildly elevated because the conjugation step in the liver is intact. So, the hallmark is a rise in indirect bilirubin. If the issue were post-hepatic (obstructive), direct bilirubin would rise due to impaired excretion into the gut; in hepatocellular jaundice, both forms can be elevated because conjugation is impaired.

5. Which artery runs along the top of the pancreas toward the spleen?

A. Gastroduodenal artery

B. Common hepatic artery

C. Splenic artery

D. Superior mesenteric artery

The arteries supplying the pancreas and spleen include the splenic artery, which travels along the superior border of the pancreas as it moves toward the spleen. This long, tortuous vessel branches from the celiac trunk and runs in the splenorenal ligament, giving pancreatic branches along its course and finally reaching the spleen at its hilum. That path—hugging the top edge of the pancreas and heading toward the spleen—is distinctive for the splenic artery. In contrast, the gastroduodenal artery runs along the stomach/duodenum side near the head of the pancreas, the common hepatic artery stays higher up to feed the liver and stomach regions, and the superior mesenteric artery travels inferiorly to supply the midgut, not along the pancreas's superior border toward the spleen.

6. Which organism is responsible for syphilis?

A. Staphylococcus aureus

B. Treponema pallidum

C. Streptococcus pyogenes

D. Mycobacterium tuberculosis

Syphilis is caused by Treponema pallidum, a slender corkscrew-shaped spirochete that lives in human tissues. This organism is transmitted mainly through sexual contact (and can be passed from mother to fetus). It's notable because it's difficult to grow in ordinary laboratory media, so diagnosis often relies on specialized serologic tests or direct visualization with dark-field microscopy. The other organisms listed are responsible for different diseases—Staphylococcus aureus for many skin and systemic infections, Streptococcus pyogenes for strep throat and related illnesses, and Mycobacterium tuberculosis for tuberculosis. Treponema pallidum uniquely fits syphilis.

SAMPLE

7. Outline the hepatic lymph drainage pathway.

A. To the stomach

B. Lymphatics accompany the portal triad to hepatic lymph nodes, then to celiac nodes, and finally to the thoracic duct via cisterna chyli.

C. Directly to the inferior vena cava

D. To the spleen

The pathway tested is how liver lymph exits and travels to the venous system. Lymphatic vessels in the liver run with the portal triad in the hepatoduodenal ligament toward the porta hepatis, collecting interstitial lymph from the liver. From there, lymph moves to hepatic lymph nodes at the porta hepatis, then onward to the celiac (coeliac) nodes. From the celiac nodes, lymph drains into the cisterna chyli and then up the thoracic duct, which delivers lymph into the left venous angle near the junction of the left subclavian and internal jugular veins. This sequence—hepatic lymphatics with the portal triad to hepatic lymph nodes, to celiac nodes, and finally to the thoracic duct via the cisterna chyli—fits the standard hepatic drainage route. Direct drainage to the inferior vena cava is not the lymphatic path, and drainage to the spleen or stomach is not the primary hepatic drainage route.

8. Which laboratory finding is characteristic of multiple myeloma?

A. Increased serum albumin

B. Bence Jones proteinuria

C. Decreased serum globulin

D. Normal sedimentation rate

In multiple myeloma, malignant plasma cells produce a monoclonal immunoglobulin along with excess free light chains that circulate in the blood and are filtered into the urine. These free light chains are what show up as Bence Jones proteinuria. They are small enough to pass through the kidney filter and are detected with urine protein electrophoresis or immunofixation, helping to confirm the presence of a monoclonal gammopathy typical of this disease. This finding stands out because serum albumin is not specifically increased in this condition—if anything, albumin may be normal or reduced as disease burden grows. Serum globulin, reflecting immunoglobulins, is typically increased due to the monoclonal protein rather than decreased. The erythrocyte sedimentation rate is usually elevated because of the high levels of serum proteins and rouleaux formation, not normal.

9. Which vessels run along the pancreas's superior border?

A. Hepatic artery

B. Splenic vessels

C. Superior mesenteric artery

D. Gastroduodenal artery

Vessels that run along the pancreas's superior border are the splenic vessels. The splenic artery travels along the upper edge of the pancreas within the splenorenal ligament, giving branches to the pancreas as it heads toward the spleen. The splenic vein lies close to the pancreas as well, ultimately joining with the superior mesenteric vein to form the portal vein. This pattern distinguishes them from the other vessels listed, which do not trace the pancreas's superior border in the same way. The hepatic artery runs in the hepatoduodenal ligament toward the liver, the superior mesenteric artery travels inferiorly to supply midgut structures, and the gastroduodenal artery courses near the head of the pancreas but not along its entire superior border.

SAMPLE

10. Which migraine type is most common in the general population?

A. Basilar

B. Common

C. Classic

D. Status migrainosus

Most people who migraine without aura, historically called the "common" migraine, experience no sensory or visual disturbances before the headache. Aura is present only in a minority of sufferers, and when it is, that pattern is the classic migraine. Basilar migraine is a rare form with brainstem-type aura symptoms, and status migrainus is a prolonged, severe attack that's also uncommon. Because migraine without aura makes up the vast majority of cases, it is the most common type in the general population.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://visceralcletirat3.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE