WEST COAST EMT PROGRAM PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://westcoastemtprog.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1
Table of Contents 2
Introduction 3
How to Use This Guide 4
Questions 5
Answers 8
Explanations 10
Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **In which of the following situations should the EMT splint an injured limb in the position of deformity?**

   A. If resistance is encountered or the patient experiences severe pain

   B. If transport time to the hospital is greater than 20 to 30 minutes

   C. When distal circulation and neurological functions are absent

   D. If a traction splint will be used to immobilize the injured extremity

2. **In a patient with rapid and shallow respirations, what is the EMT's priority intervention?**

   A. Administer oral glucose

   B. Provide supplemental oxygen

   C. Obtain a full set of vitals

   D. Start CPR immediately

3. **When is gastric distention most likely to occur during ventilation?**

   A. When the airway is completely obstructed.

   B. If the patient is ventilated too quickly.

   C. In patients who are intubated.

   D. With minimal tidal volume delivery.

4. **If you arrive at a critically ill patient and the closest hospital is 25 miles away, what should you do?**

   A. Load him into the ambulance and perform all treatment en route.

   B. Perform a detailed secondary assessment, check vital signs, and transport rapidly.

   C. Manage threats to airway, breathing, circulation, and consider an ALS request.

   D. Administer oxygen and gather as much medical history as possible.

5. **What action should be taken when inadvertently administering glucose to a stable diabetic patient?**

   A. Notify law enforcement for an incident report.

   B. Document the error and report it to your supervisor.

   C. Exclude the intervention from PCR as it did not cause harm.

   D. Contact medical control and notify them of the error.

SAMPLE

6. What is a common risk factor for an elderly person after a fall?

A. Development of heart disease

B. Increased likelihood of a pulmonary embolism

C. Increased risk of hip fractures

D. Potential for neurological deterioration

7. During a transport of a patient with blunt abdominal trauma, if the patient's level of consciousness decreases, you should:

A. Reassess his vital signs and notify the receiving hospital.

B. Suction his oropharynx and increase the oxygen flow rate.

C. Perform a comprehensive secondary assessment.

D. Insert an airway adjunct and begin BVM ventilation.

8. What type of tissue connects bones to other bones?

A. Tendons

B. Cartilage

C. Ligaments

D. Muscle

9. What action should be taken if a patient shows signs of shock after an abdominal trauma?

A. Give the patient fluids before transport.

B. Perform CPR immediately.

C. Monitor vital signs and maintain a high level of alertness.

D. Reassess the patient's condition every 5 minutes.

10. For transporting an unresponsive female patient from upstairs, what is the safest method?

A. Secure her to a stair chair with the strongest provider at the foot end.

B. Secure her to a fracture board with the strongest provider at the foot end.

C. Secure her to a fracture board with the strongest provider at the head end.

D. Secure her to a stair chair with the strongest provider at the head end.

SAMPLE

Answers

SAMPLE

1. A
2. B
3. B
4. C
5. D
6. C
7. D
8. C
9. C
10. C

SAMPLE

Explanations

SAMPLE

1. In which of the following situations should the EMT splint an injured limb in the position of deformity?

A. If resistance is encountered or the patient experiences severe pain

B. If transport time to the hospital is greater than 20 to 30 minutes

C. When distal circulation and neurological functions are absent

D. If a traction splint will be used to immobilize the injured extremity

The appropriate situation to splint an injured limb in the position of deformity is when resistance is encountered or the patient experiences severe pain. This practice is important because attempting to realign a limb that is severely deformed can cause further injury, exacerbate pain, or lead to complications such as damage to nerves and blood vessels. Preserving the limb in its current position, particularly if it is the most comfortable for the patient, helps to minimize movement and trauma to the injury during transport. In circumstances where a person shows obvious signs of distress or pain when trying to change the position of the limb, it indicates that the injury is likely complex and could be compounded by further handling. Therefore, it's crucial to avoid altering the position until the patient receives professional evaluation and care. The other contexts, such as longer transport times or absent circulation and neurological function, might lead to different protocols, like prioritizing circulation or different splinting techniques. However, the priority remains to avoid causing additional harm and discomfort to the patient whenever they exhibit significant pain or resistance to movement.

2. In a patient with rapid and shallow respirations, what is the EMT's priority intervention?

A. Administer oral glucose

B. Provide supplemental oxygen

C. Obtain a full set of vitals

D. Start CPR immediately

In a scenario where a patient exhibits rapid and shallow respirations, the priority intervention for the EMT is to provide supplemental oxygen. This is crucial because rapid and shallow respirations indicate that the patient may not be effectively ventilating, which can lead to inadequate oxygen exchange and subsequent hypoxia. By administering supplemental oxygen, the EMT helps to ensure that the patient receives enough oxygen to maintain adequate oxygen saturation and support vital organ function. Supplemental oxygen can help address the immediate need for oxygenation, which is often more critical in cases of poor respiratory effort. While obtaining vitals is important for overall assessment, it should follow the provision of oxygen since the patient's breathing pattern directly impacts their oxygen levels and overall stability. Administering oral glucose is specific to hypoglycemic emergencies and would not address the immediate risks posed by inadequate ventilation. Starting CPR is necessary only if the patient exhibits signs of cardiac arrest, so it is not the immediate intervention unless the patient becomes unresponsive and pulseless.

SAMPLE

3. When is gastric distention most likely to occur during ventilation?

A. When the airway is completely obstructed.

B. If the patient is ventilated too quickly.

C. In patients who are intubated.

D. With minimal tidal volume delivery.

Gastric distention is primarily a result of air entering the stomach during ventilation, which is more likely to occur when a patient is ventilated too quickly. When ventilation is administered rapidly, there is a tendency for the air to be pushed into the stomach rather than into the lungs, especially if the airway is not properly positioned or if there are other complications such as an obstructed airway. Ventilating a patient too quickly can lead to an increased pressure that forces air into the esophagus instead of the trachea, contributing to gastric distention. This situation can be exacerbated by inadequate relaxation of the diaphragm or ineffective airway positioning. In contrast, an obstructed airway, while serious, does not directly correlate with gastric distention; rather, it typically prevents proper ventilation altogether. Intubated patients, while requiring careful monitoring, are generally less likely to experience gastric distention if proper techniques are followed. Finally, minimal tidal volume delivery refers to the volume of air being moved and does not inherently lead to gastric distention when ventilation is performed correctly.

4. If you arrive at a critically ill patient and the closest hospital is 25 miles away, what should you do?

A. Load him into the ambulance and perform all treatment en route.

B. Perform a detailed secondary assessment, check vital signs, and transport rapidly.

C. Manage threats to airway, breathing, circulation, and consider an ALS request.

D. Administer oxygen and gather as much medical history as possible.

Choosing to manage threats to airway, breathing, and circulation while considering an Advanced Life Support (ALS) request is the correct approach in this scenario. This decision is vital because it prioritizes stabilizing the patient's immediate life-threatening conditions, which is critical for critically ill patients. Airway, breathing, and circulation are the foundational aspects of patient care in emergency situations, and addressing any issues in these areas should always be the first step. In a situation where the hospital is 25 miles away, time is of the essence. By ensuring that airway, breathing, and circulation are secure, you can minimize the risk of further deterioration during transport. Additionally, considering an ALS request is essential if the patient's condition warrants advanced interventions that may be beyond the scope of Basic Life Support (BLS) staff. This ensures that more comprehensive care can be provided either during transport or upon arrival at the hospital. The other options involve performing treatments or assessments that could delay the transport of the patient. While gathering medical history or performing a detailed secondary assessment can be important, in the case of a critically ill patient, immediate stabilization takes precedence over thorough assessment or treatment strategies that could occur en route. Administering oxygen is beneficial but should also be done within the context of ensuring the

SAMPLE

5. What action should be taken when inadvertently administering glucose to a stable diabetic patient?

A. Notify law enforcement for an incident report.

B. Document the error and report it to your supervisor.

C. Exclude the intervention from PCR as it did not cause harm.

D. Contact medical control and notify them of the error.

When administering glucose to a stable diabetic patient, careful communication is critical, especially in the event of an error. Contacting medical control and notifying them of the error serves multiple important functions. First, it ensures that a medical professional is aware of the situation and can provide guidance on how to proceed. This can include instructions for monitoring the patient for any unforeseen effects from the inadvertent glucose administration, as well as advice on subsequent care. Additionally, reporting to medical control helps maintain a clear record of the incident. This transparency is essential for any future treatment decisions and for the overall quality assurance of the EMS operation. It is a standard practice to keep lines of communication open to ensure the safety of the patient and to uphold the integrity of the EMS system. This approach encourages accountability and can facilitate necessary follow-ups. The other options, while they may seem reasonable, do not prioritize patient safety and rigorous clinical practice in the same way. Failing to report the error can prevent important patient safety measures and undermine trust in the EMS process.

6. What is a common risk factor for an elderly person after a fall?

A. Development of heart disease

B. Increased likelihood of a pulmonary embolism

C. Increased risk of hip fractures

D. Potential for neurological deterioration

A common risk factor for an elderly person after a fall is the increased risk of hip fractures. This demographic often has decreased bone density due to conditions like osteoporosis, making bones more fragile and susceptible to fractures during falls. When an elderly person falls, the impact can easily surpass the strength of their weakened bones, particularly in the hip area, which is notably vulnerable due to its structure and weight-bearing function. Additionally, fractures can lead to significant complications, including increased morbidity, prolonged hospital stays, and decreased mobility, which can further exacerbate health issues for the elderly, prolonging recovery and affecting overall quality of life. Understanding this risk is essential in fall prevention strategies and managing post-fall care for older adults.

SAMPLE

7. During a transport of a patient with blunt abdominal trauma, if the patient's level of consciousness decreases, you should:

A. Reassess his vital signs and notify the receiving hospital.

B. Suction his oropharynx and increase the oxygen flow rate.

C. Perform a comprehensive secondary assessment.

D. Insert an airway adjunct and begin BVM ventilation.

In the context of a patient with blunt abdominal trauma experiencing a decrease in their level of consciousness, prioritizing airway management is crucial. When a patient's consciousness is lowered, there is a significant risk of airway compromise due to decreased muscle tone and potential inability to protect their airway. By inserting an airway adjunct, you are ensuring that the airways remain open, which is vital for effective ventilation. Using a Bag-Valve-Mask (BVM) ventilation is also appropriate because it allows for assisted breathing to maintain adequate oxygenation and ventilation when the patient's respiratory drive is compromised. This intervention addresses the immediate risk to the patient's airway and breathing, which is prioritized in any emergency situation. While reassessing vital signs is important and notifying the receiving hospital is crucial for continuity of care, those actions do not directly address the urgent risk posed by the decreased level of consciousness to the patient's airway. Similarly, suctioning and increasing oxygen flow can be part of the management plan but do not directly secure the airway. Comprehensive secondary assessments can be performed once the airway is stabilized, making airway management the most critical and immediate response in this scenario.

8. What type of tissue connects bones to other bones?

A. Tendons

B. Cartilage

C. Ligaments

D. Muscle

The correct choice is ligaments, which are specialized connective tissues that connect bones to other bones at a joint. They provide stability to the joints while allowing for some movement, ensuring that bones remain properly aligned and that the joints function effectively during activities. Ligaments are made up of dense bundles of collagen fibers, which give them strength and elasticity. Tendons, on the other hand, connect muscles to bones, playing a crucial role in movement rather than connecting bones to other bones. Cartilage serves as a cushioning material at joint surfaces and can also form structures like the nose and ears, but it does not connect bones directly to one another. Muscle tissue is involved in the contraction and movement of the body, and does not have a role in connecting bones. Thus, ligaments are specifically the type of tissue that fulfill the requirement of connecting bones to each other.

SAMPLE

9. What action should be taken if a patient shows signs of shock after an abdominal trauma?

A. Give the patient fluids before transport.

B. Perform CPR immediately.

C. Monitor vital signs and maintain a high level of alertness.

D. Reassess the patient's condition every 5 minutes.

When a patient exhibits signs of shock after experiencing abdominal trauma, monitoring vital signs and maintaining a high level of alertness is crucial. This approach is vital because shock can lead to rapid deterioration of the patient's condition. By continuously monitoring vital signs, such as heart rate, blood pressure, and respiratory rate, the EMT can identify any significant changes or trends that may indicate worsening shock or other complications. Additionally, staying alert allows the EMT to quickly recognize any new symptoms or changes in the patient's status, which is essential for effective management and timely intervention as the situation evolves. Recognizing that shock can result from various causes, such as internal bleeding or organ damage after trauma, ensures that the patient receives the appropriate care and is adequately prepared for transport to a medical facility where surgical intervention may be necessary. While administering fluids is a common step in treating shock, it typically occurs in conjunction with transport, rather than before, particularly in cases where immediate transport to a hospital is warranted. CPR is specifically indicated for cardiac arrest situations, and unnecessary reassessment without responsive intervention may lead to delays in critical care. Therefore, maintaining vigilance through monitoring is the best immediate action following signs of shock in this scenario.

10. For transporting an unresponsive female patient from upstairs, what is the safest method?

A. Secure her to a stair chair with the strongest provider at the foot end.

B. Secure her to a fracture board with the strongest provider at the foot end.

C. Secure her to a fracture board with the strongest provider at the head end.

D. Secure her to a stair chair with the strongest provider at the head end.

The correct approach to transporting an unresponsive female patient from upstairs is to secure her to a fracture board with the strongest provider at the head end. This method is optimal for several critical reasons. Firstly, using a fracture board allows for the stabilization of the patient's spine and body, which is vital when dealing with an unresponsive individual. It provides a rigid surface that can help prevent movement, which is essential in case of potential spinal injuries. Positioning the strongest provider at the head end is strategic for both safety and control. This position ensures that the strongest provider can better manage the patient's head and neck during the descent. Since the head is the heaviest part of the body, stabilizing it effectively is crucial to prevent further injury, especially if there is any concern about spinal issues. It also enables better communication with the rest of the team during the transport, as the head-end provider can guide the movement and ensure that the descent is controlled and coordinated. Furthermore, this method helps maintain the patient's airway, which is particularly important for any unresponsive patient. By having the strongest member at the head, there is greater assurance that the patient's airway can be monitored and managed effectively throughout the transport. In contrast, using a stair chair or placing the strongest provider

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://westcoastemtprog.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE