WESTERN GOVERNORS UNIVERSITY (WGU) NURS6800 D116 ADVANCED PHARMACOLOGY FOR THE ADVANCED PRACTICE NURSE PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://wgu-nurs6800d116.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **A psychiatric nurse explains that therapeutic effects may not occur for several weeks with an antidepressant. The nurse understands this is likely due to what?**

   A. Changes in the brain as a result of prolonged drug exposure

   B. The blood-brain barrier blocking effects

   C. Tolerance development

   D. Rapid onset of adverse effects

2. **What does directly observed therapy (DOT) aim to ensure in tuberculosis treatment?**

   A. To shorten treatment duration.

   B. To monitor liver function.

   C. To ensure adherence to the medication regimen.

   D. To avoid need for culture results.

3. **Why are ACE inhibitors contraindicated in bilateral renal artery stenosis and what monitoring is essential?**

   A. ACE inhibitors can cause severe renal failure in bilateral renal artery stenosis; monitor creatinine and potassium.

   B. They improve renal function in bilateral RAS; monitor blood pressure.

   C. They are safe in all forms of kidney disease.

   D. They cause hyperkalemia but good kidney perfusion.

4. **A patient with Alzheimer's disease is taking a cholinesterase inhibitor. Which side effects would most concern the nurse due to potential cardiovascular risk?**

   A. Bradycardia and fainting

   B. Dizziness, headache, nausea

   C. Cough and hoarseness

   D. Weight gain and edema

5. **What is the shared primary mechanism of action for ACE inhibitors and ARBs in blood pressure management?**

   A. Inhibit HMG-CoA reductase

   B. Block the RAAS

   C. Block the beta-adrenergic receptors

   D. Activate PPAR gamma

SAMPLE

6. A patient taking bismuth subsalicylate [Pepto-Bismol] has a black tongue on examination. What should the NP do?

A. Reassure the patient that this is an expected side effect.

B. Consider this a sign of GI bleeding.

C. Suspect hepatotoxicity.

D. Assess for drug overdose.

7. Which vaccine should not be administered to an immunocompromised child?

A. Varicella vaccine

B. DTaP vaccine

C. Hib vaccine

D. Polio vaccine

8. When planning care for an advanced heart failure patient, with whom should end-of-life discussions most appropriately occur?

A. With the patient and family, guided by patient preferences

B. Only with the cardiology team in hospital

C. With the patient's attorney only

D. Not discussed unless patient requests

9. Regarding laxatives, which statement best reflects their general use?

A. Laxatives should generally not be used long-term.

B. Laxatives are the best option for chronic constipation over years.

C. Laxatives have no adverse effects.

D. Laxatives cure all forms of diarrhea.

10. Which prognosis best describes Stage D heart failure?

A. Poor prognosis with limited response to conventional therapies

B. Excellent prognosis with diuretics

C. No impact from device therapy

D. Always fully reversible with medication

SAMPLE

Answers

SAMPLE

1. A
2. C
3. A
4. A
5. B
6. A
7. A
8. A
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. A psychiatric nurse explains that therapeutic effects may not occur for several weeks with an antidepressant. The nurse understands this is likely due to what?

A. Changes in the brain as a result of prolonged drug exposure

B. The blood-brain barrier blocking effects

C. Tolerance development

D. Rapid onset of adverse effects

Therapeutic effects from antidepressants take weeks because the brain must undergo adaptive changes after the drug is started. While the medication quickly increases monoamine levels in the synapse, the mood improvements come from slower downstream processes—altered receptor sensitivity, changes in gene expression, and increased neurotrophic signaling like brain-derived neurotrophic factor that promote neuroplasticity and hippocampal neurogenesis. These neuroadaptive changes unfold over time, explaining the delay before clinical benefit appears. The other ideas don't fit this pattern: the blood-brain barrier doesn't account for a delayed response, tolerance would more likely reduce effect rather than delay it, and rapid onset of adverse effects doesn't explain a gradual improvement in mood.

2. What does directly observed therapy (DOT) aim to ensure in tuberculosis treatment?

A. To shorten treatment duration.

B. To monitor liver function.

C. To ensure adherence to the medication regimen.

D. To avoid need for culture results.

Directly observed therapy is about ensuring that patients take their TB medications as prescribed. TB treatment requires daily, multi-drug therapy for an extended period, and lapses in taking pills can lead to relapse and the development of drug resistance. Having a healthcare worker or trained observer watch the patient swallow each dose helps catch missed doses, provides support, and keeps the patient on the full regimen, which improves adherence and cure rates and reduces transmission. This approach isn't meant to shorten the treatment duration or replace safety monitoring (liver function checks are still done when indicated). It also doesn't affect the need for diagnostic cultures; DOT's purpose is strictly to ensure adherence to the prescribed regimen.

SAMPLE

3. Why are ACE inhibitors contraindicated in bilateral renal artery stenosis and what monitoring is essential?

A. ACE inhibitors can cause severe renal failure in bilateral renal artery stenosis; monitor creatinine and potassium.

B. They improve renal function in bilateral RAS; monitor blood pressure.

C. They are safe in all forms of kidney disease.

D. They cause hyperkalemia but good kidney perfusion.

The key idea is that ACE inhibitors can dangerously reduce kidney function when both renal arteries are narrowed because the kidneys rely on angiotensin II to keep glomerular filtration pressure high. Blocking angiotensin II removes this efferent arteriolar constriction, lowering glomerular capillary pressure and GFR. In bilateral renal artery stenosis there isn't a healthy downstream kidney to compensate, so this can precipitate acute kidney injury. The risk is compounded by the potential for hyperkalemia due to reduced aldosterone-mediated potassium excretion. Because of this, the essential monitoring after starting or increasing an ACE inhibitor is measuring serum creatinine to detect a decline in kidney function and checking potassium levels to catch hyperkalemia. Baseline values should be obtained, with follow-up labs in the first one to two weeks (and sooner if symptoms appear), then periodically thereafter. Blood pressure should also be monitored, but the critical safety signals are the creatinine and potassium changes.

4. A patient with Alzheimer's disease is taking a cholinesterase inhibitor. Which side effects would most concern the nurse due to potential cardiovascular risk?

A. Bradycardia and fainting

B. Dizziness, headache, nausea

C. Cough and hoarseness

D. Weight gain and edema

Cholinesterase inhibitors boost acetylcholine in both the central and peripheral nervous systems, which increases parasympathetic (vagal) effects on the heart. Activation of cardiac muscarinic receptors slows the SA node and slows AV nodal conduction, so the heart rate can drop. In older adults, especially those with underlying conduction abnormalities or who are taking other medications that slow heart rate, this can lead to significant bradycardia and even fainting from reduced cerebral perfusion. That cardiovascular risk—bradycardia with possible syncope—is why the combination of a slow pulse and fainting is the most concerning side effect. Other common adverse effects like dizziness, headache, and nausea can occur with these drugs but do not carry the same immediate cardiovascular risk. Cough and hoarseness aren't typical cardiovascular concerns with this class, and weight gain with edema isn't characteristic. Clinically, monitor heart rate and rhythm, assess for symptomatic bradycardia or syncope, and review concurrent AV-nodal depressants to determine if dose adjustment or discontinuation is needed.

SAMPLE

5. What is the shared primary mechanism of action for ACE inhibitors and ARBs in blood pressure management?

A. Inhibit HMG-CoA reductase

B. Block the RAAS

C. Block the beta-adrenergic receptors

D. Activate PPAR gamma

Both ACE inhibitors and ARBs lower blood pressure by interrupting the renin-angiotensin-aldosterone system. ACE inhibitors block the conversion of angiotensin I to angiotensin II, reducing angiotensin II–mediated vasoconstriction and aldosterone release. ARBs block the angiotensin II type 1 receptor, preventing angiotensin II from producing its usual effects. Either way, there is less vasoconstriction and less aldosterone-driven sodium and water retention, leading to lower blood pressure. The other options involve different drug classes—statins affect cholesterol synthesis, beta-blockers blunt sympathetic stimulation, and PPAR gamma activators are used for insulin sensitivity—so they don't share the same primary mechanism as ACE inhibitors and ARBs.

6. A patient taking bismuth subsalicylate [Pepto-Bismol] has a black tongue on examination. What should the NP do?

A. Reassure the patient that this is an expected side effect.

B. Consider this a sign of GI bleeding.

C. Suspect hepatotoxicity.

D. Assess for drug overdose.

Discoloration of the tongue is a known, harmless effect of bismuth subsalicylate. The bismuth can react with sulfur compounds in saliva and on the tongue to form black bismuth sulfide, giving the tongue a dark appearance. This does not indicate bleeding, liver injury, or overdose, so the appropriate action is to reassure the patient and continue the medication as directed. If new symptoms suggestive of GI bleeding (such as vomiting blood or black, tarry stools) or salicylate toxicity (for example ringing in the ears, persistent confusion, or rapid breathing) appear, those would warrant further evaluation.

7. Which vaccine should not be administered to an immunocompromised child?

A. Varicella vaccine

B. DTaP vaccine

C. Hib vaccine

D. Polio vaccine

Live vaccines should be avoided in immunocompromised individuals because their weakened immune systems may not control replication of the attenuated pathogens, risking infection with the vaccine strain. Varicella vaccine is a live attenuated vaccine, containing a weakened varicella-zoster virus. In an immunocompromised child, this can lead to varicella illness or more severe complications from disseminated infection, so it should not be given. The other vaccines listed are inactivated or toxoid-based and do not contain replicating organisms, so they are generally safe to administer to patients with compromised immunity, depending on their specific immune status and clinician judgment. DTaP and Hib are inactivated formulations, and polio is typically given as the inactivated polio vaccine in modern practice; these do not pose the same risk as a live vaccine in this context.

SAMPLE

8. When planning care for an advanced heart failure patient, with whom should end-of-life discussions most appropriately occur?

A. With the patient and family, guided by patient preferences

B. Only with the cardiology team in hospital

C. With the patient's attorney only

D. Not discussed unless patient requests

End-of-life planning should be patient-centered, beginning with the patient and family and guided by the patient's own preferences. This approach honors autonomy and ensures that care goals reflect what is most important to the patient, whether that means focusing on comfort, pursuing disease-modifying therapies, or a combination of both. In advanced heart failure, where prognosis can be uncertain and trajectories can shift, involving the patient and family in conversations about goals of care, the desired level of intervention, and plans for advance directives or hospice ensures that subsequent treatment choices—including medications, escalation or de-escalation of therapy, and hospital vs. palliative care options—are aligned with the patient's values. Relying solely on the cardiology team, involving only medical-legal professionals, or waiting for the patient to request discussion undermines patient autonomy and can miss expressed preferences. These conversations are most effective when they are ongoing, revisited as the situation evolves, and supported by the broader care team to provide consistent, compassionate guidance that centers the patient's wishes.

9. Regarding laxatives, which statement best reflects their general use?

A. Laxatives should generally not be used long-term.

B. Laxatives are the best option for chronic constipation over years.

C. Laxatives have no adverse effects.

D. Laxatives cure all forms of diarrhea.

Laxatives are intended for short-term relief of constipation rather than a long-term solution. Using them chronically can lead to dependence, reduced natural bowel motility, and electrolyte or fluid imbalances, especially in older adults or those with kidney issues. They can also mask an underlying cause of constipation, delaying diagnosis. The safer, general approach is to address lifestyle factors—adequate fiber and fluids, regular activity—and to reassess the need for laxatives, using the lowest effective dose for the shortest time under supervision. Some situations, such as opioid-induced constipation or certain neuromuscular disorders, may require longer-term management, but that's not the typical course. Laxatives can have adverse effects like cramping, bloating, dehydration, and electrolyte shifts, and they do not cure diarrhea.

10. Which prognosis best describes Stage D heart failure?

A. Poor prognosis with limited response to conventional therapies

B. Excellent prognosis with diuretics

C. No impact from device therapy

D. Always fully reversible with medication

Stage D describes advanced heart failure with persistent symptoms despite optimized medical therapy. In this stage the disease has progressed to a point where conventional treatments provide limited benefit, and mortality and morbidity remain high. Diuretics help relieve congestion but don't improve long-term survival, so an excellent prognosis with diuretics isn't accurate. Device therapies can help some patients by reducing arrhythmias or improving heart function, but they do not reverse the overall trajectory for most with Stage D, so saying there's no impact from device therapy isn't correct. Reversibility with medication alone is not typical at this stage, as the heart failure tends to be progressive and often requires consideration of advanced therapies or palliative approaches. All of this makes the description of a poor prognosis with limited response to conventional therapies the best fit for Stage D.

EXAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://wgu-nurs6800d116.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE