WOMEN HEALTH NURSE PRACTITIONER (WHNP) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://womenhealthnursepractitioner.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which of the following therapies is NOT recommended for long-term management of abnormal uterine bleeding?

A. Copper intrauterine device

B. Levonorgestrel intrauterine system

C. Tranexamic acid

D. Hormonal contraceptives

2. What characterizes cystocele among pelvic organ prolapses?

A. Descent of the bladder into the vagina

B. Descent of the rectum into the vagina

C. Descent of the uterus into the vaginal canal

D. Descent of the post-vaginal wall

3. In which condition is 17α-hydroxyprogesterone caproate injection specifically indicated?

A. A woman with a history of a prior singleton delivery before 37 weeks

B. A woman admitted for preterm labor

C. A woman with a preterm premature rupture of membranes

D. A woman experiencing normal pregnancies

4. What is the recommended screening for cervical cancer in women aged 21-29?

A. Pap smear alone every 3 years

B. HPV testing every year

C. Co-testing (Pap and HPV) every 5 years

D. No screening recommended

5. What type of ovarian cyst occurs when an ovarian follicle is stimulated but fails to reach maturity?

A. Follicular cyst

B. Corpeus luteum cyst

C. Benign ovarian cyst

D. Cystic teratoma

SAMPLE

6. What herbal supplement is commonly used for menopausal symptoms?

A. Ginseng

B. Black cohosh

C. St. John's Wort

D. Valerian root

7. When counseling a woman on contraceptive options, what should the nurse practitioner emphasize as most important?

A. Cost of the methods

B. Effectiveness and individual lifestyle

C. Availability of the methods

D. Popularity among peers

8. Which condition significantly increases the risk of umbilical cord prolapse during labor?

A. Dilation 3 cm

B. Polyhydramnios

C. Single gestation

D. Vertex presentation

9. When treating acute bacterial rhinosinusitis in a pregnant patient, which medication should be avoided?

A. Amoxicillin

B. Azithromycin

C. Levofloxacin

D. Doxycycline

10. What is the primary focus of a Women Health Nurse Practitioner (WHNP)?

A. Providing care to children

B. Providing care to women throughout their lifespan

C. Providing emergency medical services

D. Providing care solely during pregnancy

SAMPLE

Answers

SAMPLE

1. A
2. A
3. A
4. A
5. C
6. B
7. B
8. B
9. C
10. B

SAMPLE

Explanations

SAMPLE

1. Which of the following therapies is NOT recommended for long-term management of abnormal uterine bleeding?

A. Copper intrauterine device

B. Levonorgestrel intrauterine system

C. Tranexamic acid

D. Hormonal contraceptives

The correct answer indicates that the copper intrauterine device (IUD) is not recommended for long-term management of abnormal uterine bleeding. The primary role of the copper IUD is as a form of contraception, and while it can create a local inflammatory response that may lead to heavier menstrual bleeding, it is not typically used to manage abnormal uterine bleeding over the long term. While some patients may experience changes in their menstrual patterns due to the copper IUD, it is known to sometimes exacerbate the very problem it is intended to help manage. In contrast, hormonal contraceptives, the levonorgestrel intrauterine system, and tranexamic acid have established roles in the management of abnormal uterine bleeding. Hormonal contraceptives can effectively regulate menstrual cycles and reduce excessive bleeding. The levonorgestrel intrauterine system releases progestin, which softens the endometrial lining and significantly reduces menstrual blood loss, making it a suitable long-term option. Tranexamic acid is an antifibrinolytic agent that helps to reduce bleeding during menstruation and can be used as needed, but it is not considered a long-term management option like hormonal therapies.

2. What characterizes cystocele among pelvic organ prolapses?

A. Descent of the bladder into the vagina

B. Descent of the rectum into the vagina

C. Descent of the uterus into the vaginal canal

D. Descent of the post-vaginal wall

Cystocele is characterized by the descent of the bladder into the anterior wall of the vagina. This occurs when the supportive tissues between a woman's bladder and vagina weaken, allowing the bladder to bulge into the vaginal canal. This condition can be caused by factors such as childbirth, obesity, and aging, which all contribute to the weakening of the pelvic floor muscles and supportive tissues. In contrast, other conditions like rectocele refer to the descent of the rectum into the vagina, uterine prolapse involves the descent of the uterus into the vaginal canal, and descents in the posterior vaginal wall pertain to the weakening of the structures supporting these organs. Understanding the specific characteristics of each type of prolapse is crucial for diagnosis and treatment, particularly within the scope of women's health and preventive care.

SAMPLE

3. In which condition is 17α-hydroxyprogesterone caproate injection specifically indicated?

A. A woman with a history of a prior singleton delivery before 37 weeks

B. A woman admitted for preterm labor

C. A woman with a preterm premature rupture of membranes

D. A woman experiencing normal pregnancies

The use of 17α-hydroxyprogesterone caproate injection is specifically indicated for women with a history of a prior singleton delivery before 37 weeks. This medication is primarily indicated for the prevention of recurrent preterm birth in women who have had a previous singleton preterm birth. The administration of this injectable medication aims to reduce the risk of subsequent premature deliveries by improving uterine quiescence and enhancing cervical competence through its progestational effects. In clinical practice, it's important to note that this treatment isn't typically initiated during active preterm labor or in cases of preterm premature rupture of membranes, as these conditions require different management strategies. Additionally, the injection is not indicated in women experiencing normal pregnancies, as there is no history or indication that would necessitate intervention to prevent preterm labor. This specificity in its use helps ensure that the treatment is both effective and appropriate for the populations at risk.

4. What is the recommended screening for cervical cancer in women aged 21-29?

A. Pap smear alone every 3 years

B. HPV testing every year

C. Co-testing (Pap and HPV) every 5 years

D. No screening recommended

The recommended screening for cervical cancer in women aged 21-29 is a Pap smear alone every three years. This recommendation is based on guidelines that focus on the efficacy of the Pap smear test, which detects precancerous changes in cervical cells. At this age, the immune system is generally more capable of clearing human papillomavirus (HPV) infections, which are linked to cervical cancer. Therefore, the emphasis is on regular Pap testing rather than HPV testing, which is recommended as a primary screening tool only for women 30 and older. The three-year interval for Pap testing strikes a balance between effectively monitoring cervical health while minimizing unnecessary procedures from potential false-positive results in a younger population, who may experience transient HPV infections. Regular screenings help ensure early detection of abnormalities that may develop into cervical cancer while considering the demographic risks and biological factors pertinent to younger women.

SAMPLE

5. What type of ovarian cyst occurs when an ovarian follicle is stimulated but fails to reach maturity?

A. Follicular cyst

B. Corpeus luteum cyst

C. Benign ovarian cyst

D. Cystic teratoma

The type of ovarian cyst that forms when an ovarian follicle is stimulated but does not reach maturity is a follicular cyst. These cysts develop during the normal menstrual cycle when the follicle, which is responsible for releasing an egg, grows but fails to ovulate. As a result, the fluid that surrounds the immature follicle accumulates, leading to the formation of a cyst. In contrast, a corpus luteum cyst arises after ovulation when the follicle transforms into the corpus luteum, which is responsible for producing hormones necessary for maintaining early pregnancy. Benign ovarian cyst is a more general term that can refer to various non-cancerous cysts, while a cystic teratoma is a type of germ cell tumor that can contain different tissue types but is not directly related to the failure of follicle maturation. Each of these options represents different mechanisms and processes in ovarian function.

6. What herbal supplement is commonly used for menopausal symptoms?

A. Ginseng

B. Black cohosh

C. St. John's Wort

D. Valerian root

Black cohosh is recognized as a commonly used herbal supplement for managing menopausal symptoms, particularly hot flashes, mood swings, and other related discomforts. Its active compounds have been studied for their ability to modulate hormonal activity and help balance estrogen's effects in the body, which is particularly beneficial during the transitional menopausal phase. Ginseng, while sometimes associated with enhancing energy and alleviating stress, does not specifically target menopausal symptoms and thus is not as widely recommended for this purpose. St. John's Wort is primarily used for its antidepressant effects and may not provide relief from the physical symptoms of menopause. Valerian root, known for its sedative properties and often used to aid with sleep, does not specifically address the range of menopausal symptoms as effectively as black cohosh. In summary, black cohosh stands out due to its targeted action in alleviating menopausal symptoms, making it the most recognized and widely used herbal supplement for this condition.

SAMPLE

7. When counseling a woman on contraceptive options, what should the nurse practitioner emphasize as most important?

A. Cost of the methods

B. Effectiveness and individual lifestyle

C. Availability of the methods

D. Popularity among peers

Emphasizing effectiveness and individual lifestyle when counseling a woman on contraceptive options is crucial because these factors play a significant role in the success of any contraceptive method. Effectiveness refers to the reliability of the method in preventing pregnancy, which is a primary concern for many women. Understanding the success rates of different contraceptive methods helps women make informed choices that align with their needs and expectations. Individual lifestyle considerations are equally important; different methods may fit better with a woman's daily routine, health status, and personal preferences. For instance, a woman with a busy lifestyle might prefer a long-acting reversible contraceptive like an IUD or implant, while another may favor a method that allows for more control, such as the pill. Counseling that incorporates both effectiveness and lifestyle facilitates a tailored approach, ensuring that the chosen contraceptive method aligns with the woman's health goals and personal circumstances, ultimately leading to better adherence and satisfaction with the contraceptive method selected. By focusing on these components, the nurse practitioner helps empower women to choose a method that is most likely to be effective and feasible within their context, enhancing the likelihood of successful use.

8. Which condition significantly increases the risk of umbilical cord prolapse during labor?

A. Dilation 3 cm

B. Polyhydramnios

C. Single gestation

D. Vertex presentation

Polyhydramnios, or an excess of amniotic fluid, significantly increases the risk of umbilical cord prolapse during labor. The abundance of fluid allows for greater fetal movement, which can lead to a situation where the umbilical cord slips ahead of the presenting part of the fetus. This is particularly critical during labor as the head descends into the birth canal. If the presenting part does not adequately engage, the cord can become trapped between the fetus and the cervix, compromising fetal blood flow and oxygenation. In the case of dilation being at a low level, such as 3 cm, there is typically still a considerable amount of time before delivery, and the fetus is not yet engaged in the birth canal. This reduces the likelihood of cord prolapse at that stage. In situations of single gestation and vertex presentation, the positional stability of the fetus further mitigates the risks associated with umbilical cord prolapse, as these scenarios generally involve a well-engaged fetal head that acts as a barrier to the cord, preventing it from slipping past. Polyhydramnios creates a different dynamic altogether that can lead to an increased risk for this complication.

SAMPLE

9. When treating acute bacterial rhinosinusitis in a pregnant patient, which medication should be avoided?

A. Amoxicillin

B. Azithromycin

C. Levofloxacin

D. Doxycycline

Acute bacterial rhinosinusitis in pregnant patients requires careful consideration when choosing appropriate medications due to potential risks to the fetus. Levofloxacin, which is a fluoroquinolone antibiotic, is generally avoided during pregnancy. This is primarily because fluoroquinolones have been associated with risks such as cartilage damage in developing joints and potential effects on the fetus's musculoskeletal development. In contrast, both amoxicillin and azithromycin are often considered safe for use during pregnancy, as they have established safety profiles. Doxycycline, however, is typically avoided in pregnant patients because it can affect the development of teeth and bones in the fetus due to its action as a tetracycline antibiotic. Understanding the safety profiles of these antibiotics is crucial for effective treatment while minimizing risks during pregnancy.

10. What is the primary focus of a Women Health Nurse Practitioner (WHNP)?

A. Providing care to children

B. Providing care to women throughout their lifespan

C. Providing emergency medical services

D. Providing care solely during pregnancy

The primary focus of a Women Health Nurse Practitioner (WHNP) is to provide care to women throughout their lifespan. This comprehensive approach includes addressing various aspects of women's health, from adolescence through menopause and into older age. WHNPs are trained to manage a wide range of health issues that specifically affect women, including reproductive health, preventive care, chronic disease management, and reproductive life planning. The versatility of the WHNP role enables practitioners to offer holistic care that considers the physical, emotional, and social aspects of health, which is essential for promoting overall wellness in women. This role is distinct from those focused solely on specific populations or conditions, such as children or emergency services. While some WHNPs may provide care related to pregnancy, their scope is much broader, encompassing all health issues relevant to women at different stages of life.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://womenhealthnursepractitioner.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE