WOMEN'S HEALTH END OF ROTATION (EOR) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://womenshealtheor.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the definition of oligomenorrhea?

A. Heavy menstrual bleeding

B. Lack of menstrual periods for several months

C. Irregular or infrequent menstrual periods

D. Severe menstrual cramps

2. In which situation is a pelvic ultrasound particularly indicated?

A. Suspicion of ovarian cysts

B. Evaluation of menstrual irregularities

C. Confirmation of pregnancy

D. Assessment of uterine fibroids

3. In what part of the female reproductive system does fertilization usually occur?

A. Uterus

B. Ovaries

C. Fallopian tubes

D. Cervix

4. What is a major potential complication of untreated chlamydia in women?

A. Uterine fibroids

B. Fibrocystic breast changes

C. Pelvic inflammatory disease (PID)

D. Premenstrual syndrome (PMS)

5. What condition could result in the need for a myomectomy?

A. Cervical cancer

B. Uterine fibroids

C. Endometriosis

D. Pelvic inflammatory disease

6. What screening test is recommended for detecting cervical cancer?

A. Ultrasound

B. CT scan

C. Pap smear

D. MRI

SAMPLE

7. What is typically observed in a pregnancy exam at 28 weeks?

A. Fundus at the pubic symphysis

B. Fundus at the umbilicus

C. Fundus above the umbilicus

D. Fundus below the pubic symphysis

8. Which of the following is a contraindication to prescribing oral hormonal contraceptives?

A. History of iron deficiency anemia

B. History of dysmenorrhea since age 15

C. Positive family history of ovarian cancer

D. History of smoking in a patient older than age 35

9. What is the typical presentation of uterine fibroids?

A. Painful urination

B. Menorrhagia

C. Night sweats

D. Weight gain

10. Which of the following is a major risk factor for an ectopic pregnancy?

A. Alcohol intake

B. Young maternal age

C. History of salpingitis

D. Low dose oral contraceptive use

SAMPLE

Answers

SAMPLE

1. C
2. A
3. C
4. C
5. B
6. C
7. C
8. D
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. What is the definition of oligomenorrhea?

A. Heavy menstrual bleeding

B. Lack of menstrual periods for several months

C. Irregular or infrequent menstrual periods

D. Severe menstrual cramps

Oligomenorrhea refers to irregular or infrequent menstrual periods, typically characterized by a menstrual cycle that occurs more than 35 days apart but less than six months. It signifies a disruption in the normal menstrual rhythm and can be associated with various underlying conditions, including hormonal imbalances, overweight or obesity, excessive exercise, stress, and polycystic ovary syndrome (PCOS). This definition aligns with the chosen answer, highlighting that oligomenorrhea indicates not just the absence of menstruation for a protracted period but rather a pattern of cycles that occurs less frequently than expected. Understanding this concept is crucial in evaluating and managing patients who present with menstrual irregularities, helping clinicians pinpoint potential diagnoses and treatment options. The other definitions provided do not accurately capture the essence of oligomenorrhea; heavy menstrual bleeding refers to menorrhagia, lack of menstrual periods for several months describes amenorrhea, and severe menstrual cramps pertain to dysmenorrhea.

2. In which situation is a pelvic ultrasound particularly indicated?

A. Suspicion of ovarian cysts

B. Evaluation of menstrual irregularities

C. Confirmation of pregnancy

D. Assessment of uterine fibroids

A pelvic ultrasound is particularly indicated in the suspicion of ovarian cysts because it allows for direct visualization of the ovaries and assessment of any cystic structures present. This imaging modality is non-invasive and provides detailed information regarding the size, shape, and characteristics of any cysts, helping to differentiate between functional and pathological cysts. It can also assess for signs of complications, such as rupture or hemorrhage. This makes it an essential tool in managing suspected ovarian cysts, providing necessary information for appropriate clinical decision-making and follow-up. While pelvic ultrasound is also useful for evaluating menstrual irregularities, confirming pregnancy, and assessing uterine fibroids, each of these situations may not be as specifically indicated solely for the ultrasound when compared to the direct diagnostic capability it provides for ovarian cysts. For instance, other tests may precede confirmation of pregnancy, and menstrual irregularities may involve additional assessments beyond imaging. Nonetheless, the distinct advantage of ultrasound in visualizing ovarian structures affirms its particular indication in the context of suspected ovarian cysts.

SAMPLE

3. In what part of the female reproductive system does fertilization usually occur?

A. Uterus

B. Ovaries

C. Fallopian tubes

D. Cervix

Fertilization typically occurs in the fallopian tubes, which serve as a crucial site for the merging of sperm and egg. After ovulation, when an egg is released from the ovary, it enters the fallopian tube where it can be met by sperm. The fallopian tubes provide an optimal environment for this process, as they are equipped with cilia that help transport the fertilized egg towards the uterus for implantation. Understanding the function of the fallopian tubes highlights their role in human reproduction. They not only facilitate the encounter between sperm and egg but also are the pathway through which the fertilized egg travels to reach the uterus, an essential step for implantation and further development into a fetus. In contrast, while the uterus is crucial for gestation, it is not where fertilization takes place. The ovaries are responsible for the production of eggs but are not involved in the fertilization process itself. The cervix serves as a canal for sperm to enter the uterus and does not play a role in the fertilization of the egg. Hence, the fallopian tubes are specifically designed to support fertilization, making this choice the correct answer.

4. What is a major potential complication of untreated chlamydia in women?

A. Uterine fibroids

B. Fibrocystic breast changes

C. Pelvic inflammatory disease (PID)

D. Premenstrual syndrome (PMS)

In women, untreated chlamydia can lead to significant reproductive health complications, with pelvic inflammatory disease (PID) being one of the most serious outcomes. Chlamydia is a sexually transmitted infection that can ascend from the cervix and uterus into the pelvic cavity, impacting the fallopian tubes, ovaries, and peritoneum. PID can result in chronic pelvic pain, ectopic pregnancy, and infertility due to scarring and damage to reproductive organs. This complication is particularly concerning because many women with chlamydia may be asymptomatic or have mild symptoms, which can lead to delays in diagnosis and treatment. Early identification and management of chlamydia are crucial to prevent the progression to PID and its associated complications. In contrast, uterine fibroids, fibrocystic breast changes, and premenstrual syndrome are not direct consequences of chlamydia infection. While these conditions are relevant to women's health, they do not stem from untreated chlamydia like pelvic inflammatory disease does. Therefore, understanding the severe implications of untreated chlamydia emphasizes the importance of screening and treatment for this common infection.

SAMPLE

5. What condition could result in the need for a myomectomy?

A. Cervical cancer

B. Uterine fibroids

C. Endometriosis

D. Pelvic inflammatory disease

A myomectomy is a surgical procedure specifically aimed at removing uterine fibroids, which are benign growths originating from the muscle layer of the uterus. Uterine fibroids, also referred to as leiomyomas, can cause a variety of symptoms including heavy menstrual bleeding, pelvic pain, and pressure symptoms depending on their size and location. When these fibroids lead to significant symptoms or complications, a myomectomy may be indicated to alleviate those issues while preserving the uterus. In contrast, conditions such as cervical cancer involve the removal of cancerous tissue from the cervix, which often necessitates different surgical procedures such as a hysterectomy or cone biopsy rather than a myomectomy. Endometriosis is managed through various means including hormonal therapy and may sometimes require surgical intervention, but it does not specifically involve myomectomy since it deals with endometrial-like tissue outside the uterus rather than fibroids. Pelvic inflammatory disease primarily involves infection of the female reproductive organs and would be managed differently, often with antibiotics and potential surgical intervention for complications, but not with a myomectomy. Therefore, the specific focus on the removal of uterine fibroids qualifies B as the correct answer.

6. What screening test is recommended for detecting cervical cancer?

A. Ultrasound

B. CT scan

C. Pap smear

D. MRI

The recommended screening test for detecting cervical cancer is the Pap smear. This test is specifically designed to collect cells from the cervix to check for precancerous changes and the presence of cervical cancer. The Pap smear is vital because it can identify abnormal cervical cells before they develop into cancer, allowing for earlier intervention and treatment, which significantly increases the chances of successful outcomes. In conjunction with the Pap smear, the HPV (human papillomavirus) test may also be performed to determine if high-risk types of HPV are present, as persistent infection with these viruses is a major risk factor for developing cervical cancer. Other screening methods, such as ultrasound, CT scans, and MRI, are not appropriate for routine cervical cancer screening, as they serve different diagnostic purposes and are not effective in detecting cellular changes in cervical tissue. Ultrasound is more commonly used for assessing structural abnormalities of the reproductive organs, CT scans are primarily for imaging internal organs in detail, and MRI is often used for evaluating soft tissue structures. Thus, the Pap smear remains the gold standard for cervical cancer screening.

SAMPLE

7. What is typically observed in a pregnancy exam at 28 weeks?

A. Fundus at the pubic symphysis

B. Fundus at the umbilicus

C. Fundus above the umbilicus

D. Fundus below the pubic symphysis

At around 28 weeks of pregnancy, it is typical for the fundus, which is the top part of the uterus, to be found above the level of the umbilicus. This corresponds to the time when the uterus is expanding significantly to accommodate the growing fetus. By the third trimester, the fundal height measurement is an important indicator of fetal growth and development. In a healthy pregnancy, around 28 weeks, the fundus is expected to reach approximately 2 to 3 centimeters above the umbilicus. This indicates that the pregnancy is progressing normally, as the uterus continues to rise and expand throughout the latter stages of gestation. In contrast, the other options do not align with typical findings at this stage of pregnancy. The fundus being at the pubic symphysis or at the level of the umbilicus would be more characteristic of earlier in the pregnancy, while the fundus below the pubic symphysis would indicate a potential issue, as the uterus would be much smaller than expected at this advanced gestational age. Thus, the observation of the fundus being above the umbilicus at 28 weeks is a key marker of proper fetal development and gestational progress.

8. Which of the following is a contraindication to prescribing oral hormonal contraceptives?

A. History of iron deficiency anemia

B. History of dysmenorrhea since age 15

C. Positive family history of ovarian cancer

D. History of smoking in a patient older than age 35

Oral hormonal contraceptives are commonly used for various purposes, including contraception and managing menstrual-related disorders. However, certain medical conditions and lifestyle factors can contraindicate their use due to increased risks of complications, especially thromboembolic events. In the case of a patient who is older than age 35 and has a history of smoking, the risk for cardiovascular issues, including blood clots, significantly increases. Smoking has a well-established relationship with cardiovascular disease and can enhance the thrombotic effects of hormonal contraceptives, creating a heightened risk of serious complications such as deep vein thrombosis (DVT) and pulmonary embolism. Therefore, this specific scenario represents a contraindication for prescribing oral hormonal contraceptives, as the potential risks outweigh the benefits. Other options listed, such as a history of iron deficiency anemia, dysmenorrhea, or a positive family history of ovarian cancer, do not generally pose the same level of risk regarding the use of oral contraceptives. Iron deficiency anemia, while it requires medical management, does not prevent the safe use of hormonal contraception. Similarly, a history of dysmenorrhea or a family history of ovarian cancer would not contraindicate the use of hormonal contraceptives, as these conditions are not

SAMPLE

9. What is the typical presentation of uterine fibroids?

A. Painful urination

B. Menorrhagia

C. Night sweats

D. Weight gain

Uterine fibroids, also known as leiomyomas, are non-cancerous growths of the uterus that often present with specific symptoms, one of the most common being menorrhagia, or heavy menstrual bleeding. Women with fibroids may experience increased menstrual flow or prolonged periods due to the fibroids' ability to disrupt normal uterine function and anatomy. The presence of these tumors can distort the endometrial lining and lead to excessive bleeding. In contrast, while painful urination can occur if fibroids exert pressure on the bladder, it is not a typical symptom. Night sweats and weight gain are not classic symptoms associated with uterine fibroids; these may be indicative of other health issues or hormonal imbalances rather than directly related to the presence of fibroids. Thus, menorrhagia stands out as the primary and most characteristic symptom linked to uterine fibroids.

10. Which of the following is a major risk factor for an ectopic pregnancy?

A. Alcohol intake

B. Young maternal age

C. History of salpingitis

D. Low dose oral contraceptive use

A major risk factor for ectopic pregnancy is a history of salpingitis, which refers to inflammation of the fallopian tubes. This condition can lead to scarring, adhesions, or blockage of the tubes, significantly increasing the likelihood of an embryo implanting outside of the uterine cavity. The damage caused by salpingitis can result from previous infections, particularly sexually transmitted infections, which may compromise the structural integrity of the fallopian tubes and impede the normal movement of the embryo toward the uterus. Other factors that could influence the risk of ectopic pregnancy include previous surgery on the fallopian tubes, certain infertility treatments, or pelvic inflammatory disease. However, among the options provided, a history of salpingitis stands out as a definitive risk factor due to its direct pathological impact on the fallopian tubes, which play a critical role in the normal implantation of a fertilized egg.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://womenshealtheor.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE